

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/





HARVARD COLLEGE LIBRARY




GIFT OF

JAMES STURGIS PRAY


Charles Eliot Professor of Landscape Architecture


SAN

To be kept in the main collection of the College Library



, ` . •





U. S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

OF

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,

BY

SAMUEL HENSHAW.

Parts I, II, and III.

THE MORE IMPORTANT WRITINGS

OF

BENJAMIN DANN WALSH

AND

CHARLES VALENTINE RILEY.

WASHINGTON:

GOVERNMENT PRINTING OFFICE.

1890.

č

BIBLIOGRAPHY

OF

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE, BY $^{\bullet}$

SAMUEL HENSHAW.

PART I.

THE MORE IMPORTANT WRITINGS

of

BENJAMIN DANN WALSH.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1889.

Z 1,889

FARVARD COLLEGE LIBRARY
GIFT OF

JAMES STURGIS PRAY

15 Sep. 1922

S 7510.89

V

FORTY-SEVENTH CONGRESS, FIRST SESSION.

CONGRESS OF THE UNITED STATES,
IN THE HOUSE OF REPRESENTATIVES,

June 24, 1882.

Resolved, By the House of Representatives (the Senate concurring) that there be printed, for the use of the Department of Agriculture, one thousand copies of a special report, entitled "Bibliography of Economic Entomology."

Attest:

EWD. McPherson, Clerk.

Passed the Senate July 6, 1882.

F. E. SHOBER,
Acting Secretary.

A true copy.

EWD. McPherson, Clerk.

2

LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,
DIVISION OF ENTOMOLOGY,
Washington, D. C., June 12, 1889.

SIR: In the absence of Professor Riley, who is now at the Paris Exposition in official capacity, I beg to submit for publication Parts I, II, and III of the Bibliography of American Economic Entomology, which was ordered by Congress in July, 1882. Owing to various unforeseen delays this work has not yet been completed. Its preparation was originally assigned by Professor Riley to Mr. B. Pickman Mann, formerly of this Division, and upon the severance of his connection with the Department, in the fall of 1885, it was turned over in an incomplete and fragmentary condition to Mr. Samuel Henshaw, of Boston, who has added greatly to it and practically rewritten the portions already prepared.

The extent of the work renders it advisable to publish it in several parts, and as there is unquestionably a more urgent need on the part of working economic entomologists for a bibliography of the writings of B. D. Walsh and C. V. Riley than of others, on account of their great number and value, these portions have been finished first. Their immediate publication will probably antecede the publication of the final part only by a few months.

Very respectfully, yours,

L. O. HOWARD,

Acting Entomologist.

Hon. J. M. Rusk, Secretary of Agriculture.





TABLE OF CONTENTS TO PARTS I, II, AND III.

	Page.
Introduction	7
The writings of B. D. Walsh (1860-1873)	9
The joint writings of B. D. Walsh and C. V. Riley (1868-1869)	51
The writings of C. V. Riley (1863–1888)	97
Systematic index of the new names:	
By B. D. Walsh and by Walsh and Riley	373
By C. V. Riley	377
Index	381
ĸ	



INTRODUCTION TO PARTS I, II, AND III.

The object of the present lists is to record the principal entomological writings of B. D. Walsh and C. V. Riley. Few will be likely to criticize the association of the names of Walsh and Riley in a list of this kind, so largely devoted to economic entomology, or to question the desirability of collating their writings in advance of others.

As recognized authorities upon entomology, especially from an economic standpoint, their writings have been extensively quoted sometimes with, but frequently without, credit in the agricultural and horticultural journals of the day. These secondary references, though of a certain value when the original is inaccessible, have, as a rule, been omitted. In citing such as are given I have used my personal judgment. To have given all would have increased the size of the list considerably with but a slight increase in its usefulness.


Occasionally, a publisher to meet some popular demand puts the writings of an author into book form without the knowledge of the latter. Such an occurrence has happened more than once with Dr. Riley's writings. To mention but one: In 1877 Messrs. George Rutledge & Sons, of London, published "The Colorado beetle, with suggestions for its repression and methods of destruction," with Dr. Riley's name upon the title-page as author. This is a partial reprint of Potato Pests entirely unsanctioned and published without knowledge of the author. (See London Times, October 17, 1887.) These publishers' reprints are omitted from this list. The "S.-b." after the reference of many of the articles from the agricultural and other journals refers to the series of scrap-books used in divisional work.

A biographical sketch of Dr. Walsh is given in the second volume of the American Entomologist, Vol. II, No. 3, pp. 65-68, January, 1870.

The chief facts in Dr. Riley's life may be found in the "Common-wealth of Missouri" (1875) and the National Farmer, September 20, 1883.

'In the preparation of the list I have freely used all data accumulated by the Division, notably the work of Mr. B. P. Mann and a private list of Dr. Riley's.

Many persons, both within and without the Department, have kindly answered my inquiries.—S. H.



BIBLIOGRAPHY OF THE MORE IMPORTANT CONTRIBUTIONS TO AMERICAN ECONOMIC ENTOMOLOGY.

THE WRITINGS OF B. D. WALSH (1860-1873).

- 1. Walsh, B. D. Bark-lice. < Western Rural. S.-b. No. 3, p. 11. Habite of Lecanium compared with Aspidiotus [=Mytilaspis].
- Walsh, B. D. Entomological notes. <Prairie Farmer, 17 May, 1860, [v. 21], n. s., v. 5, pp. 308-309, figs. S.-b. No. 1, pp. 42-43.
 - Notice of Coccus harrisii n. sp. [=Chionaspis furfurus]; figure of its scales; description and figure of Capsus oblineatus [=Lygus pratensis] infesting apple, quince, and pear trees; figure of larva case and image and description of Phycita nebulo [=Acrobasis indiginella]; figures of image and of injuries of Amphicerus bicaudatus; habits of the same and of Scolytus [=Xyleborus] pyri, Saperda bivittata [=candida] and Chrysobothris femorata; means against the last.
- 3. Walsh, B. D. "Cutworms." < Prairie Farmer, 9 August, 1860, [v. 22], n. s., v. 6, pp. 82-83, fig. S.-b. No. 1, pp. 84-85.
 - Account of the larvæ of Lachnosterna quercina [=fusca] of Elateridæ of Agrotidæ and of Pangus [=Harpalus] culiginosus; figures of larvæ and images of the same or allied species; also of Heteronychus [=Ligyrus] relictus, Lopha [=Bembidium] 4-maculatum and Agonoderus pallipes.
- 4. Walsh, B. D. [Ips quadrisignatus, Say.] < Prairie Farmer, 1860, [v. 22], n. s., v. 6, p. —.
 - Ips 4-signatus [= fasciatus] attacks growing ears of sweet corn.
- 5. Walsh, B. D.] "Insect Life." < Proc. Agric. Conv.; Ann. meeting Ill. Nat. Hist. Soc.; Comm. Exerc. State Normal Univ. [Bloomington, Ill.], 1860, pp. 11-12.
 - Report, compiled from Chicago papers of address delivered before the Illinois Natural History Society; comparison between injurious insects and an invading army; necessity of appropriations for carrying on entomological work; economic entomology in Europe; importance of beneficial insects.

ţ

BIBLIOGRAPHY

OF

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,
BY

SAMUEL HENSHAW.

PART I.

THE MORE IMPORTANT WRITINGS

OF

BENJAMIN DANN WALSH.

WASHINGTON: GOVERNMENT PRINTING OFFICE. 1889.

- 24. WALSH, B. D. List of the Pseudoneuroptera of Illinois contained in the cabinet of the writer, with descriptions of over forty new species, and notes on their structural affinities. <Proc. Acad Nat. Sci. Phil., September, 1862, pp. 361-402.
 - One hundred and nine (109) species are enumerated; two (2) new genera and forty (40) new species are described. For a list of the same see the Systematic Index.
- 25. WALSH, B. D. [Erythroneura tricincta Fitch.] < St. Louis Valley Farmer, October, 1862, pp. 305-306, fig.

 Erythroneura [=Typhlocyba] tricincta on grape-vines.
- 26. WALSH, B. D. Grasshoppers and locusts. < Journ. Ill. State Agric. Soc., November, 1862, pp. 1-3. Reprint: < Trans. Ill. State Agric. Soc., v. 5, pp. 497-499.
 - Injuries of locusts in the Mississippi Valley; differences between locusts and Cicadas.
- 27. WALSH, B. D. On the genera of Aphidæ found in the United States. < Proc. Ent. Soc. Phil., December, 1862, v. 1, pp. 294-311, figs. 1-8.
 - Synoptic separation of the genera; list of seventy (70) species found in the United States, with food habitat; Aphis [= Nectarophora] rudbeckiæ Fitch, A. vitis ? Scop., A. maidis ? Fitch, Lachnus caryæ Harris, Thelaxes [= Colopha] ulmicola Fitch are described; one (1) new genus and eleven (11) new species are described; see the Systematic Index for a list of the same; the figures show venation and other details. Two (2) new ants, Formica aphidicola and F. [= Lasius] latipes, are described.
- 28. WALSH, B. D. [Observations on Papilio glaucus and P. turnus.] < Proc. Ent. Soc. Phil., February, 1863, v. 1, pp. 349-352.

 Reasons for considering turnus and glaucus identical; the latter a dimorphic
 - Q of the former; distribution of the two forms; citation of similar case among Pierida and of analogous one among Dytiscida; description of larva from which a black Q was raised.
- 29. WALSH, B. D. [Katydid eggs.] < Prairie Farmer, 28 February, 1863, [v. 27], n. s., v. 11, p. 132, fig.
 Oviposition of katydid eggs on apple-twigs.
- 30. Walsh, B. D. Fire-blight. < Prairie Farmer, 4 April, 1863, [v. 27], n. s., v. 11, p. 212, fig. S.-b., No. 1, p. 10.
 - Description and figures of eggs of Chloroneura malefica [= Empoasca viride-scens] and the slits in which they are deposited in apple-twigs; cause of fire-blight; suggestion of remedies.
- 31. WALSH, B. D. Insects injurious to fruit trees. < Prairie Farmer, 2 May, 1863, [v. 27], n. s., v. 11, p. 276, fig. S.-b., No. 1, p. 9.
 - Fruit and foliage of pear-trees destroyed by Capsus oblineatus [=Lygus pratensis]; description and figure of imago; habits, food-plants, and remedies.
- 32. WALSH, B. D. The ten-striped spearman. < Prairie Farmer, 6 June, 1863, [v. 27], n. s., v. 11, p. 356, fig. S.-b., No. 1, p. 7.
 - History, ravages, and means against Doryphora 10-lineata; figure of imago; mentions other injurious Phytophaga.

- 33. WALSH, B. D. The plum-gouger; a new foe of the plum. < Prairie Farmer, 13 June, 1863, [v. 27], n. s., v. 11, pp. 372–373, figs. 1–3. S.-b., No. 1, p. 6.
 - Habits, description, and figures of Anthonomus? prunicida n. sp. [= Coccotorus scutellaris], injurious to plums; comparison with Conotrachelus nonuphar; figure of the latter.
- 34. WALSH, B. D. The plum-gouger. < Prairie Farmer, 11 July, 1863, [v. 28], n. s., v. 12, p. 21, 2 figs. S.-b., No. 1, p. 10.
 - Distribution and ravages of Anthonomus prunicida [= Coccotorus scutellaris]; list of insects found on plum-trees; description of Conotrachelus puncticollis n. sp. [= C. geminatus]; usefulness of Reduvius raptatorius [= Sinea diadema]; figures A. prunicida and R. raptatorius.
- 35. Walsh, B. D. A new fruit foe. < Prairie Farmer, 18 July, 1863, [v. 28], n. s., v. 12, p. 37, fig. S.-b., No. 1., p. 11.
 - Description and figure of imago of *Epicarus imbricatus*; injuries to fruit trees and gooseberry-bushes; supposed habits of this and allied species; hibernation of the larva of *Conotrachelus*; description of *C. cratægi* n. sp.
- 36. Walsh, B. D. Leaf-hopper. < Prairie Farmer, 25 July, 1863, [v. 28], n. s., v. 12, p. 53.
 - Proconia [= Oncometopia] undata injurious to grape-vines in southern Illinois; its eggs laid in twigs.
- 37. Walsh, B. D. Locust borers. < Prairie Farmer, 15 August, 1863, [v. 28], n. s., v. 12, p. 101. S.-b., No. 1, p. 12.
 - Description and habits of Clytus [= Cyllene] robiniæ; Clytus of Carya does not attack Robinia; Cossus robiniæ mentioned; larvæ and imagos of Nitidulidæ found under decaying bark.
- 38. Walsh, B. D. Insect friends and insect foes. The twice-stabbed lady-bird. < Prairie Farmer, 1863, [v. 28], n. s., v. 12; 22 August, p. 117, figs. 1, 2; 29 August, p. 133, figs. 3, 4. S.-b., No. 1, pp. 11-12.
 - Description and figure of imago of Chilocorus bivulnerus; figure of larva; destruction of injurious insects by predaceous insects; brief description of and means against Lytta cinerea [= Macrobasis unicolor], found on potato, English bean, and apple, and L. [= Epicauta] vittata, on potato; figures Lytta atrata [= Epicauta pennsylvanica], found on aster and potato, and larva-cases of Solenobia sp. from under bark of apple-trees; note on allies of Solenobia.
- 39. WALSH, B. D. Observations on certain N. A. Neuroptera, by H. Hagen, M. D., of Kænigsberg, Prussia; translated from the original French MS., and published by permission of the author, with notes and descriptions of about twenty new N. A. species of Pseudoneuroptera. < Proc. Ent. Soc. Phil., October, 1863, v. 2, pp. 167-272, fig.</p>
 - Dr. Hagen's comments relate to the species described in No. 24; the author's notes supplement these and describe three (3) new genera and twenty-three (23) new species; for a list of these, see the Systematic Index; descriptions of previously described species, with notes on their habits, distribution, etc., are also given; Batis interlineata [= Siphlurus femoratus] is proposed

- 39. WALSH, B. D.—Continued.
 - (p. 190) in case B. [= S.] femorata Walsh is distinct from B. [= S.] femorata Say; in a note (p. 255) Ophiogomphus mainensis n. sp. is described from Packard's manuscript; the venation of the Odonata and the terminal abdominal characters of Stalis infunata are figured.
- 40. Walsh, B. D. On certain remarkable or exceptional larvæ, coleopterous, lepidopterous, and dipterous, with descriptions of several new genera and species, and of several species injurious to vegetation, which have been already published in agricultural journals. < Proc. Bost. Soc. Nat. Hist., February, 1864, v. 9, pp. 286-318.
 - Habits of the larva of Cotalpa lanigera, Pelidnota punctata, Cratonychus [= Melanotus] incertus, and Xyloryctes satyrus; comparative characters of the larva and imago of Halisidota antiphola n. sp. [= tessellaris]; description of larva, pupa, and imago of Sphingicampa (n. g.) distigma n. sp. [= bicolor]; characters and affinities of Dryocampa [= Sphingicampa] bicolor?; description of imagos of Limacodes scapha and L.? [= Phobetron] hyalinus n. sp., and of larva of the last and L.? [= Phobetron?] tetradactylus n. sp.; description of the larva, pupa, and imago of Hipparchiscus n. g. venustus n. sp. [= Aplodes mimosaria]; habits and description of the larva of an undetermined Tabanus, description of the pupa; description of the larva, pupa, and imago of Midas fulvipes n. sp., habits of its larva. On pp. 309-318 descriptions of several new species, with brief notes of food-habits, etc., are reprinted from various agricultural journals.
- 41. WALSH, B. D. On dimorphism in the hymenopterous genus Cynips; with an appendix containing hints for a new classification of Cynipidæ and a list of Cynipidæ, including descriptions of several new species, inhabiting the oak-galls of Illinois. < Proc. Ent. Soc. Phil., March, 1864, v. 2, pp. 443-500, fig.
 - Detailed observations proving the frequent occurrence of dimorphism in the Cynipida; Cynips [= Amphibolips] aciculata, a dimorphic form of C. q. spongifica; concludes from analogy "that aciculata Q generates galls which produce by parthenogenesis & spongifica exclusively, and that Q Q spongifica coupling in June with these & & oviposit in the same month in the young buds of the oak, the eggs lying dormant till the following spring, when some of the eggs produce Q spongifica in June, and some Q aciculata in the autumn or early in the following spring, which last in their turn, as before mentioned, generate & spongifica to appear in the following June;" interesting details concerning the history of the group are given; relations between the true gall-flies (Psenides) and parasitic Cynipides (Inquilines); classification and characters of the same; anatomical structure and homologies of the family; the list includes fifteen (15) species inhabiting the various oaks, with descriptions of their galls and of several new species; seven (7) species of Inquilina are described; for the new species, see the Systematic Index; the figures illustrate the anatomy of the abdomen and ovipositor.
- 42. Walsh, B. D. The four-humped Curculio. A new foe of the apple. <Valley Farmer. Reprint: <Prairie Farmer, 27 August, 1864, [v. 30], n. s., v. 14, p. 131, 2 figs. S.-b., No. 1, p. 28.
 - Description and figures of Anthonomus quadrigibbus, a fee of the apple; comparison with Constrachelus nenuphar and Anthonomus prunicida [= Cocoptorus scutellaris].

43. Walsh, B. D. On the pupa of the ephemerinous genus Batisca Walsh. < Proc. Ent. Soc. Phil., August, 1864, v. 3, pp. 200-206, fig.

Description and figures of the pupa of Batisca obesa Say.

ì

- 44. Walsh, B. D. On certain entomological speculations of the New England school of naturalists. < Proc. Ent. Soc. Phil., August-September, 1864, v. 3, pp. 207-249.
 - Discussion of statements, chiefly by Prof. L. Agassiz, upon questions in general entomology. The memoir is divided into sections; in the first, the distribution of insects in North America is discussed in opposition to the views of Agassiz; in the second section the Darwinian theory of the origin of species is discussed and some remarks on the variation of several species are added; in the third and fourth sections the statements of Prof. Agassiz regarding the worm-like larva stage of insects and the resemblances between the pupæ of insects and the Crustacea are discussed and criticised; in the fifth section the author argues that the relative rank of the different orders of insects must be determined from a consideration of all the characters of each order and not from the significance of any one character; the sixth section is a review of Dana's classification of insects on the principle of cephalization; and in the seventh section several erroneous statements and generalizations in the same author's paper on classification are pointed out.
- 45. Walsh, B.D. On phytophagic varieties and phytophagic species. < Proc. Ent. Soc. Phil., November, 1864, v. 3, pp. 403-430.
 - Many species feed exclusively upon a single food-plant, while other species feed upon many kinds; correlated with this are certain larval or imaginal differences; when these different forms interbreed they are considered the same species and are termed phytophagic varieties; these when separated with a single kind of food-plant form phytophagic races, and these again by a continuation of their isolation form phytophagic species. Phytophagous forms are most abundant where the image is wingless. Diapheromera velii, Tingts [= Gargaphia] tilia, and T. [=G.] amorpha are described as new; the following are mentioned in the remarks upon species and varieties: Haltica [= Disonycha] alternata, Chrysomela scalaris, Clytus [= Cyllene] robinia, C. [= C.] pictus, Dryocampa [= Sphingicampa] bicolor, Sphingicampa distigma [= bicolor], Bombyx [= Sericaria] mori, Halisidota tessellaris, H carya, H. antiphola [= tessellaris], Cynips [= Amphibolips] q. spongifica, C. [= Andrious] q. punotata, C. [= A.] q. podagra, and C. [= Amphibolips] q. inanis.
- 46. WALSH, B. D. On the insects, coleopterous, hymenopterous, and dipterous, inhabiting the galls of certain species of willow. Part 1st—Diptera. <Proc. Ent. Soc. Phil., December, 1864, v. 3, pp. 543-644.
 - Structural characters, habits, metamorphoses, etc., of *Cecidomyidæ*; remarks on the unity of habits in genera; synopsis of cecidomyidous galls of *Salix*; descriptions of new species, their galls and inquilines; for a list of the new species, see the *Systematic Index*. See No. 197 for Part 2d.
- 47. WALSH, B. D. The borer. <Colman's Rural World. Reprint; <Prairie Farmer, 6 May, 1865, v. 15, p. 355.

Notice of soap as a means against borers.

- 48. Walsh, B. D. The new potato-bug and its natural history. < Pract. Ent., 30 October, 1865, v. 1, pp. 1-4.
 - Record of the spread eastwards of *Doryphora* 10-lineata; description of the egg; food-plants, enemies, and means against the same; comparative characters of *D. juncta* and *D.* 10-lineata; species of *Meloida* injurious to the potato.
- 49. [Walsh, B. D.] The joint-worm. < Pract. Ent., 27 November, 1865, v. 1, pp. 11, 12.
 - Reasons for disbelieving that the joint-worm is the larva of one of the Chalcidida.
- 50. Walsh, B. D. On phytophagic varieties and phytophagic species, with remarks on the unity of coloration in insects. <Proc. Ent. Soc. Phil., November-December, 1865, v. 5, pp. 194-216, fig. See: <Amer. Journ. Sci., September, 1865, ser. 2, v. 40, pp. 282-284. <Ann. and Mag. Nat. Hist., November, 1865, ser. 3, v. 16, pp. 383, 384.
 - Argues for the origin of races and species by phytophagic isolation; refers to Datana ministra, D. contracta, Halisidota tessellaris, H. antiphola [= H, tessellaris], H. harrisii [= H. tessellaris], Arhopalus [= Cyllene] pictus, A. [= C.] robinia, A. infaustus [= C. decorus], Callidium antennatum, C. janthinum, Conotrachelus nenuphar, Doryphora 10-lineata, and D. juncta; unity of coloration explained by assumption of a genetic connection between various species; figure showing bullse on wing of Ichneumon.
- 51. W[ALSH], B. D. [Cicada years.] < Pract. Ent., 25 December, 1865, v. 1, pp. 18-19.
 - Answer to inquiry of M. S. Hill; Cicada districts of the United States, as given by Fitch in N. Y. Rept. I, p. 39; habits of C. [= Tibicen] septendecim.
- 52. Walsh, B. D. Insects injurious to vegetation in Illinois. <Trans. Ill. State Agric. Soc., 1865, v. 5, pp. 469-483, figs.
 - Notes on injurious insects an Illinois in 1861; life-history and ravages of Leucania unipuncta.
- 53. Walsh, B. D. The "Thrips" of the vine-growers. What is it? <Pract. Ent., 25 December, 1865, v. 1, pp. 20-21.
 - The Thrips of fruit-growers probably not a true Thrips; food-habits of Thrips.
- 54. WALSH, B. D. The grub-worm. <Colman's Rural World, December, 1865. S.-b., No. 1, p. 51.
 - Habits, ravages, and means against Lachnosterna quercina [=fusca]; its larva compared with that of Ligyrus relictus; vernacular names of the Lacknosterna and of Allorhina nitida.
- 55. Walsh, B. D. Borers. < Pract. Ent., 29 January, 1866, v. 1, pp. 25-31, figs. 1-8.
 - Remarks on the name "borer;" figures, natural history, and mean's against Saperda bivittata [= candida], Chrysobothris femorata, Bostrichus [= Amphicerus] bicaudatus, Ægeria [= Sannina] exitiosa, A. tipuliformis, Clytus [= Cyllene] robinia, C. [= C.] pictus, and Cerasphorus [= Chion] cinctus; need of State aid against injurious insects.

- 56. [Walsh, B. D.] [Attagenus? injuring feathers.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of R. Parnell; ravages of and means against Dermestides.
- 57. [Walsh, B. D.] [Codling moth.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of Isaac Hicks; natural history and means against Carpocapsa pomonella.
- 58. [Walsh, B. D.] [Aphididæ.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of J. H. Foster, jr.; comparative characters of Eriosoma [= Schizoneura] lanigera and Pemphigue pyri [= Schizoneura lanigera].
- 59. [Walsh, B. D.] [Mites mistaken for plant-lice.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.
 - Answer to inquiry of W. M. Smith; distinctive characters of Insecta and Arachnida; description and habitat of the eggs of Aphie mali.
- 60. [Walsh, B. D.] [Saddle-back caterpillar.] < Pract. Ent., 26 February, 1866, v. 1, p. 34.</p>
 Answer to inquiry of S. P. M.; description of the larva and image of Empretia stimulea; urticating properties of the larva.
- 61. W[ALSH], B. D. [Red-humped prominent.] < Pract. Ent., 26 February, 1866, v. 1, p. 35.</p>
 Answer to inquiry of E. D. Wright; life-habits of Notodonta [= Œdomasia]
 - concinna.
- WALSH, B. D. The joint-worm. < Pract. Ent., 26 February, 1866,
 v. 1, pp. 37-38.
 Habitats of the larva of Cocidomyia destructor and of Eurytoma sp.; criticism of A. Fitch's report upon the latter.
- 63. W[ALSH, B. D.]. Entomology all a humbug. <Pract. Ent., 26 February, 1866, v. 1, p. 39.</p>
 Fewness of entomologists and the multiplicity of their duties the reason why
- so little has been effected against noxious insects.

 64. W[ALSH], B. D. [Cocoons of Ichneumon-flies.] < Pract. Ent., 26
 March, 1866, v. 1, p. 46.
 - Answer to inquiry of M. S. Hill; life-habits of the Microgasters.
- 65. W[ALSH], B. D. [Mass of eggs.] < Pract. Ent., 26 March, 1866, v. 1, p. 46.
 - Answer to inquiry of T. R. Payne; characters and life-habits of Orgyia loucostigma.
- 66. [WALSH, B. D.] [Apple-tree caterpillar; parasites.] < Pract. Ent., 26 March, 1866, v. 1, p. 46.
 - Answer to inquiries of G. E. Brackett; food-plants of Clisiocampa americana; characters and habits of Tachinida; habits of secondary parasites.
- 67. W[ALSH], B. D. Apple-tree borers. < Pract. Ent., 26 March, 1866, v. 1, p. 47.
 - Reprint, with remarks, of "Apple-tree borers" (Journ. of Commerce); extract from A. Fitch, of larval habits of Saperda bivittata [= candida].

 2 ENT

- 68. Walsh, B. D. The white grub. <Pract. Ent., 30 April, 1866, v. 1, pp. 60-62.
 - Descriptions and habits of the larval states of Lachnosterna queroins [=fueca] and Ligyrus relictus; ravages and food-habits of the former; the hog as a white-grub destroyer.
- 69. Walsh, B. D. [Apple-tree scales.] < Pract. Ent., 30 April, 1866, v. 1, p. 64.
 - Answer to inquiry of L. S. Pennington; descriptions and ravages of Mytilaspis pomorum and Chionaspis furfurus; Coccinellidæ as a means against the former; petroleum as a remedy against bark-lice.
- 70. WALSH, B. D. [Clothes-moths.] < Pract. Ent., 30 April 1866, v. 1, p. 64.
 - Answer to inquiry of S. Canby; habits and means against Tineida.
- 71. WALSH, B. D. [Poplar borer.] < Pract. Ent., 30 April, 1866, v. 1, p. 64.
 - Answer to inquiry of P. C. Truman; larva and image of Saperda calcurata infesting cottonwood.
- 72. WALSH, B. D. [Worms in flour and rye.] < Pract. Ent., 30 April, 1866, v. 1, p. 64.
 - Answer to inquiry of L. D. Hunt; descriptions and habits of Tenebrio mollior and Dermestes lardarius.
- 73. Walsh, B. D. Popular remedies for noxious insects. < Pract. Ent., 28 May, 1866, v. 1, pp. 71-74.
 - Criticism of various worthless remedies against injurious insects.
- 74. WALSH, B. D. [Serica iricolor Say.] < Pract. Ent., 28 May, 1866, v. 1, p. 77.
 - Answer to inquiry of B. F. Seibert; description, ravages, and habitat of Serioa iricolor.
- 75. Walsh, B. D. [Measuring-worms.] < Pract. Ent., 28 May, 1866, v. 1, p. 77.
 - Answer to inquiry of T. Siveter; habits of Ennomos magnaria [=alnisria]; food-plants, habits, and means against E. [=Eudalimia] subsignaria and Anisopteryx [=Paleacrita] vernata.
- 76. WALSH, B. D. [Lygus pratensis.] < Pract. Ent., 25 May, 1866, v. 1, pp. 77-78.
 - Answer to inquiry of H. B. Howarth; habits and odors of Lygus pratencie; changes insects pass through.
- 77. WALSH, B. D. [Apple-tree insects.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.
 - Answer to inquiry of C. Cooke; an unnamed tineid moth raised from cocoons off twigs of apple-tree; *Hemiteles* sp. and *Microgaster* sp. bred from cocoons attached to eggs of *Clisiocampa americana*.
- 78. Walsh, B. D. [Destructive current-worm.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.
 - Answer to inquiry of W. M. Smith; cocoon and sexual characters of Nematus ribis [= ribesii].

- 78. WALSH, B. D. [Egg slits made by Homoptera.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.
 - Answer to inquiry of C. Dadant; slits in pear-twigs caused by Chloroneura malefica [= Empoases viridescens]; Proconia [= Oncometopia] undata oviposits in grape-vines.
- 80. WALSH, B. D. [Plant-lice.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.
 - Answer to inquiry of J. Flournoy; characters, habits, enemies, and means against Aphididæ.
- 81. WALSH, B. D. [Insects injuring plum-trees.] < Pract. Ent., 28 May, 1866, v. 1, p. 78.
 - Answer to inquiry of A. C. Hammond; *Egeria* [=Sannina] exitiosa and an elaterid larva attacking plum-trees; the latter probably attracted by decayed matter, the work of the peach borer.
- 82. WALSH, B. D. Prof. Dana and his entomological speculations. <Proc. Ent. Soc. Phil., May-June, 1866, v. 6, pp. 116-121.
 - Rejoinder to J. D. Dana's reply to criticism upon the classification of insects.
- 83. WALSH, B. D. Clover-worms. < Pract. Ent., 25 June, 1866, v. 1, pp. 82-83.
 - Summary of published observations concerning Asopia costalis; points which need explanation; probable remedy.
- 84. WALSH, B. D. Click-beetles. < Pract. Ent., 25 June, 1866, v. 1, p. 83.
 - Note to W.'s "Click-beetles;" snapping of Elateridæ; Jaeger's Life of N. Am. Insects and Emmons' N. Y. Report unreliable.
- 85. Walsh, B. D. Cut-worms. < Pract. Ent., 25 June, 1866, v. 1, pp. 85-86.
 - Definition, food-habits, and means against cut-worms; climbing habits; cut-worms in California. See Nos. 229, 281.
- 86. WALSH, B.D. The canker-worm. Finding a mare's nest. < Pract. Ent., 25 June, 1866, v. 1, p. 87.
 - Criticism of communication in Western Rural; improbability of the report that Anisopteryx [= Palescrita] vernata deposite her eggs on the ground.
- 87. WALSH, B. D. Driving nails into fruit trees. < Pract. Ent., 25 June, 1866, v. 1, pp. 87-88.
 - Uselessness of nails as a means against borers; extracts from various sources, with comments.
- 88. Walsh, B. D. The new potato-bug. <Pract. Ent., 25 June, 1866, v. 1, pp. 88-89.
 - Records Doryphora 10-lineata at Bloomington and Athens, Ill.; rate of its eastern progress; need of legislative action against injurious insects.
- 89. WALSH, B. D. [The painted borer.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.
 - Answer to inquiry of T. C. Wright; sexual difference in the antenne of Clytus [= Cyllone] pictus bred from hickory.

90. Walsh, B. D. [Bug allied to the chinch-bug.] < Pract. Ent., 25
June, 1866, v. 1, p. 89.

Answer to inquiry of H. B. Howarth; comparative characters of Blissus leucopterus and an unnamed lygacid of similar habits.

- 91. Walsh, B. D. [Canker-worms.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.
 - Answer to inquiry of C. H. W. Wood; means against Anisopteryx.
 - 92. Walsh, B. D. [Apple-tree tineid.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.

Answer to inquiry of C. Cooke; food-plant of Bucculatrix pomifoliella.

- 93. WALSH, B. D. [Spittle insect.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.
 - Answer to inquiry of M. V. B. Hathaway; life-habits of species of Aphrophora.
- 94. WALSH, B. D. [Insects named.] < Pract. Ent., 25 June, 1866, v. 1, p. 89.
 - Answer to inquiry of E. Hall; occurrence of *Doryphora* 10-lineata at Athens, Ill.; *Hippodamia* [= *Megilla*] maculata devours the eggs of the same; two species of *Cassida* affect the sweet-potato.
- 95. WALSH, B. D. [Longicorn borers.] < Pract. Ent., 25 June, 1866, v. 1, p. 90.
 - Answer to inquiry of Dr. Trimble; food-habits of Orthosoma cylindricum, [=brunneum], Prionus laticollis, and P. imbricornis.
- 96. WALSH, B. D. [Canker-worms.] < Pract. Ent., 25 June, 1866, v. 1, p. 90.
 - Answer to inquiry of F. K. Phœnix; season of oviposition of Anisopteryx.
- 97. WALSH, B. D. [White-pine scale.] < Pract. Ent., 25 June, 1866, v. 1, p. 90.

Answer to inquiry of H. Shimer; Aspidiotus [= Chionaspis] pinifolii infesting Pinus strobus.

- 98. WALSH, B. D. Doctors differ. < Pract. Ent., 30 July, 1866, v. 1, p. 96.
 - Criticism of communications in agricultural papers; tanners' oil injurious to trees; sulphur plugged in trees does not protect the leaves; sheep in an orchard do not keep away Conotrachelus nenuphar.
- 99. WALSH, B. D. The grain plant-louse. < Pract. Ent., 30 July, 1866, v. 1, pp. 96-97.

The insect attacking small grains in Georgia is probably a plant-louse.

100. Walsh, B. D. Popular names for insects. < Pract. Ent., 30 July, 1866, v. 1, p. 97.

The use of the same vernacular name for different insects causes uncertainty in identifying and in suggesting means against injurious insects.

101. Walsh, B. D. Scientific names. < Pract. Ent., 30 July, 1866, v. 1, pp. 97-99.

Explains the necessity of using Latin for scientific names,

- 102. Walsh, B. D. [Grape-vine insects.] < Pract. Ent., 30 July, 1866, v. 1, pp. 99-100.
 - Answer to inquiry of C. S. Jackson; habits of an undescribed *Fidia*; injuries to vines by plant-louse (*Aphis vitis t*); enemies of *Aphidida*.
- 103. Walsh, B. D. [Wire-worms.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.
 - Answer to inquiry of A. D. Chesebro; summer fallowing as a means against wire-worms.
- 104. Walsh, B. D. [Bark-lice.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.
 - Answer to inquiry of L. E. Harmon; life-history and means against Coccide.
- 105. WALSH, B. D. [Tortoise beetles.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.
 - Answer to inquiry of C. K. Yant; food-plants and larval habits of Cassidida.
- 106. Walsh, B. D. [Grape bark-louse.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.
 - Answer to inquiry of J. Bird; description and ravages of Lecanium [=Pul-vinaria] vitis.
- 107. Walsh, B. D. [Plum bark-louse.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.
 - Answer to inquiry of O, B. Douglas; description of Lecanium sp.
- 108. Walsh, B. D. [Saw-fly eggs.] < Pract. Ent., 30 July, 1866, v. 1, p. 100.
 - Answer to inquiry of L. D. Morse; position of saw-fly eggs on oak-leaf; characters of the larvæ of Tenthredinidæ.
- 109. Walsh, B. D. [Wheat midge.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.
 - Answer to inquiry of R. F.; ravages, description, habits, and means against Cocidomyia [=Diplosis] tritici.
- 110. WALSH, B. D. [Wheat midge.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.
 - Answer to inquiry of M. H. Boye; Cecidomyia [=Diplosis] tritici not infrequently passes from larval to imaginal state in the same season; means against the same.
- 111. Walsh, B. D. [Grape galls.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.
 - Answer to inquiry of J. H. Foster, jr.; description of the galls and larvæ of Cecidomyidæ.
- 112. Walsh. B. D. [Apple-tree web-caterpillar.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.
 - Answer to inquiry of Marion Hobart; description of the eggs of Clisiocampa americana; food-plants of the same and of Hyphantria textor [= cunea].

- 113. Walsh, B. D. [Grape-vine beetles.] < Pract. Ent., 30 July, 1866, v. 1, p. 101.
 - Answer to inquiry of M. S. Hill; food-habits of the larva and image of Pelidnota punctata and of Anomala lucicola.
- 114. WALSH, B. D. [Bot-flies.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.
 - Answer to inquiry of J. B. Fisher; habits and hosts of several species of Estridæ; parasite infesting Triplax [= Tritoma] thoracica. See No. 153.
- 115. Walsh, B. D. [Spruce-tree saw-flies.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.
 - Answer to inquiry of J. Barratt; description of cocoon and image of Lopky-rus abietis; means against the same. See Pract. Ent., v. 1, p. 114.
- . 116. WALSH, B. D. [Stings of insects.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.

 Answer to inquiry of A. O. Brickman; effects of the stings of bees and other
 - insects.

 117. WALSH, B. D. [Blackberry scale.] < Pract. Ent., 30 July, 1866,
 - v. 1, p. 102.

 Answer to inquiry of G. E. Brackett; characters of Lecanium sp. infesting blackberry.
 - 118. WALSH, B. D. [Eggs in sumach.] < Pract. Ent., 30 July, 1866, v. 1, p. 102.
 - Answer to inquiry of W. M. Smith; eggs of Orohelimum sp.? or Xiphidium sp.? in pith of sumach.
 - 119. W[ALSH], B. D. The "new potato-bug" in Maine. < Pract. Ent., 27 August, 1866, v. 1, p. 105.

 Extract from Maine Farmer, 26 July, 1866, with comments; the larva of
 - Lema trilineata mistaken for that of Doryphora 10-lineata.

 120. Walsh, B. D. [The army-worm.] < Pract. Ent., 27 August, 1866, v. 1, p. 107.
 - Extract from Western Rural, 21 July, 1866, with comments; army-worm mistaken for cauker-worm; need of precision in using pepular names.
 - 121. W[ALSH], B. D. [Fire-cure for potato-beetles.] < Pract. Ent., 27 August, 1866, v. 1, pp. 107-108.
 - Extract from letter from M. S. Hill, with comments; burning straw effective as a means against *Meloids* infesting potato-vines; its use against *Doryphora* 10-lineata; hot water as a means against the onion-maggot and peach-borer.
 - 122. Walsh, B. D. A new humbug. < Pract. Ent., 27 August, 1866, v. 1, pp. 108-110.
 - Exposure of proposed remedy against Cecidomyia destructor; life-habits of the same.
 - 123. W[ALSH], B. D. The striped bug. < Pract. Ent., 27 August, 1866, v. 1, p. 110, figs. 1, 2.
 - Extract from Western Rural, 21 July, 1866, with comments; characters, habits, means against, and figures of *Diabrotica vittata* and *D. 12-punctata*. See No. 148.

- 124. Walsh, B. D. [Grape-vine caterpillars.] < Pract. Ent., 27 August, 1866, v. 1, p. 111.
 - Answer to inquiry of B. Borden; description of the early stages of *Procris* [=Acoloithus] falsarius; characters and habits of the species of *Procris*.
- 125. Walsh, B. D. [Squash-vine insects.] < Pract. Ent., 27 August, 1866, v. 1, p. 111.
 - Answer to inquiry of J. Cope; description and habits of *Epilachna borealis*; habits of *Coocinellidæ*; description and means against the larva of *Trochilium cucurbitæ* [= Melittia ceto].
- 126. Walsh, B. D. [Injurious insects.] < Pract. Ent., 27 August, 1866, v. 1, p. 111.
 - Answer to inquiry of T. Conard; larval habits and means against Lema trilineata; work of the grain plant-louse (Nectarophora granaria) and of Trochilium oucurbitæ [= Melitia ceto]; mention of some of the enemies of bees.
- 127. WALSH, B. D. [Grape-leaf galls.] < Pract. Ent., 27 August, 1866, v. 1, pp. 111, 112.
 - Answer to inquiry of E. Daggy; description of the leaf-gall of *Phylloxera* vitifoliæ [= vastatrix]; reasons for including it in the Coccidæ; dipterous enemy of the same. See No. 154.
- 128. Walsh, B. D. [Grape-vine insects.] < Pract. Ent., 27 August, 1866, v. 1, p. 112.
 - Answer to inquiry of W. H. S.; description and means against an undescribed gall on the tendrils and leaf-stalks of a grape-vine; life-habits of parasitic *Chaloididæ*. See No. 160.
- 129. Walsh, B. D. [Grape-vine beetles.] < Pract. Ent., 27 August, 1866, v. 1, pp. 112, 113.
 - Answer to inquiry of W. J. Lawrence; food-habits of *Pelidnota punctata* and of *Colaspis flavida*.
- 130. Walsh, B. D. [Corydalis cornutus.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of J. S. Lewis; natural history and habits of Corydalis cornutus.
- 131. Walsh, B. D. [Tobacco-moth.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of F. W. Noble; transformations and food-plants of Sphinx [= Protoparce] carolina; food-plants of S. 5-maculata [= P. celeus].
- 132. Walsh, B. D. [Datana ministra.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of S. S. Lacy; food-plants and transformations of Datana ministra.
- 133. Walsh, B. D. [Grape-vine and fuschia beetles.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of F. K. Phonix; Colaspis flavida injurious to grape-vine shoots; Haltica exapta [= carinata] destructive to fuschia; means against flea-beetles.

- 134. WALSH, B. D. [Blister-beetles.] < Pract. Ent., 27 August, 1866, v. 1, p. 113.
 - Answer to inquiry of J. Barber; food-plants of Lytta marginata [= Epicauta cinerea], L. [= E.] vittata, L. atrata [= E. pennsylvanica], and L. cinerea [= Macrobasis unicolor].
- 135. Walsh, B.D. [Colaspis flavida; Lema trilineata.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.
 - Answer to inquiry of J. F. Benner; Colaspis flavida destructive to grape-vines; distribution and method of breeding Lema trilineata.
- 136. Walsh, B. D. [Green worms on gooseberries and currants; Lema trilineata.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.
 - Answer to inquiry of H. Burt; gooseberries and currents injured by larva of a saw-fly?; effects of excessive rain, heat, and drought; food-plants of Lema trilineata.
- 137. Walsh, B. D. [Galls.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.
 - Answer to inquiry of H. Shimer; note of galls made by Rhodites radioum, Trypeta solidaginis, and Thelaxes [= Colopha] ulmicola.
- 138. WALSH, B. D. [Potato and corn insects.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.
 - Answer to inquiry of T. H. Parsons; scarcity of Lema trilineata in Illinois; Doryphora 10-lineata destroyed by turkeys, but not by fowls; Gortyna sp.? injurious to young Indian corn; rules for breeding insects.
- 139. Walsh, B. D. [Cocoons on apple-trees.] < Pract. Ent., 27 August, 1866, v. 1, p. 114.
 - Answer to inquiry of M. S. Hill; description of cocoon and image of Attacus cocropia; the interaction of organisms.
- 140. Walsh, B. D. Imported insects; the gooseberry saw-fly. <Pract. Ent., 29 September, 1866, v. 1, pp. 117-125, fig.
 - List of some injurious European insects imported into America and of American insects imported into Europe; reasons for the increase and abundance of noxious European insects in America, and why American species do not flourish in Europe; natural history, description, and means against Nematus ventricosus [=ribesii]; figure of currant-leaf, showing method of deposition of eggs and work of larva; natural history and description of larva and imago of Pristiphora grossulariæ n. sp.; number of saw-flies infesting the gooseberry and currant in Europe.
- 141. WALSH, B. D. [Borers.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of W. H. R. Lykins; western distribution of Clytus [=Cyllene] robinia.
- 142. Walsh, B. D. [Wheat midge.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of C. P. Wickersham; destructiveness of *Cecidomyia* [= Diplosis] tritici; nature of its coccon.

- 143. Walsh, B. D. [Crickets.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of Marion Hobart; characters and food-habits of Acheta [=Gryllus] abbreviatus; katydids not exclusively vegetarians; Ecanthus niveus found feeding on plant-lice.
- 144. Walsh, B. D. [Walking-stick.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of L. W. Taylor; characters and habits of Spectrum [= Diapheromera] femorata.
- 145. Walsh, B. D. [Apple-tree insects.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of H. B. Beegle; characters and food-plants of *Limentitis* ursula; description and food-plants of the larva of *Lagoa opercularis*; characters of the imago.
- 146. Walsh, B. D. [Mites on grasshoppers.] < Pract. Ent., 29 September, 1866, v. 1, p. 126.
 - Answer to inquiry of W. Riddell; Astoma [= Trombidium] locustarum infesting Caloptenus femur-rubrum.
- 147. W[ALSH], B. D. Grasshoppers and locusts. < Pract. Ent., October, 1866, v. 2, pp. 1-5, 22.
 - Description, migration, and ravages of Caloptenus spretus; limits of its range and means of preventing its increase; confusion caused by the use of popular names of insects. On p. 22 a "note," accidentally omitted, gives the comparative length of the wing-covers and antennæ of Caloptenus spretus and C. femur-rubrum.
- 148. W[ALSH], B. D. The striped cucumber-bug. <Pract. Ent., October, 1866, v. 2, p. 5.
 - Diabrotica vittata attacks German asters. See No. 123.
- 149. W[ALSH], B. D. Fire blight. < Pract. Ent., October, 1866, v. 2, p. 7.
 - Extract from Horticulturist, with comments; cause of fire-blight unknown; not produced by Scolytus [= Xyleborus] pyri.
- 150. W[ALSH], B. D. The Hessian fly. < Pract. Ent., October, 1866, v. 2, p. 7.
 - Extract from Colman's Rural World, with comments; late sowing as a means against Cecidomyia destructor.
- 151. [Walsh, B. D.] [Injurious caterpillars; plant-lice enemies.] < Pract. Ent., October, 1866, v. 2, pp. 7-8.
 - Answer to inquiry of T. McGraw; habits, descriptions, and food-plants of larva of Dryocampa senatoria and of Datana ministra; food-habits of larva allied to D. ministra; description and habits of larva of Scymnus sp.; S. kamorrhous? bred from gall of Thelaxes [=Colopha] ulmicola.
- 152. [Walsh, B. D.] [Tomato-worm.] < Pract. Ent., October, 1866, v. 2, p. 8.
 - Answer to inquiry of A. A. Jackson; description of the larva and pups of Protoparce celeus; food-plants and harmlessness of the larva.

October, 1866, v. 2, p. 8. (Pract. Ent.,

Answer to inquiry of J. B. Fisher; larvæ found on swallows (see No. 114) belong to the Muscidæ and not to the Estridæ; extract from letter of Osten Sacken on larval habits of Muscidæ found with birds.

154. [Walsh, B. D.] [Bark-louse enemy.] < Pract. Ent., October, 1866, v. 2, p. 8.

Answer to inquiry of E. Daggy; see No. 127; according to Osten Sacken the bark-louse enemy belongs to Leucopis; systematic position of the genns.

155. [Walsh, B. D.] [Parasitized tomato-worm.] < Pract. Ent., October, 1866, v. 2, pp. 8-9.

Answer to inquiry of T. C. Wright; habits of Microgaster, parasitic on Protoparce celeus; habits of ichneumonized larvæ.

156. [Walsh, B. D.] [Beetles on buttercup and azalea.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of C. H. Peck; identification of Galeruca [=Adimonia] rufosanguinea, found on Ranunculus acris, and of Prasocuris varipes on Azalea nudiflora; synoptic table, by J. L. Leconte, of the species of Prasocuris.

157. [Walsh, B. D.] [Flying bug.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of E. E. Sheldon; Aphodius mistaken for Cocidomyia destructor.

158. [Walsh, B. D.] [Parasitic cocoon.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of H. W. Howarth; description of cocoons made by Microgaster and Pezomachus; habits of Ichneumonidæ.

159. [Walsh, B. D.] [Oak-bark louse.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of T. Meehan; undetermined Coccus on red-oak.

160. [Walsh, B. D.] [Grape-galls.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of W. H. S.; see No. 128; larva of *Leucopis* sp., an enemy to grape-vine gall-louse; probable identity of the maker of the tendril and leaf-galls; varieties of grape infested by *Phylloxera vastatrix*.

161. [WALSH, B. D.] ["Saddle-back."] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of T. M. Harvey; food-plants, description, and urticating properties of *Empretia stimulea*.

162. [Walsh, B. D.] [Corn-worm.] < Pract. Ent., October, 1866, v. 2, p. 9.

Answer to inquiry of J. B. Ellis; method of work of two lepidopterous larves.

163. [WALSH, B. D.] [Insects named.] < Pract. Ent., October, 1866, v. 2, p. 10.

Answer to inquiry of W. C. Fish; comparative characters of *Entilia sinuata* and *E.* [= *Publilia*] concava.

- 164. [Walsh, B. D.] [Grape insects.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of J. H. Garman; description and habits of larva of Procris [= Harrisina] americana; Proconia [= Oncometopia] undats and several species of Erythroneura [= Typhlocyba] as grape enemies.
- 165. [Walsh, B. D.] [Datana ministra.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of J. B.; increase of injuries caused by *Datana ministra*; distinctive characters of the larvæ of *Geometridæ*.
- 166. [Walsh, B. D.] [Iulus marginatus.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of T. Wiggins; characters and food of *Iulus* [= Spirobolus] marginatus.
- 167. [WALSH, B. D.] [Midge,] <Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of W. G. Morris; characters and habits of the larva and image of Chironomus sp.
- 168. [Walsh, B. D.] [Rat-tailed larva.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of J. A. Lapham; characters and habitats of the larvæ of rat-tailed Syrphidæ.
- 169. [Walsh, B. D.] [Beetle on thistle.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of W. S. Robertson; characters and food-plants of Phyllobrotica [= Diabrotica] longicornis.
- 170. [Walsh, B. D.] [Grape-vine Aphis.] < Pract. Ent., October, 1866, v. 2, p. 10.
 - Answer to inquiry of H.; grape-vine injured by Aphie vitie.
- 171. W[ALSH], B. D. The new potato-bug. <Pract. Ent., November, 1866, v. 2, pp. 13-16, fig.
 - Eastern progress of *Doryphora* 10-lineata; rate of travel; hand-picking the only effective remedy; description of a horse machine for their destruction; pecuniary loss inflicted by the species.
- 172. W[ALSH], B. D. The canker-worm < Pract. Ent., November, 1866, v. 2, p. 16.
 - Tarring trees effective against Paleacrita vernata.
- 173. W[ALSH], B. D. The canker-worm again. <Pract. Ent., November, 1866, v. 2, pp. 16-17.
 - Critical review of article in New England Farmer; value of tarring as a means against Paleacrita vernata.
- 174. Walsh, B. D. Notes by Benj. D. Walsh. < Pract. Ent., November, 1866, v. 2, pp. 19-20.
 - Remarks on statement of H. Shimer in his paper, "The grape leaf gall-coccus" (Pract. Ent., pp. 17-19); number of eggs to each 2; tendril and leaf-galls probably produced by the same insect; old leaves entirely free from bark-lice by the end of summer; characters of the pupa of *Leucopis*; systematic position and food-habits of *Thysanoptera*; criticism of paper by H. Shimer in Prairie Farmer, 3 November, 1861.

- 175. W[ALSH], B. D. The striped bug. < Pract. Ent., November, 1866, v. 2, p. 20.
 - Remarks on note by A.; effects of plaster as a means against Diabrotics vittats.
- 176. [Walsh, B. D.] [Currant insects; Psocus venosus.] < Pract. Ent., November, 1866, v. 2, p. 20.
 - Answer to inquiry of I. Hicks; occurrence of Pristiphora grossularia in the Eastern States; Ellopia [= Eufitchia] ribearia injurious to current; date of introduction of "sulphur-cure" on peach-trees; characters and habits of Procus venosus.
- 177. [Walsh, B. D.] [Worm in apple.] < Pract. Ent., November, 1866, v. 2, pp. 20-21.
 - Answer to inquiry of C. Ward; description and ravages of larva of Trypeta pomonella?
- 178. [Walsh, B. D.] [Insects attracted to light. | < Pract. Ent., November, 1866, v. 2, p. 21.
 - Answer to inquiry of Practical Pemologist; conditions favorable for attracting insects to light; kinds most frequently attracted; the proportion of injurious species attracted.
- 179. [Walsh, B. D.] [Zebra caterpillar.] < Pract. Ent., November, 1866, v. 2, p. 21.
 - Answer to inquiry of J. H. Parsons; description of the larva of *Mamestra picta*; times of transformation.
- 180. [Walsh, B. D.] [Inquiries answered.] < Pract. Ent., November, 1866, v. 2, p. 21.
 - Answer to inquiry of J. Pettit; identification of several beetles; galls on golden-rod made by Euryptychia saligneana and Trypeta solidaginis; characters of short-winged form of Micropus [=Blissus] leucopterus; its occurrence in Canada.
- 181. [Walsh, B. D.] [Worm in corn.] < Pract. Ent., November, 1866, v. 2, pp. 21-22.
 - Answer to inquiry of J. B. Ellis; description of larva, habits, transformations, and means against an undetermined noctuid.
- 182. [Walsh, B. D.] [Bag-worms.] < Pract. Ent., November, 1866, v. 2, p. 22.
 - Answer to inquiry of C. P. Wickersham; characters and synonymy of Thyridopteryx ephemeræformis.
- 183. [Walsh, B. D.] [Ichneumon cocoons.] < Pract. Ent., November, 1866, v. 2, p. 22.
 - Answer to inquiry of S. A. N.; characters of cocoons of Microgaster sp. ?
- 184. WALSH, B. D. Note. < Proc. Ent. Soc. Phil., December, 1866, v. 5, p. 260.
 - Withdraws assertion attributed to C. R. Osten Sacken.
- 185. W[ALSH], B. D. The old-fashioned potato-bugs. <Pract. Ent., December, 1866, v. 2, pp. 25-27, figs.
 - Natural history and means against Lema trilineata, Lytta [= Epicauta] vittata, L. atrata [= E. pennsylvanica], L. marginata [= E. cinerea], and L. cinerea [= Macrobasis unicolor]; figures L. trilineata and E. vittata.

- 186. W[ALSH], B. D. Klippart's wheat plant. < Pract. Ent., December, 1866, v. 2, pp. 27-29.
 - Adverse criticism of the entomological portion of the above-named book.
- 187. W[ALSH], B. D. Trimble's insect enemies of fruit and fruit trees. <Pract. Ent., December, 1866, v. 2, pp. 29-30.

 Favorable review of the above-named book.
- 188. W[ALSH], B. D. Answer to the above, by B. D. W. < Pract. Ent., December, 1866, v. 2, pp. 31-33, figs.
 - Answers to questions asked in J. S. Houghton's "Insects in the Orchard" (Pract. Ent., pp. 30-31); food-habits and hibernation of Conotrackelus nanuphar; comparative characters and figures of Aspidiotus conchiformis [=Mytilaspis pomorum] and Coccus harristi [=Chionaspis furfurus]; natural history of C. harristi; means against and enemies of the two species; figures Chilocorus bivulnerus and Hippodamia [=Megilla] maculata; food-habits and synonymy of Lytta [=Pomphopæa] sayi; flowing as a means against root-feeding insects; inefficacy of salt, lime, ashes, etc., against the same; insects attacking evergreens not likely to injure fruit trees; cannibal habits of spiders; habits of bees and wasps.
- 189. [Walsh, B.D.] [Cicada; gooseberry-worm.] < Pract. Ent., December, 1866, v. 2, p. 33.
 - Answer to inquiry of M. S. Hill; Cicada [= Tibicen] septendecim compared with an undescribed species; ravages of Pristiphora grossulariæ? on gooseberry bushes.
- 190. [WALSH, B. D.] [Vespidæ.] < Pract. Ent., December, 1866, v. 2, pp. 33, 34.

 Answer to inquiry of M. Hobart; life-habits of Vespidæ.
- 191. [Walsh, B. D.] [Meal-worms: Timber-borers.] < Pract. Ent., December, 1866, v. 2, p. 34.
 - Answer to inquiry of M. C. D.; characters, food-habits, and means against *Tenebrio molitor* and *T. obscurus*; characters of *Ptinus brunneus* infesting pine timber; corrosive sublimate as a means against timber insects.
- 192. [Walsh, B. D.] [Corn-worm.] < Pract. Ent., December, 1866, v. 2, p. 34.
 - Answer to J. B. Ellis; number of broads of *Heliothis armigera* in the Northern and Southern States.
- 193. [Walsh, B.D.] [Thousand-legged worms.] < Pract. Ent., December, 1866, v. 2, pp. 34-35, fig.
 - Answer to inquiry of G. W. Robinson; description and figure of Iulus multistriatus n. sp. [= Cambala annulata]; habits and means against the same.
- 194. [Walsh, B. D.] [Cranberry-galls, etc.] < Pract. Ent., December, 1866, v. 2, p. 35.
 - Answer to inquiry of W. C. Fish; occurrence of galls made by *Cecidomyia* sp. on leaves of cranberry; identification of several insects.
- 195. [Walsh, B. D.] Another humbug. < Pract. Ent., December, 1866, v. 2, p. 35.
 - Extract, with comments, from Cultivator and Country Gentleman, 22 November, 1866, exposing worthlessness of P. B. Sheldon's composition for destroying borers,

- 196. [Walsh, B. D.] Notices. < Pract. Ent., December, 1866, v. 2, p. 35.
 - . Mention of several agricultural journals.
- 197. Walsh, B. D. On the insects, coleopterous, hymenopterous, and dipterous, inhabiting the galls of certain species of willow. Part 2d and last. < Proc. Ent. Soc. Phil., December, 1866—January, 1867, v. 6, pp. 223–288.
 - See No. 46 for Part 1st. Supplementary notes on the Cocidomyida (gall-makers and guest-files); criticism of statements of Fitch and Harris; discussion of structure and habits of Tenthredinida and insects generally; synopsis of tenthredinidous willow-galls; descriptions of the galls, their makers and inquilines; for the new species described, see the Systematic Index; remarks on species and varieties; figures fore-wings of Tenthredo and Ichneumon.
- 198. W[ALSH], B. D. Plant-lice—their friends and enemies. < Pract. Ent., January, 1867, v. 2, pp. 37-44, figs.
 - Method of work and mode of reproduction of Aphididæ; their relations to ants; plant-lice enemies; figures Aphie mali and several enemies of plant-lice; means against Aphididæ.
- 199. W[ALSH], B. D. Birds versus insects. < Pract. Ent., January, 1867, v. 2, pp. 44-47.
 - Insectivorous birds not necessarily beneficial; a bird must be shown to eat at least thirty (30) times as many injurious insects as it does beneficial ones before it can be considered useful.
- 200. [Walsh, B. D.] [Insects named.] < Pract. Ent., January, 1867, v. 2, p. 47.
 - Answer to inquiry of J. Pettit; identification of many species, mostly beetles; characters and food-habits of Clorus nigripes [=4-guttatus]; habitat of Leptura capitata and Tingis [=Corythuca] ciliata.
- 201. [Walsh, B. D.] [Apple-tree insects.] < Pract. Ent., January, 1867, v. 2, p. 47.
 - Answer to inquiry of M. Hobart; increasing distribution of Aspidiotus conchiformis [= Mytilaspis pomorum]; enemies of the same; characters of the eggs of Aphis mali.
- 202. [Walsh, B. D.] [Work in entomology.] < Pract. Ent., January, 1867, v. 2, p. 47.
 - Answer to G. Scarborough; advice for the study of entomology by beginners.
- 203. W[ALSH], B. D. The true *Thrips* and the bogus *Thrips*. < Pract. Ent., February, 1867, v. 2, pp. 49-52, figs.
 - Natural history, characters, habits, and figures of *Thrips* sp.; characters of the larva and image of *Haltica chalybea*; figure of image; habits of species of *Erythroneura* [= Typhlocyba]; description and figures of E. [=T.] vitis.
- 204. W[ALSH], B. D. Universal remedies. < Pract. Ent., 1867, v. 2, p. 52.
 - Exposure of proposed universal remedy against fruit-tree insects.
- 205. W[ALSH], B. D. Poisoning noxious insects. <Pract. Ent., February, 1867, v. 2, pp. 52-53.
 - Sugaring with poison as a means against Noctuides.

- 206. W[ALSH], B. D. Remarks by B. D. W. < Pract. Ent., February, 1867, v. 2, p. 54.
 - Comments on S. S. Rathvon's "Bag-worms" (Pract. Ent., pp. 53-54); distribution and food-plants of Thyridoptoryx ophemeræformis.
- 207. W[ALSH], B. D. Habits of the tree-cricket (*Ecanthus niveus*). <Pract. Ent., February, 1867, v. 2, p. 54, figs.
 - Characters of eggs and method of deposition; characters, food-habits, and figures of the imago.
- 208. [Walsh, B. D.] Importing European parasites. < Pract. Ent., February, 1867, v. 2, pp. 54-55.
 - Imaginary correspondence between Asa Fitch and John Curtis on the importation of parasitic insects.
- 209. [Walsh, B. D.] [Screw-worm.] < Pract. Ent., February, 1867, v. 2, p. 55.
 - Answer to inquiry of L. D. Morse; undetermined larva found in osageorange seed; hominivorous habits of *Lucilia macellaria*.
- 210. [Walsh, B. D.] [Attacus cecropia.] < Pract. Ent., February, 1867, v. 2, p. 55.
 - Answer to inquiry of T. T. Smith; characters of larva and image of Attacus occropia; food-plants; method of issuance from cocoon; nature of fluid discharged when emerging from cocoon.
- 211. [Walsh, B. D.] [Apple worm.] < Pract. Ent., February, 1867, v. 2, pp. 55-56.
 - Answer to inquiry of W. C. Fish; characters of the pups of Sciara mali.
- 212. [Walsh, B. D.] [Oak-gall.] < Pract. Ent., February, 1867, v. 2, p. 56.
 - Answer to inquiry of W. Muir; characters of an undetermined cynipid gall on oak.
- 213. [Walsh, B. D.] [Cicada: Grape-vine Procris.] < Pract. Ent., February, 1867, v. 2, p. 56.
 - Answer to inquiry of M. S. Hill; variations in the image of Cloada [= Tibicen] septendecim and in the song of the same; characters of the larva of Procris [= Harrisina] americana.
- 214. [WALSH, B. D.] [Nitidulidæ.] < Pract. Ent., February, 1867, v. 2, p. 56.
 - Answer to inquiry of W. H. S.; food-habits of Ips fasciatus and Nitidula bipustulata.
- 215. [Walsh, B. D.] [Turnip enemy.] < Pract. Ent., February, 1867, v. 2, p. 56.
 - Answer to inquiry of F. T. Pember; characters and food-habits of *Polydesmus complanatus*.
- 216. [Walsh, B. D.] [Wheat midge.] < Pract. Ent., February, 1867, v. 2, p. 57.
 - Answer to inquiry of C. P. Wickersham; nature of the membrane enveloping the full-grown larva of *Diplosis tritici*.

- 217. [Walsh, B. D.] [Queries answered.] < Pract. Eut., February, 1867, v. 2, p. 57.
 - Answer to inquiry of R. Middleton; toads, frogs, and spiders beneficial; effects of the stings of various insects.
- 218. [Walsh, B. D.] [Scales.] < Pract. Ent., February, 1867, v. 2, p. 57.
 - Answer to inquiry of M. M. S.; characters and ravages of scales on oleander and orange; means against *Cocoidæ*.
- 219. [WALSH, B. D.] [Hundred-legged worm.] < Pract. Ent., February, 1867, v. 2, p. 57.
 - Answer to inquiry of S. P. Monks; identification of *Polydesmus virginiensis*; distinctive characters of Myriapoda.
- 220. [Walsh, B. D.] [Bark borers.] < Pract. Ent., February, 1867, v. 2, pp. 57-58.
 - Answer to inquiry of C. V. Riley; synoptic table and food-habits of several species of *Scolytus*; descriptions of *S. fagi* n. sp. and *S. caryæ* n. sp. [=4-spinosus].
- 221. [Walsh, B. D.] [Aleurodes.] < Pract. Ent., February, 1867, v. 2, p. 58.
 - Answer to inquiry of D. F. C.; characters and food-plants of an undescribed species of Alewodes.
- W[ALSH], B. D. The critic criticized. < Pract. Ent., February, 1867, v. 2, p. 58. Answer [by C. V. Riley]. < Prairie Farmer, 16 March, 1867, [v. 35], n. s., v. 19, p. 169. S.-b., No. 2, p. 58. Criticism of article by C. V. Riley (Prairie Farmer, 19 January, 1867, [v. 35], n. s., v. 19, p. 37).
- 223. [Walsh, B. D.] Obituary. < Pract. Ent., February, 1867, v. 2, p. 58.
 - Notice of the late Brackenridge Clemens.
- 224. W[ALSH], B. D. Wire-worms. < Pract. Ent., March, 1867, v. 2, pp. 61-62, figs.
 - Characters of Myriapoda and larvæ of Elateridæ, popularly called wire-worms; food-habits of larva of Ludius attenuatus; figure of larva and image of the same; confusion caused by the use of the term "wire-worms" by correspondents of the Country Gentleman.
- 225. W[ALSH], B. D. Imported insects.—The onion-fly. < Pract. Ent., March, 1867, v. 2, p. 64, figs.
 - Figures of larva and image of Anthonyia coparum and of Ortalis [= Tritoxa] flexa; comparative characters of images; distribution and means against the two species.
- 226. W[ALSH], B. D. [Means against cut-worms.] < Pract. Ent., March, 1867, v. 2, p. 66.
 - Becommends, in reply to J. Townley's inquiry, sugaring with poisoned molasses as a means against cut-worms.

- 227. W[ALSH], B. D. Entomology indeed run mad. <Pract. Ent., March, 1867, v. 2, pp. 66, 67.
 - Reprint of "The bud-worm" (N. C. Rural Journ., September, 1866), with criticism and correction of some of the errors.
- 228. W[ALSH], B. D. The imported gooseberry saw-fly. <Pract. Ent., March, 1867, v. 2, p. 67.
 - Reason for re-appearance in June and for the non-appearance of the second broad of Nematus ventricosus [=ribesii] in New York in 1866.
- 229. W[ALSH], B. D. Tree cut-worms. < Pract. Ent., March, 1867, v. 2, p. 67.
 - Cut-worm described (Pract. Ent., v. 1, pp. 85-86) identified as Hadena chenopodii [= Mamestra trifoli]. See Nos. 85, 281.
- 230. W[ALSH], B. D. Doctoring fruit trees again. <Pract. Ent., March, 1867, v. 2, pp. 67, 68.
 - Extract from Industrial Gazette, 15 December, 1866, with criticism; calomel, sulphur, and substances insoluble in water not absorbed into the circulation of trees.
- 231. W[ALSH,] B. D. The grape-vine Colaspis (Colaspis flavida Say). < Pract. Ent., March, 1867, v. 2, pp. 68-69, fig.
 - Ravages of *Colaspis flavida*; comparison with allied species; remarks on the law of priority and on certain insects becoming suddenly injurious; figures *C. flavida*.
- 232. W[ALSH], B. D. Another universal remedy. <Pract. Ent., March, 1867, v. 2, p. 69.
 - Extract from reprint (f) in Cultivator and Country Gentleman, 10 January, 1867, with criticism of proposed remedy against insects injurious to trees.
- 233. W[ALSH], B. D. Borers. The plug-ugly theory. <Pract. Ent., March, 1867, v. 2, p. 69.
 - Criticism of theory that a plug driven into a hole in a tree is death to borers.
- 234. W[ALSH], B.D. A mass of mistakes. < Pract. Ent., March, 1867, v. 2, p. 70.
 - Correction of errors in "Remedy for the borer" (Prairie Farmer, 9 February, 1867).
- 235. W[ALSH], B. D. Hop-growing in the West. < Pract. Ent., March, 1867, v. 2, p. 70.
 - Extract from Country Gentleman, 31 January, 1867, showing ravages of Phorodon humuli in Vermont; danger of its introduction into the Western States.
- 236. W[ALSH], B. D. Thousand legged worms. < Pract. Ent., March, 1867, v. 2, p. 70.
 - Extract from letter of J. H. Parsons, with comments; injuries to onions by Iulus multistriatus [= Cambala annulata]; I. multistriatus undistinguishable from 1. caruleo-cincius.

- 237. W[ALSH], B. D. Fighting the Curculio. < Pract. Ent., March, 1867, v. 2, p. 71.
 - Extract from Genesee Farmer for 1853, p. 125; ridicule of additions to "jarring" as a means against Constructelus nonuphar.
- 238. W[ALSH], B. D. ["Bushels of butterflies."] < Pract. Ent., March, 1867, v. 2, p. 71.
 - Extract from Cincinnati Gazette, August, 1866, with comments; occurrence of dead images of Attacus occropia? in large numbers ("10 bushels") in Newport, Ohio; food-plants of the larva.
- 239. [Walsh, B. D.] [Scab on potato.] < Pract. Ent., March, 1867, v. 2, pp. 71-72.
 - Answer to inquiry of T. L. J. Baldwin; account of blister-like scabs on potatoes, probably caused by *Sciara* sp.; means against the same.
- 240. [Walsh, B. D.] [Mass of eggs.] < Pract. Ent., March, 1867, v. 2, p. 72.
 - Answer to inquiry of E. Daggy; egg-masses of Clisiocampa americana; food-plants of C. americana and C. sylvatica [=disstria].
- 241. [Walsh, B. D.] [Food of katydids.] < Pract. Ent., March, 1867, v. 2, p. 72.
 - Answer to inquiry of C. M.; food of Microcentrum retinervis largely carnivorous.
- 242. [Walsh, B. D.] [Caterpillar nest on wild cherry.] < Pract. Ent. March, 1867, v. 2, p. 72.
 - Answer to inquiry of H. Morey; undetermined larva on wild cherry, food-plants of Hyphantria textor [=cunea].
- 243. [Walsh, B. D.] [Insect eggs.] < Pract. Ent., March, 1867, v. 2, p. 72.
 - Answer to inquiry of P. Ferris; undetermined eggs of Lepidopters and Hemipters on apple-tree twigs.
- 244. [Walsh, B. D.] [Attacus polyphemus, etc.] < Pract. Ent., March, 1867, v. 2, pp. 72-73.
 - Answer to inquiry of M. M. S.; variations in Attacus [= Teles] polyphemus; habits and times of transformations of Arctia [= Pyrrharctia] isabella; references to history of Dryocampa rubicunda.
- 245. [WALSH, B. D.] [Cossus robiniæ, etc.] < Pract. Ent., March, 1867, v. 2, p. 73.
 - Answer to inquiry of J. Townley; food-habits and injuries of Cossus robinies; result of freezing insects; characters of some wood-borers.
- 246. [Walsh, B. D.] [Museum pests.] < Pract. Ent., March, 1867, v. 2, p. 73.
 - Answer to inquiry of F. L. Van Arsdale; means against Dermestide.
- 247. [Walsh, B. D.] [Katydid eggs.] < Pract. Ent., March, 1867, v. 2, p. 73.
 - Answer to C. M. B.; characters of eggs of Microcentrum retinervis.

- 248. [Walsh, B. D.] [Squash-vine borer.] < Pract. Ent., March, 1867, v. 2, p. 73.
 - Answer to inquiry of E. Nason; ravages and means against Trochilium cucurbita [= Melittia octo].
- 249. [Walsh, B. D.] [Apple-twig borers, etc.] < Pract. Ent., March, 1867, v. 2, pp. 73-74.
 - Answer to inquiry of A. W. Brumbaugh; work of Bostrichus [= Amphicerus] bicaudatus; characters and habits of Taniopteryx fasciata; ravages, habits, and means against Macrodactylus subspinosus.
- 250. [Walsh, B. D.] [Hessian fly.] < Pract. Ent., March, 1867, v. 2, p. 74.
 - Answer to inquiry of E. E. Sheldon; hibernation of the larva of Cecidomyia destructor.
- 251. [WALSH, B. D.] [Tree-cricket.] < Pract. Ent., March, 1867, v. 2, p. 74.
 - Answer to inquiry of J. M. Cole; food-habits of *Œcanthus niveus*; characters of its eggs.
- 252. [Walsh, B. D.] [Tineids in bee-hives.] < Pract. Ent., March, 1867, v. 2, p. 74.
 - Answer to inquiry of J. H. Hunt; characters of case and larva of undetermined tineid.
- 253. W[ALSH], B. D. Jumping to conclusions. < Pract. Ent., March, 1867, v. 2, p. 74.
 - Criticism of article "THE potato-bug" (Wisconsin Farmer, 2 March, 1867).
- 254. W[ALSH], B. D. The common Curculio and its allies. < Pract. Ent., April, 1867, v. 2, pp. 75-81.
 - Characters of the Rhyncophora; characters, ravages, life-history, food-habits, number of broods, and means against Conotrachelus nenuphar; ravages, habits, characters, and means against Anthonomus prunicida [= Coccotorus scutellaris] and A. 4-gibbus; mention of additional injurious weevils; comparative characters of, the three species.
- 255. W[ALSH], B. D. Remarks by B. D. W. < Pract. Ent., April, 1867, v. 2, p. 82.
 - Comments on C. V. Riley's article, "The imported apple-tree bark-louse" (Prairie Farmer, 23 March, 1867, [v. 35], n. s., v. 19, p. 184).
- 256. W[ALSH], B. D. Confessing the corn. < Pract. Ent., April, 1867, v. 2, p. 82.
 - Acknowledgment of mistake in quoting from Prairie Farmer.
- 257. [WALSH, B. D.] [Bibio albiponnis.] < Pract. Ent., April, 1867, v. 2, p. 83.
 - Answer to inquiry of H. Burt; habits of larva of Bibio albipennis.
- 258. [Walsh, B. D.] [Lice on horses.] < Pract. Ent., April, 1867, v. 2, p. 83.
 - Answer to inquiry of L. West; comparative characters of Pediculina and Mallophaga.

- 259. [Walsh, B. D.] [Cocoon on wild cherry.] < Pract. Ent., April, 1867, v. 2, p. 83.
 - Answer to inquiry of A. A. Baker; sexual differences of Attacus promethea; food-plants of its larva.
- 260. [Walsh, B. D.] [Lepidopterous case-bearer.] < Pract. Ent., April, 1867, v. 2, p. 83.
 - Answer to inquiry of J. H. Hunt; larval characters of case-bearing Lepidoptera.
- 261. [Walsh, B. D.] ["Thousand-legged worms."] < Pract. Ent., April, 1867, v. 2, p. 83.
 - Answer to inquiry of J. H. Parsons; Iulus virgatus eating dead seeds; injurious habits of species of Polydesmus.
- 262. [WALSH, B. D.] [Toads: Glorified squash-bug.] < Pract. Ent., 1867, v. 2, p. 83.
 - Answer to inquiry of F. C. Hill; food of toads; poisonousness of Prionotus novemarius [= Prionidus cristatus].
- 263. [WALSH, B. D.] [Galls on blackberry.] < Pract. Ent., April, 1867, v. 2, pp. 83-84.
 - Answer to inquiry of J. H. Tice; identification of the gall of Diastrophus nebulosus; genera of Cynipidæ confined to single plants.
- 264. [Walsh, B. D.] [Bark-lice.] < Pract. Ent., April, 1867, v. 2, p. 84.
 - Answer to inquiry of Dr. Houghton; comparative harmlessness of Coccus karrisii [= Chionaspis furfurus]; means against Coccidæ.
- 265. [Walsh, B. D.] [Plant-lice.] < Pract. Ent., April, 1867, v. 2, p. 84.
 - Answer to inquiry of E. Orton; description of eggs of Lachnus strobi; means against Aphidida.
- 266. [Walsh, B. D.] [Bag-worms.] < Pract. Ent., April, 1867, v. 2, pp. 84-85.
 - Answer to inquiry of J. Murphy; food-habits and means of dispersal of Thyridopteryx ephemeræformis.
- 267. [Walsh, B. D.] [Insects injurious to grape.] < Pract. Ent., April, 1867, v. 2, p. 85.
 - Answer to inquiry of J. Wood; supposed Curculio punctures in grapes.
- 268. [Walsh, B. D.] [Attacus promethea.] < Pract. Ent., April, 1867, v. 2, p. 85.
 - Answer to inquiry of J. B. Lyon; mention of undetermined parasite of Attacus promethea.
- 269. [Walsh, B. D.] [Borer in hickory.] < Pract. Ent., April, 1867, v. 2, p. 85.
 - Answer to inquiry of B. Norris; characters of young and mature larva of Clytus [= Cyllene] pictus.
- 270. [Walsh, B. D.] [Apple-tree plant-lice.] < Pract. Ent., April, 1867, v. 2, p. 85.
 - Answer to inquiry of W. W. Linn; identification of eggs of Aphia mali; enemies of the same.

- 271. [Walsh, B. D.] [Tulip-tree bark-louse: Basket-worm.] < Pract. Ent., April, 1867, v. 2, p. 85.
 - Answer to inquiry of I. Hicks; parasites and enemies of undescribed barklouse on tulip-tree; Thyridopteryx ephemeraformis destructive to evergreens.
- 272. W[ALSH], B. D. The grape-vine Fidia. (Fidia viticida, new species.) < Pract. Ent., May, 1867, v. 2, pp. 87-88, fig.
 - Ravages, description, and figure of *Fidia viticida* n. sp.; modification of the clasping organs of insects.
- 273. W[ALSH], B. D. Enemies of the rice crop. < Pract. Ent., May, 1867, v. 2, p. 89.
 - Injuries caused by and means adopted against the water-weevil [= Lissorhop-trus simplex] and the rice-grub = [Chalepus trachypygus.]
- 274. [Walsh, B. D.] The canker-worm once more. <Pract. Ent., May, 1867, v. 2, p. 89. Abstract: <Cultivator and Country Gentleman, 6 June, 1867, v. 29, p. 370.
 - Extract from and review of W. G. C.'s "My trials with the canker-worms: remedy" (Iowa Homestead, 13 March, 1867); sorghum as a means against Anisopteryx.
- 275. W[ALSH], B. D. None so blind as those who shut their eyes. <Pract. Ent., May, 1867, v. 2, pp. 89-90.
 - Criticism of error by editors of agricultural papers; rate of eastward progress of Doryphora 10-lineata.
- 276. W[ALSH], B. D. Self-taught entomologists. < Pract. Ent., May, 1867, v. 2, pp. 91-92.
 - Criticism of misstatements found in agricultural papers; first occurrence of Doryphora 10-lineata in Illinois in 1864.
- 277. W[ALSH], B. D. Pear-tree and apple-tree insects. < Pract. Ent., May, 1867, v. 2, pp. 92-93.
 - Food-plants of Attacus cecropia, Clisiocampa americana, and Saperda bivittata [=candida].
- 278. W[ALSH], B. D. More universal remedies. < Pract. Ent., May, 1867, v. 2, p. 93.
 - Reprint and criticism of "Destruction of insects" (Mo. Rept. Bureau Agric., February, 1867, p. 60).
- 279. W[ALSH], B. D. Hop-growing in the West. < Pract. Ent., May, 1867, v. 2, pp. 93-94.
 - Occurrence in Michigan of Phorodon humuli; its destructiveness; popular ignorance of eutomology.
- 280. W[ALSH], B. D. The wheat midge—jumping to a conclusion. <Pract. Ent., May, 1867, v. 2, p. 94.
 - Extract, with comment, from Maryland Farmer and Mechanic, August, 1865;

 Diplosis tritici not infested with parasites in America; food-habits of
 Thrips.

- 281. W[ALSH], B. D. Tree cut-worms. < Pract. Ent., May, 1867, v. 2, p. 94.
 - "Dark-sided cut-worm" an Agrotis and not Hadena chenopodii [= Mamestra trifolii]. See Nos. 85,229.
- 282. W[ALSH], B. D. Quacks and physicians. < Pract. Ent., May, 1867, v. 2, p. 95.
 - Criticism of proposed universal remedy against insects injurious to the rose (Western Rural, 30 March, 1867).
- 283. W[ALSH], B. D. The ephemeron or May-fly. < Pract. Ent., May, 1867, v. 2, p. 95.

Length of duration of life of Ephemerida.

- 284. W[ALSH], B. D. A groundless fear. < Pract. Ent., May, 1867, v. 2, p. 95.
 - Extract from Mo. Rept. Bureau Agric., February, 1867, p. 62; food-habits of larva and image of Lytta atrata [= Epicauta pennsylvanica]; places of oviposition.
- 285. W[ALSH], B. D. Fire-blight. <Pract. Ent., May, 1867, v. 2, p. 96.

Probable cause.

- 286. [Walsh, B. D.] [Insect eggs.] <Pract. Ent., May, 1867, v. 2, p. 96.
 - Answer to inquiry of W. Willock; identification of eggs of *Œœnthus niveus* and of one of the *Membracidæ* on grape twigs.
- 287. [Walsh, B. D.] [Attacus cecropia.] < Pract. Ent., May, 1867, v. 2, p. 96.

Answer to inquiry of J. B. H.; characters of Attacus cocropia.

- 288. [WALSH, B. D.] [Twig pruners.] < Pract. Ent., May, 1867, v. 2, p. 96.
 - Answer to inquiry of M. W. Philips; twigs of pear and other trees amprated in the same manner as oak twigs by Elaphidion putator [=villoum.
- 289. [WALSH, B. D.] [Blister-beetles.] < Pract. Ent., May, 1867, v. 2, p. 97.
 - Answer to inquiry of J. M. Tracy; food-habits, ravages, and means against $Lytta \ [= Pomphop \infty a]$ anea; food-plants of $L. \ [= P.]$ tarsalis; tabular separation of three species of $Lytta \ [= Pomphop \infty a]$.
- 290. [WALSH, B. D.] [Means against Curculio: *Œstridæ*.] < Pract. .

 Ent., May, 1867, v. 2, p. 97.
 - Answer to inquiry of C. Greene; ineffective means against Constructeus nonuphar; Æstridæ infesting the ox and rabbit.
- 291. [Walsh, B. D.] [Insect eggs.] < Pract. Ent., May, 1867, v. 2, p. 97.
 - Answer to inquiry of P. Ferris; identification of eggs of Clisicoampa sylvatica [—disstriu] and Aphis mali.

292. W[ALSH], B. D. The wheat midge. <Pract. Ent., June, 1867, v. 2, pp. 99-101.

Extract from letter from S. S. Rathvon, with remarks; natural history of Diplosis tritici.

293. W[ALSH], B. D. The new or Colorado potato-bug. <Pract. Ent., June, 1867, v. 2, pp. 101-102.

Distribution, habits, and extent of injuries of Doryphora 10-lineata.

294. [Walsh, B. D.] [Gall-flies; saw-flies.] < Pract. Ent., June, 1867, v. 2, pp. 102-103.

Answer to inquiry of M. W. Philips; mention of various insects which are gall-makers; their modes of life; characters of the larvæ of Cecidomyidæ and Tenthredinidæ.

295. [Walsh, B. D.] [Bark-beetles.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of W. C. Fish; identification of specimens sent; habits of Tomicus pusillus [= Pityophthorus minutissimus] and of T. ramulorum [= P. miorographus].

296. [Walsh, B. D.] [Apple-tree insects.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of S. Cassi; characters, food-plants, and ravages of Arctia [=Spilosoma] virginica; supposed work of Epicarus imbricatus.

297. [Walsh, B. D.] [Attacus cecropia.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of T. McGraw; larvæ of Attacus cocropia usually solitary.

298. [Walsh, B. D.] [Grape-vine flea-beetle.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of A. Kelley; hibernation of the image of Haltica chalybea; ravages and means against the same.

299. [Walsh, B.D.] [Sphyracephala brevicornis.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of H. B. Howarth; characters and habitat of Sphyrace-phala brevioornis.

300. [Walsh, B. D.] [May-bug.] < Pract. Ent., June, 1867, v. 2, p. 103.

Answer to inquiry of C. H. R.; characters and food-habits of the larva of Lachnosterna quercina [== fusca].

301. [Walsh, B. D.] [Blister-beetles: plant-louse enemy.] < Pract. Ent., June, 1867, v. 2, p. 104.

Answer to inquiry of Dr. Houghton; food-habits of Lytta [= Pomphopæa] amea and L. [== P.] sayi; Chilocorus bivulnerus as a means against Coccida.

302. [Walsh, B. D.] [Blackberry borers.] < Pract. Ent., June, 1867, v. 2, p. 104.

Answer to inquiry of A. S. Fuller; characters of undetermined larva (*Ægeria?*) and of *Oberca perspicillata* [= bimaculata] boring in blackberry stems; food-habits of *Ægeridæ*.

- 303. W[ALSH], B. D. Scientific nomenclature. <Pract. Ent., July, 1867, v. 2, pp. 105-106.
 - Criticism of the practice of establishing new genera upon slight characters, as illustrated by the American silk-worms.
- 304. W[ALSH], B. D. Currant plant-lice (Aphis ribis). < Pract. Ent., July, 1867, v. 2, p. 106.
 - Method of work of Aphis [= Myzus] ribis.
- 305. [Walsh, B. D.] [Lucanus elaphus.] < Pract. Ent., July, 1867, v. 2, p. 107.
 - Answer to inquiry of S. Haycraft; characters and habits of Lucanus elaphus; rarity of the Q; use of prolonged jaws in & insects.
- 306. [Walsh, B. D.] [New York weevil.] < Pract. Eut., July, 1867, v. 2, p. 107.
 - Answer to inquiry of T. Gregg; habits and distribution of Ithycerus noveboracensis.
- 307. [Walsh, B. D.] [Codling-moth.] < Pract. Ent., July, 1867, v. 2, p. 107.
 - Answer to inquiry of H. Morey; number of broods, habits, etc., of Carpocapsa pomonella.
- 308. [Walsh, B. D.] [Hickory borer, etc.] < Pract. Ent., July, 1867, v. 2, p. 107.
 - Answer to inquiry of T. Wiggins; times of emergence of Clytus [=Cyllene] pictus and C. [=C] robiniæ; habits of Corydalus cornutus.
- 309. [Walsh, B. D.] [Wasps.] < Pract. Ent., July, 1867, v. 2, p. 107.

 Answer to inquiry of G. W. Smith; food-habits of digger-wasps.
- 310. [Walsh, B. D.] | Parasitic flies, etc.] < Pract. Ent., July, 1867, v. 2, pp. 107-108.
 - Answer to inquiry of S. P. Monks; method of transformation of parasitic Hymenoptera; use of Latin plurals for English words.
- 311. [Walsh, B. D.] [Workers among Hymenoptera.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of W. V. Andrews; economy of workers among the social Hymenoptera.
- 312. [Walsh, B. D.] [Cut-worms.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of V. Abbey; means against Agrotis sp. ?
- 313. [Walsh, B. D.] [Potato-beetle enemies.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of W. Smith; scutellerid enemy of larva of Doryphora 10-lineata; katydids as an enemy of the same.
- 314. [Walsh, B. D.] [Black-bug.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of J. R. Tewksbury; characters and habits of *Pirates* [= Melanolestes] picipes.

- [Walsh, B. D.] [Burying-beetles.] < Pract. Ent., July, 1867, v. p. 108.
 - Answer to inquiry of A. D. Strong; characters of Neoropherus marginatus; habits of Silphidæ; mite parasites of Silphidæ and other beetles.
- 316. [Walsh, B. D.] [Insects named.] < Pract. Ent., July, 1867, v. 2, p. 108.
 - Answer to inquiry of M. S. Hill; identification of specimens sent; food-plants of larva of Chrysomela [= Gastroidea] cyanea.
- 317. [Walsh, B. D.] [Spring-tails.] < Pract. Ent., July, 1867, v. 2, pp. 108-109.
 - Answer to inquiry of E. Daggy; characters, habits, and means against Poduridæ.
- 318. [Walsh, B. D.] [Blackberry gall insects.] < Pract. Ent., July, 1867, v. 2, p. 109.
 - Answer to inquiry of Aculeus; number of abdominal segments of *Diastrophus* and *Aulax*; reasons for considering the former the gall-maker and the latter the guest-fly.
- 319. [Walsh, B. D.] [Pine and cedar longicorns.] < Pract. Ent., July, 1867, v. 2, p. 109.
 - Answer to inquiry of J. Barratt; sexual characters and food-habits of Callidium antennatum and C. janthinium.
- 320. [Walsh, B. D.] [White-pine plant-louse, etc.] < Pract. Ent., July, 1867, v. 2, p. 109.
 - Answer to inquiry of C. H. Peck; Lachnus strobi on Pinus strobus; undetermined bark-louse, with enormous lateral thoracic plates, found on Rhus glabra.
- 321. [Walsh, B.D.] [Cut-worms.] < Pract. Ent., July, 1867, v. 2, p. 109.
 - Answer to inquiry of J. Townley; food-habits of climbing cut-worms.
- 322. [WALSH, B. D.] [Current plant-louse.] < Pract. Ent., July, 1867, v. 2, pp. 109, 110.
 - Answer to inquiry of R. L. Walker; Coccinellida as a means against Aphie [= Myzue] ribis.
- 323. [WALSH, B. D.] [Beetles.] < Pract. Ent., July, 1867, v. 2, p. 110.

 Answer to inquiry of I. A. Plucke; characters and food-habits of Tetraopes tornator [= tetraophihalmus] and of T. femoratus; comparative characters of Chrysomela bigsbyana and of C. philadelphica; characters of Chrysomelidæ and Coccinellidæ.
- 324. W[ALSH], B. D. Fertilizing plants. <Pract. Ert., July, 1867, v. 2, p. 110.
 - Part played by insects in fertilizing plants.
- 325. W[ALSH], B. D. The peach-worm. < Pract. Ent., July, 1867, v. 2, p. 110.
 - Larvæ of Ephestia zeæ [=interpunctella] injurious to dried peaches; foed-plants and enemies of the same.

- 386. W[ALSH], B. D. Valedictory. <Pract. Ent., August-September, 1867, v. 2, p. 111.
 - Note of thanks for aid received in editing the Practical Entomologist.
- 327. W[ALSH], B. D. The State entomologist of Illinois. <Pract. Ent., August-September, 1867, v. 2, p. 111. Reprint: <Can. Farmer, 1 October, 1867, v. 4. S.-b., No. 2, p. 98.
 - Bill for appointment of State entomologist of Illinois not carried into effect.
- 328. W[ALSH], B. D. The three so-called army-worms. < Pract. Ent., August-September, 1867, v. 2, pp. 111-114.
 - Confusion caused by the use of the same popular name for Anomis [= Aletia] xylina, Leucania unipuncta, and Clisiccampa sylvatica [= disstria]; habits, etc., of the three species; food-habits of their larvæ; characters, parasites, and means against C. sylvatica [= disstria].
- 329. W[ALSH], B. D. The little Turk and its crescent. < Pract. Ent., August-September, 1867, v. 2, pp. 114-115.

 Comments on letter from F. C. Hill; oviposition of Constructed semapher.
- 830. [Walsh, B. D.] Apple-worm. (Carpocapsa pomonella.) < Pract. Ent., August-September, 1867, v. 2, p. 115.

 Prevalence of C. pomonella in 1867; food-habits of its larva.
- 331. W[ALSH], B. D. Spindle-worms. < Pract. Ent., August-September, 1867, v. 2, pp. 115-116.</p>
 Habits, etc., of Achatodes zew and of Gortyna nitela.
- 332. W[ALSH], B. D. A plant growing out of an insect. < Pract. Ent., August-September, 1867, v. 2, p. 116.
 - Larvæ of Lacknosterna fusca infested with Cordyceps turned up by the plow in large numbers; larvæ supposed to have eaten poisonous seed which has germinated after killing the larvæ; sowing the seed as a means against the larvæ.
- 333. [Walsh, B. D.] The imported gooseberry saw-fly. <Pract. Ent., August-September, 1867, v. 2, p. 116.
 - Changes in the larva of Nematus ventricosus [= ribesii] during the last moult; ravages in Columbia County, N. Y.; extent of spread from Rochester, N. Y.
- 334. W[ALSH], B. D. The Colorado potato-bug. <Pract. Ent., August-September, 1867, v. 2, p. 116.
 - Spread of Doryphora 10-lineata into southern Michigan and western Indiana.
- 335. W[ALSH], B. D. Apple-tree plant-lice. (Aphis mali.) < Pract. Ent., August-September, 1867, v. 2, pp. 116-117. Effects of early frosts on Aphis mali.
- 336. W[ALSH], B. D. The tent-caterpillar of the apple-tree. (Clisio-campa americana.) < Pract. Ent., August-September, 1867, v. 2, p. 117.
 - Scarcity of C. americana in 1867 due to the abundance of an egg parasite.

- 337. [Walsh, B. D.] A new foe of the corn. < Pract. Ent., August-September, 1867, v. 2, pp. 117-118.
 - Ravages and description of Sphenophorus zee n. sp. [=sculptilis]; habits of the larva and image.
- 338. [Walsh, B. D.] [Rhyssa atrata.] < Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of H. K. Smith; Rhyssa [= Thalessa] atrata ovipositing in hickory.
- 339. [Walsh, B. D.] [Grape-vine beetles.] < Prect. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of L. D. Morse; ravages and food-plants of Figia viticida and of F. longipes.
- 340. [Walsh, B. D.] [Current insects.] < Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of A. M. Burns; identification of *Egeria tipuliformis*; mention of species called currant-worms; destructiveness of *Nomatus ventricosus* [=ribesii].
- 341. [Walsh, B. D.] [Rascal leaf-crumpler.] < Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of J. M. K.; times of transformation and means against Physita nebulo [= Acrobasis indiginella].
- 342. [Walsh, B. D.] [Strawberry enemy.] < Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of C. Faxon; characters, habits, and means against Polydesmus serratus; roots of strawberry plants injured by the same.
- 343. [Walsh, B. D.] [Eight-spotted forester.] < Pract. Ent., August-September, 1867, v. 2, p. 118.
 - Answer to inquiry of A. Gilbert; characters and food-plants of Alypia octomaculata.
- 344. [Walsh, B. D.] [Maple bark-louse.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of B. W. McLain; characters of undescribed coccid on maple.
- 345. [Walsh, B. D.] [Apple-tree bark-lice.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of A. Gilbert; identification of Aspidiotus karrisii [= Chionaspis furfurus] and of A. conchiformis [= Mytilaspis pomorum] on appletrees.
- 346. [Walsh, B. D.] [Habits of larvæ and pupæ.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of M. M. S.; insects which transform under ground enter the earth in the larval stage; apparatus used by pupse in forcing their way to the surface of the ground.
- 347. [Walsh, B. D.] [Potato-beetles: Cicindela sex-guttata.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of P. B. Sibley; mention of Lytta cinerea [= Macrobasis unicolor], and of Doryphora 10-lineata as destructive to potato vines; supposed larval habits of Cicindela sex-guttata.

- 348. [Walsh, B. D.] [Wavy-striped flea-beetle.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of E. T. Snelling; food-habits of Haltica striolata [=Phyllotreta vittata].
- 349. [Walsh, B. D.] [Parasites of cut-worms.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of J. Edgerton; metamorphoses of tachinid parasites of Noctuida.
- 350. [Walsh, B. D.] [Eggs on sugar-maple.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of W. Prichard; egg-rings of Clisicoampa americana? on sugar-maple.
- 351. [Walsh, B. D.] [Click-beetles.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of J. H. Foster, jr.; food-habits of Melanotus communis and of Monocrepidius vespertinus.
- 352. [Walsh, B. D.] [Chrysobothris femorata.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of J. J. Kelly; food-habits of Chrysobothris femorata.
- 353. [Walsh, B. D.] [Gall on chestnut.] < Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of W. Kite; mention of an undetermined Phylloxera gall on the flower-catkin of the chestnut.
- 354. [Walsh, B. D.] [Grape-vine beetle: Bugs.] <Pract. Ent., August-September, 1867, v. 2, p. 119.
 - Answer to inquiry of B. F. Lazear; habits of *Pelidnota punctata*; swarming of *Corimelæna pulicaria*; defensive odors of Heteroptera.
- 355. [Walsh, B. D.] [Squash-vine insects.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of J. H. Parasses; larval habits of Diabrotica vittata; confusion caused by popular names.
- 356. [WALSH, B. D.] [Mites.] < Pract. Ent., August-September, 1867; v. 2, p. 120.
 - Answer to inquiry of C. Carpenter; characters and life-habits of Acarina.
- 357. [Walsh, B. D.] [Haltica cucumeris.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of G. S.; food-habits of Haltica [= Crepidodera] cucumeris.
- 358. [Walsh, B. D.] [Insects named.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of M. H. Boye; food-habits of Conotrachelus cratægi, Melanotus communis, Pelidnota punctata, Desmocerus palliatus, Lucidota atra, Podabrus rugulosus, and P. modestus; variation of Haltica [= Crepidodera] helxines; characters of Dolerus arvensis.

- 359. [WALSH, B. D.] [Insects named.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of H. Burt; identification of several insects; climbing habits of cut-worms.
- 360. [Walsh, B. D.] [Hickory-galls.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of W. C. Fish; characters of galls made by Phylloxera caryæglobuli and P. caryæcaulis; identification of several beetles.
- 361. [Walsh, B. D.] [Yellow swallow-tail; Rose-beetle.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of J. Barratt; identification of *Papilio turnus*; transformation and ravages of *Macrodactylus subspinosus*.
- 362. [Walsh, B. D.] [Blister-beetles.] < Pract. Ent., August-September, 1867, v. 2, p. 120.
 - Answer to inquiry of D. W. Kauffman; food-plants and characters of Macrobasis murina [= unicolor] and of Epicauta atrata [= pennsylvanica].
- 363. [Walsh, B. D.] [Tent-caterpillars.] < Pract. Ent., August-September, 1867, v. 2, p. 121.
 - Answer to inquiry of S. R. Williams; obcurrence of Clisicoampa americans on pear-trees and of C. sylvatics [==distria] on black-walnut.
- 364. [Walsh, B. D.] [Currant-worms.] < Pract. Ent., August-September, 1867, v. 2, p. 121.
 - Answer to inquiry of A. H. Mills; characters of the larva of Nematus ventricosus [=ribesii], of Ellopia [= Euftchia] ribearia, and of Pristiphora grossularia.
- 365. [Walsh, B. D.j [Potato-beetle enemies.] < Pract. Ent., August-September, 1867, v. 2, p. 121.
 - Answer to inquiry of W. Maus; description and habits of Lebia grandis; hemipterous enemies of Doryphora 10-lineata.
- 366. W[ALSH], B. D. The Colorado potato-bug. <Pract. Ent., August-September, 1867, v. 2, p. 121.
 - Extract from Mo. Rept. U. S. Agric. Bureau, 1866, p. 344; doubts the occurrence of *Doryphora* 10-lineata in Pennsylvania.
- 367. [Walsh, B. D.] Entomology. < Prairie Farmer, December, 1867, [v. 36], n. s., v. 20, pp. 358-359.
 - Address, with discussion by others, at the third annual meeting of the Southern Illinois Fruit Growers and Shippers' Association; ravages and means against Pemphigus pyri [= Schizoneura lanigera], Carpocapea pomonella, Trypeta pomonella, Semasia [= Grapholitha] pruinivora, Caliodes [= Craponius] inaqualis, Conotrachelus nenuphar.
- 368. Walsh, B. D. The grasshoppers. <Iowa Homestead, 15 January, 1868 (v. 137), p. 9. S.-b., No. 2, p. 119.
 - Range of swarms of Caloptenus spretus in Iowa in 1867.
- 369. WALSH, B. D. Birds vs. insects. <Prairie Farmer, 1868, [v. 37], n. s., v. 21; 30 May, pp. 346-347; 6 June, pp. 362-363; 13 June, pp. 378, 379.

- 376. Walsh, B. D. The seventeen-year locust. <Dixie Farmer, 11 June, 1868. S.-b., No. 3, p. 23.
 - Periodicity and local distribution of the various broads of Cicads [= Tibicsn] septendecim.
- 371. Walsh, B. D. Entomological. Do locusts sting? < Chicago Republican, 1868. S.-b., No. 3, p. 25.
 - Sting of Cloads not poisonous; sting of Stisus grandis [== Sphecius speciosus] mistaken for it; habits of the Stisus.
- 372. Walsh, B. D. The bug-hunter in Egypt. A journal of an entomological tour into south Illinois by the senior editor. <Amer. Ent., 1868, v. 1; September, pp. 6-14, figs. 1-7; October, pp. 29-36, figs. 23-29.
 - Habits and figures of Macronema sebratum, Palingenia [= Hexagenia] bilineats. Cicada sp., Stizus grandis [= Sphecius speciosus], Xylocopa carolina [= virginica], and Anthophora sponsa [= abrupta]; sting of Cicada not poisonous, that of the Stizus mistaken for it; habits of Conotrachelus nemuphar; description of Hull's Curculio-catcher; Conotrachelus anaglypticus and Colaspis sp. injurious to plum-trees; figures of Rhopalus sp. and Euryomia [= Kuphoria] melancholica injurious to pear-trees; figure and habits of Metapodius nasalus [= femoratus]; predaceous habits of Arma [= Podisus] spinosus. Evagoras viridis [= Diplodus luridus], and Stiretrus fimbriatus [= anchorago]; work of Aspidiotus harrisii [= Chionaspis furfurus] upon pear, and of Ceoidomvia [=: Diplosis] tritici in wheat; occurrence of Endrosa [= Lachnosterna] querous, Doryphorajuncta, and Trichius delta in Illinois; mite enemy of Aspidiotus conchiformis [= Mytilaspis pomorum]; injury to fruit by Apis mellifica; hot water as a means against Schizoneura lanigera; habits and figures of Chrysopa sp.; figures of Aspidoglossa subangulata, Harpalus pennsylvanious. and Evarthrus orbatus; figures and descriptions of larva of a carabid and of Chauliognathus pennsylvanicus; Curculio-feeding habits of Carabida.
- 373. Walsh, B. D. First annual report on the noxious insects of the State of Illinois. <Trans. Ill. State Hortic. Soc. for 1867, n.s., v. 1, Appendix. Separate: <Chicago, 1868, 103 pp., figs. 1-3.

TABLE OF CONTENTS.

Introductory	3
CHAPTER 1. The grape Curculio. Caliodes [= Craponius] inaqualis Say.	
Its history—Guest insects, parasites, and cannibals—Remedy	13
CHAPTER 2. The grape-leaf gall-louse. Dactylosphæra vitifoliæ, Fitch	
[= Phylloxera vastatrix].	
Peculiar to the Clinton and other cultivated varieties of Frost	
grape—Practical inference therefrom	21
CHAPTER 3. The rose-bug. Macrodactylus subspinosus Linn.	
Device for destroying it on grape-vines	24
CHAPTER 4. The grape-root borer. Egeria [= Sciapteron] polistiformis	
Harris.	
Its history—Its scientific nomenclature—Remedies	24
CHAPTER 5. The apple-worm or codling-moth. Carpocapea pomonella	
Linn.	
Double-brooded-Practical inference therefrom-Remedies	27
CHAPTER 6. The apple-maggot fly. Trypeta pomonella Walsh.	
Occurs at present only in certain Eastern States on the apple,	
though it was found long ago in Illinois on the crab-Ita history.	29

373. WALSH, B. D.—Continued. CHAPTER 7. The rascal leaf-crumpler. Physica nebulo Walsh [== Acrobasis indiginella].	
A peculiarly Northwestern species, not found either East or South— Easily destroyed	ı
CHAPTER 8. The oyster-shell bark-louse. Aspidiotus conchiformis Gmel. [:= Mytilaspis pomorum]. Its history—Loses almost all its organs when only a few days old, and becomes thenceforth as stationary as a cabbage—How it spreads from tree to tree—Mites and their natural history—Plantfeeding mites, parasitic mites, and cannibal mites—The bark-louse largely preyed on by a minute cannibal mite—Useless and useful remedies.	1
CHAPTER 9. Harris' bark-louse. Aspidiotus harrisii Walsh [= Chionaspis furfurus].	
How it differs from the preceding—The geographical distribution. 53 CHAPTER 10. The apple-root plant-louse. Pemphigus pyri Fitch [= Schisoneura lanigera]. Often confounded in Illinois with the true "Woolly plant-louse"—	
The differences—Its history—Causes a form of "rotten-root"— Its supposed cannibal foes—Remedies	5
CHAPTER 11. The plum Curculio. Conotrackelus unnupkar Herbst. Its history—Its peculiar crescent cut explained—Double-brooded— Its supposed enemies, the so-called "Curculio parasite" and the Baltimore oriole—Remedies	
CHAPTER 12. The plum-gouger. Anthonomus prunicida Walsh [= Coo- cotorus scutellaris]. Bores a round hole in the plum, instead of a crescent cut—How and why it does this—Differs in many other respects from the	
Curculio	5
Its history—Probably a guest-moth, and therefore not injurious CHAPTER 14. The hateful grasshopper. Caloptenus spreins Walsh. The Rocky Mountains its natural home—Invades in certain years certain neighboring districts, such as Texas, Missouri, Kansas, Nebraska, and Iowa—Lays there millions of eggs, which develop into barren grasshoppers only—Invasion of 1866—Damage done by the young grasshoppers in the spring of 1867—In Kansas about one-eighth of the field crops and seven-eighths of the garden crops destroyed by them—Grashopper invasions of A. D. 1820, 1856, 1857, 1864, and 1867—Probable results in 1868 of the invasion of 1867—Has never yet come within 115 miles of Illinois—Can not spread into Illinois as the Colorado potato-bug has done—Reasons why, in all human probability, it can never	
reach Illinois at all	

374. WALSH, B. D. A friend unmasked. < Amer. Ent., November, 1868, v. 1, pp. 51-52, fig. 55.

Description and figures of larva and adult of Chauliognathus pennsylvanious; characters and habits of the Telephoride.

- 375. WAISH, B. D. Appendix to the article on "Wasps and their habits," in No. 7. < Amer. Ent., April, 1869, v. 1, pp. 162-164. Habits of Chrysididæ; descriptions of Stizus [Megastizus] brevipennis n. sp., Agenia subcorticalis n. sp., A. architectus Say, A. cupida Cress., A. bombyoina Cress., Ceropalès ruftventris n. sp., and Ammophila pictipennis n. sp.; tabular separation of the genera of Sphecidæ. See No. 543.</p>
- 376. Walsh, B. D. Mr. Couper's thorn-leaf gall. <Ca. Ent., 15 April, 1869, v. 1, pp. 79-80.
 - Description of cecidomyidous leaf-gall on Cratagus; mentions of three additional Cecidomyidous, and an Acaridous gall on the same; Anthonomus cratagi inquilinous in gall on Cratagus.
- 377. WALSH, B. D. The six worst insect enemies of the fruit growers in northern Illinois. <Trans. North. 111. Hortic. Soc. for 1867—'68, pp. 91-96.
 - Treats of Carpocapsa pomonella, Aspidiotus conchiformis [=Mytilaspis pomorum],
 Anisopteryz [= Paleacrita] vernata, Saperda bivittata [= candida], Chrysobothris femorata, Conotrachelus nenuphar, and Anthonomus prunicida [= Coocotorus scutellaris].
- 378. WALSH, B. D. Squash borer. <Cultivator and Country Gentleman, 30 September, 1869, v. 34, p. 256. Extract: <Prairie Farmer, 30 October, 1869.
 - Characters of larva and image of *Egeria cucurbita* [= *Melittia ceto*]; natural history, means against, and distribution of the same.
- 379. WALSH, B. D. Apple-tree borers. <Journ. Ill. State Agric. Soc. Reprint: <Trans. Ill. State Agric. Soc., v. 5, pp. 499-501. Comparative differences between Saperda and Chrysobothris; soap as a means against the same.
- 380. Walsh, B. D. On a species of *Hemiteles* (*Ichneumonida*), ascertained by the editor to be parasitic in Canada on the imported current-worm fly (*Nematus ventricosus* Klug). <Can. Ent., 1 October, 1869, v.'2, pp. 9-12.
 - Division of Hemiteles into two groups; description of H. nemativorus n. sp.; larval and pupal history of the same by C. J. S. Bethune.
- 381. Walsh, B. D. The imported currant-worm fly (Nematus ventricosus Klug) and its parasite (Hemiteles nemativorus Walsh) < Ca. Ent., 15 November, 1869, v. 2, pp. 31-33.
 - Correction of errors in No. 380; sexual characters of *Hemiteles*; variation in the time at which the image state is assumed by hibernating insects; heteropterous enemies of *Nematus ventricosus* [= ribesii].
- 382. WALSH, B. D. Larvæ in the human bowels. One of Mr. Walsh's posthumous articles. <Amer. Ent., March, 1870, v. 2, pp. 137–141, fig. 93.
 - Characters and habits of larval *Homalomyia*; description of the larva of *H. wilsoni*, *H. leidyi*, and *H. prunivora* n. spp.; mention of recorded cases of diptera in the human bowels; figures larva of *H. wilsoni*.

- 383. Walsh, B. D. One day's journal of a State entomologist. < Amer. Ent. and Bot., May, 1870, v. 2, pp. 197-199.
 - Sketch of an average day's work of a State entomologist.
- 384. Walsh, B. D. On the group *Eurytomides* of the hymenopterous family *Chalcididæ*; with remarks on the theory of species and a description of *Antigaster*, a new and very anomalous genus of *Chalcididæ*. <Amer. Ent. and Bot., 1870, v. 2, September, pp. 297-301, fig. 1, 2; October, pp. 329-335, fig. 3-6; December, pp. 367-370, fig. 7-10.
 - Descriptions, habits, and figures of many Eurytomides; synoptic tables of the species of Eurytoma and Decatoma. For a list of the new species see the Systematic Index.
- 385. Walsh, B. D. Descriptions of North American Hymenoptera. <Trans. Acad. Sci. St. Louis, 7 May—9 June, 1873, v. 3, pp. 65–166, fig.
 - Printed posthumously, with notes by E. T. Cresson; description of many previously described *Tenthredinida* and *Ichneumonida*; five (5) new genera and forty-five (45) new species of *Ichneumonida* are described, for a list of which see the *Systematic Index*; figures venation of front wing of *Ichneumon*.

4 ENT



U. S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

OF

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,

BY

SAMUEL HENSHAW.


PART II.

THE MORE IMPORTANT JOINT WRITINGS

OF

B. D. WALSH and C. V. RILEY.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1889.



BIBLIOGRAPHY OF THE MORE IMPORTANT CONTRIBUTIONS TO AMERICAN ECONOMIC ENTOMOLOGY.

PART II.

- 386. [Walsh, B. D., and C. V. Riley.] Salutatory. To the agriculturists and horticulturists of the United States. <Amer. Ent., September, 1868, v. 1, pp. 1-3.
 - Importance and extent of the ravages of insects; practicability of the diminution of these ravages; value of entomological work; aims of the American Entomologist.
- 387. [Walsh, B. D., and C. V. Rilley.] Hogs vs. bugs. < Amer. Ent., September, 1868, v. 1, pp. 3-6.
 - Effectiveness of swine as a means against Constrachelus nemuphar and Carpocapsa pomonella; quotations showing good results from pasturing swine in fruit orchards.
- 388. [Walsh, B. D., and C. V. Riley.] Insect changes. < Amer. Ent., September, 1868, v. 1, p. 6.

 Definition of the different stages of insect transformation.
- 389. [WALSH, B. D., and C. V. RILEY.] A new bark-louse on the osage orange. <Amer. Ent., September, 1868, v. 1, p. 14, fig. 8. Seasons, food-plants, characters, and figures of Lecanium maclura n. sp., [=Pulvinaria innumerabilis] and of L. accricola n. sp. [=P. innumerabilis]; characters and economic importance of the genus Aspidiotus.
- 390. [WALSH, B. D., and C. V. RILEY.] Entomological ignorance in the South. < Amer. Ent., September, 1868, v. 1, pp. 14-16.

 Reprint and criticism of article, "How to destroy the cotton-worm—a suggestion."
- 391. [WALSH, B. D., and C. V. RILEY.] Grasshoppers. <Amer. Ent., September, 1868, v. 1, p. 16.
 Ravages of Acrididæ in Illinois, Iowa, and Missouri; mention of the species committing these ravages; means against the same.
- 392. [Walsh, B. D., and C. V. RILEY.] The old and the new philosophy. <Amer. Ent., September, 1868, v. 1, p. 17.

 Defense of the practical application of science.
- 393. [Walsh, B. D., and C. V. Riley.] The animal kingdom. <Amer. Ent., September, 1868, v. 1, pp. 17-18.

 Definition of the "four grand branches of the animal kingdom;" mention

٠.

of their respective classes.

- 394. [Walsh, B. D., and C. V. Riley.] Ad interim committees. <Amer. Eut., September, 1868, v. 1, p. 18.
 - Duties of ad interim committees of horticultural societies; the American Entomologist ready to publish any scientific discoveries or observations with regard to rots or diseases of a vegetal nature.
- 395. [WALSH, B. D., and C. V. RILEY.] On our table. <Amer. Ent., September, 1868, v. 1, p. 18.
 - Notice of W. H. Edwards's "The butterflies of North America," pt. 1; of A. S. Packard, jr.'s "Guide to the study of insects," pt. 2; and of the "Canadian Entomologist," v. 1, No. 1.
- 396. [WALSH, B. D., and C. V. RILEY.] "Fire-flies." < Amer. Ent., September, 1868, v. 1, p. 19, fig. 9.
 - Answer to inquiry of W. McC.; characters and transformations of *Photinus pyralis*; figures larva, with details of structure, pupa, and imago.
- 397. [Walsh, B. D., and C. V. Riley.] A new grape root-borer. <Amer. Ent., September, 1868, v. 1, p. 19.
 - Answer to inquiry of W. D. F. Lummis; characters and ravages of an undetermined longicorn larva; food-habits of Orthosoma cylindricum [== brunneum].
- 398. [WALSH, B. D., and C. V. RILEY.] Bugs on grape-vines mistaken for chinch-bugs. <Amer. Ent., September, 1868, v. 1, p. 19.

 Answer to inquiry of F. Hecker; characters, ravages, and food-plants of Blissus leucopterus and of Piesma oinerea.
- 399. [Walsh, B. D., and C. V. Rilley.] Leaf-hoppers of the grape. Amer. Ent., September, 1868, v. 1, p. 19.

 Answer to inquiry of F. Hecker: characters and ravages of Proconia [= On-
 - Answer to inquiry of F. Hecker; characters and ravages of Proconia [= Oncometopia] undata.
- 400. [Walsh, B. D., and C. V. Riley.] Leaf bugs. <Amer. Ent., September, 1868, v. 1, p. 19.
 - Answer to inquiry of C. H. Peck; food-habits of Tingis [= Gargaphia] tilia and of T. [= Corythuca] ciliata; punctures on leaves caused by Hemiptera.
- 401. [Walsh, B. D., and C. V. Riley.] Potato bugs. < Amer. Ent., 1868, v. 1, October, pp. 21-27, fig. 10-19; November, pp. 41-49, fig. 33-48.
 - Failure of popular authors to distinguish between the different insects infesting the potato plant; brief accounts and figures of one or more stages of the following species: Gortyna nitela, Baridius [= Trichobaris] trinotata, Sphinx 5-maculata [= Protoparce celeus], Lytta [= Epicauta] vittata, L. atrata [= E. pennsylvanica], L. marginata [= E. cinerea], L. cinerea [= Macrobasis unicolor], L. murina [= M. unicolor], Lema trilineata, and Haltica [= Crepidodera] cucumeris; migrations, habits, enemies, and means against Doryphora 10-lineata; figures of all stages of D. 10-lineata and D. juncta and of numerous enemies of the former.
- 402. [Walsh, B. D., and C. V. Riley.] "Grasshoppers." Their devastations in western Iowa and the good that has resulted from them. <Amer. Ent., October, 1868, v. 1, pp. 27-28.
 - Letter of M.C. Nickerson, with comments; seeds of Vilfa vaginæflora carried by "grasshoppers."

- 403. [Walsh, B. D., and C. V. Riley.] An apple growing on a grapevine. <Amer. Ent., October, 1868, v. 1, p. 28.
 - Extract from Richmond (Va.) Whig, with criticism; a gall of Cecidomyia ritispomum mistaken for an apple growing on a grape-vine. See Nos. 332, 436, 478.
- 404. [Walsh, B. D., and C. V. Riley.] Ticks and Texas fever. <a h

Improbability that Ixodes bovis is the cause of the Texas fever of cattle.

- 405. [Walsh, B. D., and C. V. RILEY.] Scientific symbols. <Amer. Ent., October, 1868, v. 1, p. 28. Reprint: <Op. cit., November, 1869, v. 2, p. 50.
 - Explanation of the use of the signs designating the sexes.
- 406. [Walsh, B. D., and C. V. Riley.] A swarm of butterflies. <Amer. Ent., October, 1868, v. 1, pp. 28-29, figs. 20-22.
 - Record of swarms of *Danais archippus*; food-plant of larva; figures larva, chrysalis, and imago.
- 407. [WALSH, B. D., and C. V. RILEY.] The sting of the 17-year Cicada. <Amer. Ent., October, 1868, v. 1, pp. 36-37. Communications from F. W. Collins, R. Richardson, and B. Borden, on the reputed sting of Cicada [= Tibicen] septendecim and on the habits of Stisus grandis [= Sphecius speciosus].</p>
- 408. [Walsh, B. D., and C. V. Rilley.] On our table. Amer. Ent., October, 1868, v. 1, p. 37.
 Notice of "The insect world," by L. Figuier.
- 409. [WALSH, B. D., and C. V. RILEY.] Squash-bug; its change of color. <Amer. Ent., October, 1868, v. 1, p. 37.</p>
 Answer to inquiry of J. Periam; changes of color of Coreus [= Anasa] tristis during its metamorphoses.
- 410. [Walsh, B. D., and C. V. Riley.] White grub; immunity from it next year in Clinton County, Mo. <Amer. Ent., October, 1868, v. 1, p. 37.
 - Answer to inquiry of J. P. McCartney; life-habits of Lachnosterna fusca.
- 411. [WALSH, B. D., and C. V. RILEY.] Insect enemies of the Colorado potato beetle. <Amer. Ent., October, 1868, v. 1, p. 37.

 Answer to inquiry of S. H. Kriedelbaugh; identification of three insect enemies of *Doryphora* 10-lineats.
- 412. [WALSH, B. D., and C. V. RILEY.] "Harvest-bugs" in America, misnamed "jiggers." < Amer. Ent., October, 1868, v. 1, p. 38.

 Answer to inquiry of M. McKenzie; habits of parasitic Acarina; distribution of and injuries caused by Sarcopsylla penetrans.
- 413. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., October, 1868, v. 1, p. 38.
 - Answer to inquiry of J. T. Smith; food-plants of Aphis mali and of Notodonta [= Edemasia] concinna; Campoplex [= Limneria] fugitiva parasitic on larva of Euchates egle.
- 414. [WALSH, B. D., and C. V. RILEY.] Grapes cut off by the tree cricket. <Amer. Ent., October, 1868, v. 1, p. 38, fig. 30-31.

 Answer to inquiry of J. H. Tice; food-habits of *Ecanthus niveus*; deposition

of eggs; figures both sexes.

- 415. [Walsh, B. D., and C. V. Rilley.] The Buffalo tree-hopper. <Amer. Ent., October, 1868, v. 1, p. 38.
 - Answer to inquiry of P. Rickert; characters, habits, and means against Ceresa bubalus.
- 416. [Walsh, B. D., and C. V. Riley.] A scavenger mistaken for a foe. <Amer. Ent., October, 1868, v. 1, p. 38.
 - Answer to inquiry of Simmons and Tillson; characters and habits of the larva of Copris carolina.
- 417. [Walsh, B. D., and C. V. Riley.] Insect foes of the bark-louse. <Amer. Ent., October, 1868, v. 1, p. 39, fig. 32.
 - Answer to inquiry of J. Weed; habits and figures of larva and image of Chilocorus bivulnerus; food-habits and figure of Epilachna borealis.
- 418. [Walsh, B. D., and C. V. Riley.] Best works on entomology. <Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of B. M. Reynolds; mention of several important works on entomology.
- 419. [Walsh, B. D., and C. V. Riley.] Larvæ in plum-gum. < Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of P. Rickert; unknown larva (probably dipterous) found in gum of plum tree; Mycetophila sp., bred from gum of peach-tree.
- 420. [Walsh, B. D., and C. V. Riley.] Insect foes of the hop vine. <Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of H. C. Freeman; characters and habits of larva of Saturnia [= Hyperchiria] io and of Vanessa [= Grapta] interrogationis.
- 421. [Walsh, B. D., and C. V. Rilley.] Insect foe of the apple-tree borer. <Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of J. E. Walker; undetermined carabid larva preying upon larva of Saperda candida.
- 422. [Walsh, B. D., and C. V. Riley.] Oak tree caterpillars. < Amer. Ent., October, 1868, v. 1, p. 39.
 - Answer to inquiry of W. W. Daniells; characters and habits of larva of Edema albifrons.
- 423. [Walsh, B. D., and C. V. Riley.] Bark-lice on apple trees. <Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiry of J. G. Fleck; Mytilaspis pomorum destroyed by a minute mite.
- 424. [Walsh, B. D., and C. V. Riley.] The "saddle-back" caterpillar. <Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiries of E. Baxter and O. A. Kenyon; characters, habits, and urticating properties of the larva of *Empretia stimulea*.
- 425. [Walsh, B. D., and C. V. Riley.] The regal walnut caterpillar. <Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiry of M. Copley; characters and food-plants of larva of Citheronia regalis; times of metamorphosis; characters of the imago.
- 426. [Walsh, B. D., and C. V. Riley.] The Hessian fly. < Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiry of F. D. Carson; ravages of Cooldomyia destructor; late sowing as a means of avoiding the same.

- 427. [WALSH, B. D., and C. V. RILEY.] Red cedar caterpillar. < Amer. Ent., October, 1868, v. 1, p. 40.
 - Answer to inquiry of R. Peter; habits, characters, and means against Thyridopteryx ephemeræformis.
- 428. [WALSH, B. D., and C. V. RILEY.] Wheat-midge winter killed. <Amer. Ent., October, 1868, v. 1, p. 40.

 Answer to inquiry of J. P. Alexander; cause of the scarcity of Diplosis tritioi.
- 429. [Walsh, B. D., and C. V. Riley.] Grape-vine borer. <Amer. Ent., October, 1868, v. 1, p. 40.

 Answer to inquiries of A. Barter and J. H. Hogan; undetermined cerambyoid larva injurious to grape-vines.
- 430. [WALSH, B. D., and C. V. RILEY.] Popular names and scientific names. <Amer. Ent., November, 1868, v. 1, p. 49, figs. 49-51. Indefiniteness of popular names; figures Julus sp., and the larva and image of one of the Elateridæ.
- 431. [Walsh, B. D., and C. V. Riley.] Entomological ignorance in the North. < Amer. Ent., November, 1868, v. 1, pp. 50-51, figs. 52-54.
 - Confusion existing in regard to the meaning of the word locust; habits of Aorididæ and Cicadidæ; figures types of the two families and of twig with eggs of Cicada sp.
- 432. [WALSH, B. D., and C. V. RILEY.] Tit for tat. <Amer. Ent., November, 1868, v. 1, p. 52.
 Ridicule of an absurd entomological item.
- 433. [WALSH, B. D., and C. V. RILEY.] Grasshoppers. <Amer. Ent., November, 1868, v. 1, p. 53.

 Abundance and ravages of several species of Acridida and of Acheta [= Gryllus] abbreviatus in the northern central United States, and scarcity of the same in New York in 1868.
- 434. [Walsh, B. D., and C. V. RILEY.] To keep seed peas from bugs. Amer. Ent., November, 1868, v. 1, p. 53.

 Habits of and means against Bruchus pisi.
- 435. [Walsh, B. D., and C. V. Riley.] Mind how you pack insects.

 Amer. Ent., November, 1868, v. 1, p. 54.
 Care needed to avoid the introduction of noxious insects.
- 436. [Walsh, B. D., and C. V. Riley.] The apple growing on a grape vine. <Amer. Ent., November, 1868, v. 1, p. 54.

 The "vegetable phenomenon," see Nos. 332, 403, 478, proved to be a gall.
- 437. [Walsh, B. D., and C. V. RILEY.] The late exhibition of useful and destructive insects at Paris. < Amer. Ent., November, 1868, v. 1, p. 55.
 - Notice of the formation of the Société d'Insectologie Agricole at Paris, and of the second exhibition of the society.
- 438. [Walsh, B. D., and C. V. Riley.] Destroying black ants in gardens. <Amer. Ent., November, 1868, v. 1, p. 55.

 Answer to inquiry of W. S. Patten; means against Formicidæ in gardens.

- 439. [Walsh, B. D., and C. V. Riley.] Entomological quackery. <Amer. Ent., November, 1868, v. 1, p. 56.

 Reprint of "The Curculio (Iowa Homesteed, 22d July, 1868), with comments;
 - means against Conotrachelus nenuphar.
- 440. [WALSH, B. D., and C. V. RILEY.] The cruel bug-hunters. < Amer. Ent., November, 1868, v. 1, p. 56.

 Insects are not susceptible of such feelings of pain and pleasure as are felt

Insects are not susceptible of such feelings of pain and pleasure as are felt by higher animals.

- 441. [WALSH, B. D., and C. V. RILEY.] Honey bees eating grapes. <Amer. Ent., November, 1868, v. 1, p. 56.
 - Reprint of a letter by T. W., from Ohio Farmer, with comments; Apis mellifica as an enemy of sound fruit.
- 442. [Walsh, B. D., and C. V. Riley.] Twigs amputated by some unknown animal. <Amer. Ent., November, 1868, v. 1, p. 57.

 Answer to inquiry of G. Burnside; method of work of an undetermined fruit-tree pruner [= Oncideres cingulata?].
- 443. [WALSH, B. D., and C. V. RILEY.] Twigs girdled by some animal. <Amer. Ent., November, 1868, v. 1, p. 57.

 Answer to inquiries of Judge Brown and P. Earle; method of work of an undetermined fruit-tree pruner [= Oncideres cingulata †]; trees affected.
- 444. [WALSH, B. D., and C. V. RILEY.] Hop-vine caterpillar. < Amer. Ent., November, 1868, v. 1, p. 57.</p>
 Answer to inquiry of H. J. Dunlap; larva of Grapta interrogationis feeding on hop-vines.
- 445. [WALSH, B. D., and C. V. RILEY.] "Galls" on leaves of soft maple.

 <Amer. Ent., November, 1868, v. 1, p. 57.

 Answer to inquiry of A. L. Child; characters of undescribed mite-galls and
- their architects; mode of formation of leaf-galls by mites.

 446. [Walsh, B. D., and C. V. Riley.] Unsightly galls on the cottonwood. <Amer. Ent., November, 1868, v. 1, p. 57.

Answer to inquiry of A. L. Child; character of galls made by *Pemphigus va-gabondus* and *P. populicaulis*; their effects upon the cottonwood and means against them.

- 447. [Walsh, B. D., and C. V. Riley.] Insects named. <Amer. Ent.,
 November, 1868, v. 1, p. 57.
 - Answer to inquiry of E. Baxter; Gordius aquatious a host in Orchelimum gracile; Cetonia [= Euphoria] inda destructive to the peach.
- 448. [Walsh, B. D., and C. V. Riley.] The stick-bug. <Amer. Ent., November, 1868, v. 1, p. 58. Reprint: <Can. Farmer, 15 January, 1870.
 - Description, vernacular names, and habits of Spectrum [= Diapheromera] femorata.
- 449. [Walsh, B. D., and C. V. Riley.] Woolly lice on the beech. <Amer. Ent., November, 1868, v. 1, p. 58.
 - Answer to inquiry of F. H. Guiwits; habits and food-plants of Pemphigus imbricator.

- 450. [Walsh, B. D., and C. V. Riley.] The sheep-bot or head-maggot. <Amer. Ent., November, 1868, v. 1, p. 58.
 - Answer to inquiry of R. W. Scott; *Œstrus ovis* ordinarily oviparous; sometimes the eggs hatch prematurely inside the body.
- 451. [Walsh, B. D., and C. V. RILEY.] Leaf-miners of the locust. <Amer. Ent., November, 1868, v. 1, p. 58.
 - Answer to inquiry of R. W. Scott; characters, habits, ravages, and means against Hispa scatellaris [= Odontota dorsalis].
- 452. [Walsh, B. D., and C. V. Riley.] Leaf-hoppers on celery. <Amer. Ent., November, 1868, v. 1, p. 58.
 - Answer to inquiry of R. Parnell; mention of two undescribed Tettigonida infesting early celery.
- 453. [Walsh, B. D., and C. V. Riley.] Parsnip caterpillars.—Scorpions. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of F. Brewer; food-plants of larva of Papilio asterias; habits of the imago; habits and sting of Buthus carolinianus; its occurrence in Missouri.
- 454. [Walsh, B. D., and C. V. Riley.] Noxious insects named. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of J. H. Parsons; larval characters and food-plants of Hyphantria textor [= cunea], Notodonta [= Edemasia] concinna, Arctia [= Spilosoma] virginica, Carpocapsa pomonella, and of Trypeta pomonella.
- 455. [WALSH, B. D., and C. V. RILEY.] Cut-worms destroying recently sown wheat. < Amer. Ent., November, 1868, v. 1, p. 59. Answer to inquiry of T. R. Allen; characters, habits, and means against undetermined Noctuidæ attacking wheat.</p>
- - Answer to inquiry of I. Hicks; Podieus spinosus as an enemy of Hyphantria textor [= cunea].
- 457. [Walsh, B. D., and C. V. Riley.] The preying Mantis, alias Devil's riding horse, etc. <Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of A. Pettit; habits of Mantis [=Phasmomantis] carolina; egg-mass of the same.
- 458. [Walsh, B. D., and C. V. Riley.] The pigeon Tremex. < Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of F. Brewer; characters of Tremex columba; food-habits and parasites of its larva.
- 459. [Walsh, B. D., and C. V. Rilley.] Flesh-worms. < Amer. Ent., November, 1868, v. 1, p. 59.
 - Answer to inquiry of W. J. Stuart; larval habits of an undetermined muscid-
- 460. [Walsh, B. D., and C. V. Riley.] Fall web-worm on hickory. <Amer. Eut., November, 1868, v. 1, p. 59.
 - Answer to inquiry of W. W. Daniels; transformation of Hyphantria textor [=ounea]; characters of its larva and imago.

- 461. [WALSH, B. D., and C. V. RILEY.] Butterflies named. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of W. W. Butterfield; characters of Collas philodice, Vancesa [= Grapta] comma, and Danais archippus.
- 462. [WALSH, B. D., and C. V. RILEY.] Museum pest. < Amer. Ent., November, 1868, v. 1, p. 60.

Answer to inquiry of T. W. Holt, jr.; means against Dermestidæ.

- 463. [WALSH, B. D., and C. V. RILEY.] Insects named. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of J. Weed; enemies of plant-lice and bark-lice; foodplant and characters of the larva of *Papilio turnus*.
- 464. [WALSH, B. D., and C. V. RILEY.] Grapes spoiled by something. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of J. Wood; grapes injured from some unknown cause; juice from wounded grapes as food for moths.
- 465. [Walsh, B. D., and C. V. RILEY.] White-pine worms. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of J. T. Little; food-plants of larva of Lophyrus abbotii; transformations of the same.
- 466. [Walsh, B. D., and C. V. Riley.] The Tarantula of Texas. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of J. Bell; manner in which Pompilus [= Pepsis] formosa prepares Mygale hentzii for food for its larva.
- 467. [Walsh, B. D., and C. V. Riley.] Insect foes of the apple-tree. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of G. C. Brackett; characters of larva of Carpocapsa pomonella; means against Datana ministra and Pemphigus pyri [=Schizoneura lanigera].
- 468. [Walsh, B. D., and C. V. Riley.] Grape-leaf folders eaten by spiders. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of A. A. Hilliard; larva of *Desmia maculalis* destroyed by a spider.
- 469. [WALSH, B. D., and C. V. RILEY.] Caterpillar of the Troilus butterfly. <Amer. Ent., November, 1868, v. 1, p. 60.

 Answer to inquiry of A. S. Fuller; food-habits of larva of Papillo troilus.
- 470. [Walsh, B. D., and C. V. Riley.] Potato beetles. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of P. H. Foster; identification of *Epicauta vittata* and *Macrobasis unicolor* as enemies of the potato.
- 471. [Walsh, B. D., and C. V. Riley.] Holes round the roots of young ash trees in the nursery. <Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of I. Hicks; characters and food-habits of the larva of Xyloryctes satyrus.
- 472. [Walsh, B. D., and C. V. RILEY.] Locust borers. < Amer. Ent., November, 1868, v. 1, p. 60.
 - Answer to inquiry of U. Bruner; destructiveness of the larva of Clytus [= Cyllene] robiniæ.

- 473. [WALSH, B. D., and C. V. RILEY.] The hellgrammite fly. <Amer. Ent., December, 1868, v. 1, pp. 61-62, figs. 56-57.
 - Habits of Corydalus cornutus; figures the larva, pupa, and imago; description and figure of the eggs of Belostoma americanum mistaken for those of Corydalus.
- 474. [Walsh, B. D., and C. V. Rilley.] The periodical Cicada. < Amer. Ent., December, 1868, v. 1, pp. 63-72, figs. 58-64. Extract: < Op. cit., June, 1869, v. 1, p. 202.
 - Characterization of the 13-year brood of Cicada as a new species, C = Tibicen tredecim; dimorphism of the same and of C = T.] septendecim; seasons, natural history, transformations, enemies, sting, and injuries of these species; chronological history of their several known broods; figures the several stages of C = T.] septendecim, the towers made by the pupa and twigs with eggs.
- 475. [Walsh, B. D., and C. V. Riley.] The hateful or Colorado grass-hopper, *Caloptenus spretus*, Uhler and Walsh. <Amer. Ent., December, 1868, v. 1, pp. 73-76, fig. 65.
 - Comparative characters and figures of Caloptenus spretus and C. femur-rubrum; native habitat, migrations, distribution, and ravages of the former; harmlessness of its second generation in the Mississippi Valley as compared with the first generation; improbability that it will ever become a permanent resident in that region.
- 476. [WALSH, B. D., and C. V. RILEY.] The twig-girdler, Oncideres cingulatus, Say. <Amer. Ent., December, 1868, v. 1, pp. 76-77, fig. 66.
 - Oviposition and food habits of *Oncideres oingulata*; figure of image and of amputated twig.
- 477. [WALSH, B. D., and C. V. RILEY.] An Ichneumon fly mistaken for a wasp. <Amer. Ent., December, 1868, v. 1, p. 77.

 Review of paper in Christian Advocate on "The stiletto wasp"; Pimpla [= Thalessa] atrata probably mistaken for a wasp.
- 478. [Walsh, B. D., and C. V. Riley.] A plant growing out of an insect. <Amer. Ent., December, 1868, v. 1, p. 77.
 - See Nos. 332, 403, 436; extract from Sedalia Press, with comments; larvæ of Lacknosterna fusca infested with Cordyceps; larvæ supposed to have eaten poisonous seed, which has germinated after killing the larvæ; sowing the seed as a means against the larvæ.
- 479. [Walsh, B. D., and C. V. Riley.] On our table. <Amer. Ent., December, 1868, v. 1; p. 78.
 - Notices of The butterflies of North America by W. H. Edwards.—The Canadian Entomologist.—Guide to the study of insects by A. S. Packard, jr.
- 480. [WALSH, B. D., and C. V. RILEY.] Eggs of the white-marked tussock moth. <Amer. Ent., December, 1868, v. 1, p. 79, fig. 67.

 Answer to inquiry of J. M. Hannah; characters of egg and larva of Orgyia loucostigma; sexual differences; parasites; figure of the larva.
- 481. [Walsh, B. D., and C. V. Riley.] Current borers. < Amer. Ent., December, 1868, v. 1, p. 79.
 - \nswer to inquiry of B. N. McKinstry; means against Trochilium [= Egeria] tipuliformis, T. [= Aloathæ] caudatum, and Psenocerus supernotatus.

- 482. [Walsh, B. D., and C. V. Riley.] Insect infesting grape seed. < Amer. Ent., December, 1868, v. 1, p. 79, fig. 68. Answer to inquiry of A. S. Fuller; characters and figure of larva of Isosoma
 - vitis infesting grape seed.
- 483. [Walsh, B. D., and C. V. Riley.] Museum pests again. < Amer. Ent., December, 1868, v. 1, p. 79. Answer to inquiry of J. Huggins; means against Anthreni in collections of
- natural history. 484. [Walsh, B. D., and C. V. Riley.] Grape-vine leaf-hoppers.
- <Amer. Ent., December, 1868, v. 1, p. 79. Answer to inquiry of P. C. Holmes; means against grape-vine leaf-hoppers.
- 485. [WALSH, B. D., and C. V. RILEY.] Apple-tree borer. < Amer. Ent., December, 1868, v. 1, p. 80, fig. 69. Answer to inquiries of J. T. Zimmerman, C. H. Roberts, and C. R. Babbitt:
- 486. [WALSH, B. D., and C. V. RILEY.] The murky ground beetle. < Amer. Ent., December, 1868, v. 1, p. 80.

habits, ravages, and figure of Bostrichus [= Amphicerus] bicaudatus.

- Answer to inquiry of E. T. Dale; fcod-habits and bombardier discharges of Harpalus caliginosus.
- 487. [WALSH, B. D., and C. V. RILEY.] Curculio and bark-lice. <Amer. Ent., December, 1868, v. 1, p. 80. Answer to inquiry of D. A. Compton; hibernation of Constrachelus nenuphar; soft soap as a means against Coccidæ.
- 488. [Walsh, B. D., and C. V. Riley.] Insects to be named. < Amer. Ent., December, 1868, v. 1, p. 80. Answer to inquiry of E. B. Beach; identification of several lepidopterous larvæ; food-habits of Dryocampa stigma and D. pellucida [= virginiensis].
- 489. [WALSH, B. D., and C. V. RILEY.] Young pecan trees girdled. Amer. Ent., December, 1868, v. 1, p. 80. Answer to inquiry of Mann and Redmond; means against Oncideres cinquiata.
- 490. [WALSH, B. D., and C. V. RILEY.] Insects named. < Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of E. T. Dale; identification of several Coleoptera; supposed food-habits of Tragidion fulvipenne; character of cocoons of two species of Microgaster infesting Protoparce celeus.
- 491. [WALSH, B. D., and C. V. RILEY.] Eggs of true bugs. < Amer. Ent., December, 1868, v. 1, p. 80. Answer to inquiry of S. C. Thornton; characters of eggs of Prionidus cristatus. See Amer. Ent., v. 1, pp. 96 and 187.
- 492. [WALSH, B. D., and C. V. RILEY.] Museum pest once more. Amer. Ent., December, 1868, v. 1, p. 80. Answer to inquiry of G. M. L.; larvæ of Anthrenus musæorum injuring whale-

bone, woolen goods, etc.

493. [WALSH, B. D., and C. V. BILEY.] The Tarantula of Texas again. < Amer. Ent., December, 1868, v. 1, p. 80. Answer to inquiry of C. Peabody; distribution of Mygale kentzii in Missouri.

- 494. [Walsh, B. D., and C. V. RILEY.] The Hessian fly on seed wheat. < Amer. Ent., December, 1868, v. 1, p. 80.
 - Answer to inquiry of U. Scott; broads and times of transformation of Cecidomyia destructor.
- 495. [Walsh, B. D., and C. V. Riley.] The apple-root plant-louse. Eriosoma (Pemphigus) pyri, Fitch. <Amer. Ent., January, 1869, v. 1, pp. 81-84, figs. 70-72.
 - Habits, ravages, description, enemies, and parasites of, and means against, Eriosoma pyri [=Schizoneura lanigera]; figures injured root, larva, and adult with details of structure; figures adult plant-lones found on cottonwood; description and figure of larva, puparium, and image of Pipiza radicum n. sp.
- 496. [Walsh, B. D., and C. V. Billey.] Ants' nests in gardens. Amer. Ent., January, 1869, v. 1, p. 84.
 Means against ants.
- 497. [Walsh, B. D., and C. V. Riley.] The parasites of the human animal. <Amer. Ent., January, 1869, v. 1, pp. 84-88, figs. 73-74.
 - Brief accounts of Pediculus humanus [= vestimenti], P. cervicalis [= capitis], P. [= Pthirius] pubis, Œstrus [= Dermatobia] hominis, Pulex irritans, P. [= Sarcopsylla] penetrans, Acanthia lectularia, Conorhinus sanguisuga, and Acarus [= Sarcoptes] scabici; figures and habits of Reduvius [= Opsicatus] personatus and of Pirates [= Rasahus] biguitatus; figure of Conorhinus sanguisuga; classificatory relations of Pediculina and Mallophaga.
- 498. [Walsh, B. D., and C. V. Riley.] The coffee borer. < Amer. Ent., January, 1869, v. 1, p. 88.
 Unnamed coffee borer injurious to coffee trees in Madras.
- 499. [WALSH, B. D., and C. V. RILEY.] Strawberry worms. <Amer. Ent., January, 1869, v. 1, pp. 89-91, figs. 75-76.</p>
 Description, natural history, ravages, means against, and figures of larva and image of Anchylopera [= Phoxopteris] fragaria, n. sp.; figures all stages of Emphytus [= Harpiphorus] maculatus; description of its larva and pupa, its natural history, and means against its ravages.
- 500. [WALSH, B. D., and C. V. RILEY.] Fungoid growths. <Amer. Ent., January, 1869, v. 1, pp. 91-92.</p>
 Mention of instances of the growth of fungi on living plants and animals; letter of S. H. Y. Early on the occurrence of fungoid growths on the larvæ of Lachnosterna fusca.
- 501. [WALSH, B. D., and C. V. RILEY.] Plums for the million. <Amer. Ent., January, 1869, v. 1, pp. 92-93.

 Means against Conotrachelus nonuphar; notes on varieties of plum exempt
- 502. [Walsh, B. D., and C. V. Rilley.] The "Colorado grasshopper." <Amer. Ent., January, 1869, v. 1, pp. 95-96.

from the attacks of the same.

Answer to inquiries of W. N. Byers and V. Devinny; specific names indicating particular districts can not be changed because the insect is found in other regions.

- 503. [Walsh, B. D., and C. V. Rilley.] Universal remedies. < Amer. Ent., January, 1869, v. 1, p. 97.
 - Criticism of an advertisement of "Best's patent fruit tree and vine invigorator."
- 504. [Walsh, B. D., and C. V. Biley.] Complimentary. <Amer. Ent., January, 1869, v. 1, p. 98.

 Notices of several notices of the American Entomologist.
- 505. [Walsh, B. D., and C. V. Riley.] Paper-makers. < Amer. Ent., January, 1869, v. 1, p. 98.
 - Comparison of the manufacture of paper by man and by the Vespida.
- 506. [Walsh, B. D., and C. V. RILEY.] On our table. <Amer. Ent., January, 1869, v. 1, p. 98.
 - Notices of L'Insectologie agricole-Cecil's books of natural history.
- 507. [Walsh, B. D., and C. V. Riley.] Look out for the eggs of the apple-tree plant-louse. <Amer. Ent., January, 1869, v. 1, p. 99. Abundance of eggs of Aphis mall in winter of 1868-'69, in Missouri and Illinois; means against the same.
- 508. [WALSH, B. D., and C. V. RILEY.] Greenhouse plants [= pests].

 Amer.Ent., January, 1869, v. 1, p. 99.

 Means against Aphidida, Coccida, and Totranyohus telarius.
- 509. [Walsh, B. D., and C. V. RILEY.] How to hatch pupe. <Amer. Ent., January, 1869, v. 1, p. 99.
 - Answer to inquiry of D. P. Smith; methods of raising insects.
- 510. [Walsh, B. D., and C. V. Rilley.] Drug-store pests. < Amer. Ent., January, 1869, v. 1, p. 99.
 - Answer to inquiry of J. M. Good; characters of larva of undertermined ptinid; food-habits of *Plinus brunneus*; food-habits, characters, and means against *Calandra orysæ* and *C. granaria*. (See No. 551.)
- 511. [WALSH, B. D., and C. V. RILEY.] Apple-tree worms. <Amer. Ent., January, 1869, v. 1, p. 99.

 Answer to inquiry of J. J. Thomas; habits, food-plants, and means against
- Physita nebulo [= Acrobasis indiginella].
 512. [WALSH, B. D., and C. V. RILEY.] Crane-fly larvæ. Amer. Ent.,
 January, 1869, v. 1, p. 100.
 - Answer to inquiries of R. D. Alexander and Hickman; habits of Tipula sp.; food-habits of T. trivittata.
- 513. [Walsh, B. D., and C. V. Riley.] Punctured grape canes. <Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiry of B. L. Kingsbury; description of punctures in grape canes probably caused by *Orocheris sattator*; means against tree-crickets.
- 514. [WALSH, B. D., and C. V. RILEY.] Insects to be named. <Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiry of W. R. Marine; identification of several insects found in apple and peach nurseries.
- 515. [WALSH, B. D., and C. V. BILEY.] Supposed cause of yellows in peach trees. <Amer. Ent., January, 1869, v. 1, p. 100, fig. 77.

 Answer to inquiry of C. H. Roberts; yellows of peach-trees a vegetable disease; figure of healthy and diseased limb; affected trees more subject to the attacks of insects than healthy trees.

- 516. [Walsh, B. D., and C. V. Riley.] Eggs of katydid. < Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiry of A. A. Hilliard; character of the eggs of Platyphyllum [= Cyrtophyllus] concavus.
- 517. [Walsh, B. D., and C. V. Riley.] Apple-twig borer. < Amer. Ent., January, 1869, v. 1, p. 100.
 - Answer to inquiry of J. T. Zimmerman; ravages of Bostrichus [=Amphicerus] bioaudatus.
- 518. [Walsh, B. D., and C. V. Riley.] Galls and their architects. <Amer. Ent., February, 1869, v. 1, pp. 101-110, figs. 78-90.
 - Definition, classification, and variation of galls; accounts of some galls made by Cynipidæ, Cecidomyidæ, and Aphididæ and of the gall-makers; natural history, descriptions, and figures of Cynips [= Amphibolips] q.-spongifica, C. [= A.] q.-inants, C. [= A.] q.-prunus n. sp., Cecidomyia s.-strobiloides, C. s.-brassicoides, C. v.-pomum n. sp., C. v.-coryloides n. sp., Pemphigus ragabundus, P. rhois and Colopha ulmicola; description of the image of Cynips q.-prunus and of Pemphigus ulmifusus n. sp., and of the larvæ of the new species of Cecidomyia; dimorphism of gall-makers; presence of inquilines and parasites in galls. (See No. 821.)
- 519. [Walsh, B. D., and C. V. Riley.] The bogus Colorado potatobug, Doryphora juncta, Germar. < Amer. Ent., February, 1869, v. 1, p. 110.

Food-habits of D. juncta.

- 520. [Walsh, B. D., and C. V. RILEY.] Ants and aphides. <Amer. Ent., February, 1869, v. 1, p. 110.</p>
 Inquiry of W. Batchelor, with answer; secretion of honey-dew by Aphidida
 - and harvesting of the same by ants.
- 521. [WALSH, B. D., and C. V. RILEY.] The Tarantula of Texas, Mygale hentzii, Girard. <Amer. Ent., February, 1869, v. 1, p. 11, fig. 91.
 - Figure of Mygale kentzii: habits of Pempilus [= Pepsis] formosa; quotes from G. Lincecum's "The Tarantula killers of Texas" (Amer. Nat., v. 1, pp. 137-141).
- 522. [Walsh, B. D., and C. V. RILEY.] The melancholy chafer. <Amer. Ent., February, 1869, v. 1, p. 111.

 Ravages and figure of the image of Euphoria melancholica.
- 523. [Walsh, B. D., and C. V. Riley.] Apple worms (Carpocapsa pomonella, Linn.). < Amer. Ent., February, 1869, v. 1, pp. 112-114, fig. 93.
 - Natural history, ravages, and means against Carpocapsa pomonella; figures injured apple, larva, pupa, and imago.
- 524. [Walsh, B. D., and C. V. Riley.] The asparagus beetle (Crioceris asparagi, Linn.). < Δmer. Ent., February, 1869, v. 1, pp. 114-115, fig. 94.
 - Introduction into the United States; natural history, ravages, parasites of, and means against *Crioceris asparagi*; figure of eggs, larvw, and image of the same.

- 525. [Walsh, B. D., and C. V. Riley.] A popular delusion. <Amer. Ent., February, 1869, v. 1, p. 116.
 - No insect passes through all the stages of its growth within one day; life history of *Ephemeridæ*.
- 526. [Walsh, B. D., and C. V. Riley.] The squirrel bot. <Amer. Ent., February, 1869, v. 1, v. 117.
 - Comments on paper by S. S. Rathvon; emasculation of the striped squirrel by Cuterebra buccata.
- 527. [Walsh, B. D., and C. V. RILEY.] [Periodical Cicada.] < Amer. Ent., February, 1869, v. 1, p. 117.
 - Comments on letter of R. H. Warder; oviposition of Cicada [= Tibicen] septendecim in evergreens.
- 528. [WALSH, B. D., and C. V. RILEY.] Transformations of insects.

 Amer. Ent., February, 1869, v. 1, p. 118.
 Brief statement of the stages of growth of insects.
- 529. [Walsh, B. D., and C. V. RILEY.] Do bees injure raspberries? <Amer. Ent., February, 1869, v. 1, p. 118.
 - Comments on discussion before New York Fruit Growers' Club as to the effects of bees on flowers and fruits.
- 530. [WALSH, B. D., and C. V. RILEY.] Best's fruit-tree invigorator again. <Amer. Ent., February, 1869, v. 1, p. 119.</p>
 Notice of invention of a new "invigorator" by B. Best.
- 531. [Walsh, B. D., and C. V. Riley.] The insect extinguisher, by Joseph Treat, N. J. < Amer. Ent., February, 1869, v. 1, p. 119. Critical review of pamphlet by J. Treat.
- 532. [Walsh, B. D., and C. V. Riley.] More good words. <Amer. Ent., February, 1869, v. 1, p. 119.

 Notice of favorable notices of the American Entomologist.
- 533. [Walsh, B. D., and C. V. Riley.] Lacewing fly. < Amer. Ent.,

 February, 1869, v. 1, p. 119.

 Answer to inquiry of J. Huggins; characters of Chrysops sp.; its hibernation
- as a pupa and imago.

 534. [WALSH, B. D., and C. V. RILEY.] Gigantic water-bug. < Amer. Ent., February, 1869, v. 1, p. 119.
 - Answer to inquiry of S. E. Munford; food-habits of Belostoma grandis [= americanum].
- 535. [Walsh, B. D., and C. V. Rilley.] The white-marked tussock moth again. < Amer. Ent., February, 1869, v. 1, p. 120.

 Approx to inquire of A. S. Fellow, mortion of govern perceites attacking
 - Answer to inquiry of A. S. Fuller; mention of seven parasites attacking Orgyia leucostigma.
- 536. [Walsh, B. D., and C. V. Riley.] Grasshopper eggs—will they hatch? <Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of C. J. Jones; frosts not likely to prevent the hatching of the eggs of Caloptenus spretus.
- 537. [Walsh, B. D., and C. V. Riley.] Injured apple-trees. < Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of G. C. Broadhead; means against Chrysobothris femorata.

- 538. [Walsh, B. D., and C. V. Rilley.] Bag worms. < Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of W. W. Butterfield; means against Thyridopteryx ephemeræformis.
- 539. [WALSH, B. D., and C. V. RILEY.] Eggs of the apple-tree plantlouse. <Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of M. W. Seaman; effect of frosts upon the eggs and larve of Aphis mall.
- 540. [Walsh, B. D., and C. V. Riley.] Insects named. <Amer Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of J. B. Merwin; identification of insects sent; characters of Vancesea antiopa and Cynthia [=Pyrameis] atalanta.
- 541. [WALSH, B. D., and C. V. RILEY.] White grubs. < Amer. Ent., February, 1869, v. 1, p. 120.
 - Answer to inquiry of W. C. Holmes; undetermined white-grub injuring grass and osage orange.
- 542. [WALSH, B. D., and C. V. RILEY.] The polyphemus moth, Attacus polyphemus, Linn. <Amer. Ent., March, 1869, v. 1, pp. 121-122, fig. 95.
 - Attaons [= Telea] polyphemus, A. cynthia, and Bombyx [= Sericaria] mori as silk-producers; foed-plants, description of larva, habits, and seasons of A. [= T.] polyphemus; figure of the imago; ravages of A. cecropia; antennæ mistaken for wings; mention of Lepidoptera having fissured wings.
- 543. [WALSH, B. D., and C. V. RILEY.] Wasps and their habits. <Amer. Ent., March, 1869, v. 1, pp. 122-143, figs. 96-112.
 - Structure, classification, and habits of North American predatory Hymenoptera; comparative characters of the digger and true wasps; figure to show the folding of the wing in true wasps; habits and figures of images of Chlorion coruleum, Bembex fasciata, Sphex ichneumonea, Ammophila pictipennis, Pepsis formosa, Stizus grandis [= Sphecius speciosus], S. [=S.] speciosus, Pelopœus lunatus [= cementarius], Agenia bombyoina, Trypoxylon albitarse, Ceropales rufiventris, Eumenes fraternus, Vespa maculata, and Polistes rubiginosus; figure of image of Cryptus [= Linoceras] junceus and of several nests of wasps. (See No. 375.)
- 544. [Walsh, B. D., and C. V. Riley.] Do toads eat worker bees? <Amer. Ent., March, 1869, v. 1, p. 144.
 - Reprint and review of article by C. Dadant, and of comments of editors of American Bee Journal; usefulness of toads.
- 545. [Walsh, B. D., and C. V. Riley.] Best's invigorator once again. Amer. Ent., March, 1869, v. 1, p. 145.
 - Condemnation of B. Best's patent fruit tree and vine invigorator.
- 546. [WALSH, B. D., and C. V. RILEY.] On our table. <Amer. Ent., March, 1869, v. 1, p. 146.

 Notices of Nos. 1 and 2 of Le Naturaliste Canadien.
- 547. [Walsh, B. D., and C. V. Riley.] Sugar-tree borer. <Δmer. Ent., March, 1869, v. 1, p. 146.
 - Answer to inquiry of E. Simms; characters and means against Arkopalus [=Plagionotus] speciosus.

- 548. [WALSH, B. D., and C. V. RILEY.] Swellings on apple scions. <Amer. Ent., March, 1869, v. 1, p. 146.

 Answer to inquiry of W. Colwell; dipterous galls on apple scions and basewood twigs.
- 549. [WALSH, B. D., and C. V. RILEY.] Gas-waste vs. Curculio. <Amer. Ent., March, 1869, v. 1, p. 147.

 Answer to inquiry of H.; merits of gas-waste as an insect destroyer.
- 550. [Walsh, B. D., and C. V. Riley.] Eggs of the apple-tree plant-louse again. < Amer. Ent., March, 1869, v. 1, p. 147.

 Answers to inquiries of C. Williams and W. L. French; means against Aphie mali
- 551. [WALSH, B. D., and C. V. RILEY.] Drug-store pests. <Amer. Ent., March, 1869, v. 1, p. 147.</p>
 Answer to inquiry of J. M. Good; food-habits of Lasioderma servicorns. See No. 510.
- 552. [WALSH, B. D., and C. V. RILEY.] Small galls and minings on apple-twigs. <Amer. Ent., March, 1869, v. 1, p. 147.

 Answer to inquiry of O. O. A. Gardner; characters of undetermined galls and borings in twigs of apple-trees.
- 553. [WALSH, B. D., and C. V. RILEY.] Stinking bugs. < Amer. Ent., March, 1869, v. 1, p. 147.</p>
 Answer to inquiry of C. L. Janney; habits of an undescribed Brachyrhynchus.
- 554. [Walsh, B. D., and C. V. Riley.] Eggs in peach-twigs. < Amer. Ent., March, 1869, v. 1, p. 147.

 Answer to inquiry of G. Fisher; character of eggs of *Ecanthus nivous*.
- 555. [WALSH, B. D., and C. V. BILEY.] Flat-headed apple-tree borer. <Amer. Ent., March, 1869, v. 1, p. 147.

 Answer to inquiry of B. F. Mudge; means against Chrysobothris femorata.
- 556. [WALSH, B. D., and C. V. RILEY.] A bundle of entomological queries. < Amer. Ent., March, 1869, v. 1, p. 148.</p>
 Answer to inquiry of W. W. Butterfield; mentions several desirable entomological books.
- 557. [WALSH, B. D., and C. V. RILEY.] Grape-berry moth. < Amer. Ent., March, 1869, v. 1, p. 148.
 Answer to inquiry of M. C. Read; means against Penthina vitivorana [= Eudemis botrana].
- 558. [WALSH, B. D., and C. V. RILEY.] Hairy caterpillar. < Amer. Ent., March, 1869, v. 1, p. 148.</p>
 Answer to inquiry of T. S. Gold; characters of the larva and image of Arctia [= Pyrrharctia] isabella.
- 559. [WALSH, B. D., and C. V. RILEY.] Borer in plum-twigs. <Amer. Ent., March, 1869, v. 1, p. 148.

 Answer to inquiry of W. Caldwell; undetermined borer, allied to the oakpruner, in plum-twigs. See No. 606.

- 560. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., March, 1869, v. 1, p. 148.
 - Answer to inquiry of W. W. Butterfield; identification of insects sent; comparative characters of Arhopalus [= Cyllene] pictus and A. [= C.] robiniæ; variation in Clytus [= Neoclytus] capræa; generic characters of Arhopalus and Clytus. (See No. 582.)
- 561. [Walsh, B. D., and C. V. Riley.] Pear-root borer. < Amer. Ent., March, 1869, v. 1, p. 148.
 - Answer to inquiry of P. Earle; larva of *Prionus laticollis* injurious to the pear and grape.
- 562. [Walsh, B. D., and C. V. Rilley.] Insects named. < Amer. Ent., March, 1869, v. 1, p. 148.
 - Answer to inquiry of M. H. Boye; Geyllotalpa longipennis [= columbia] a dimorphic form of G. borealis.
- 563. [Walsh, B. D., and C. V. Rilley.] The joint-worm (*Isosoma hordei* Harris). <Amer. Ent., April, 1869, v. 1, pp. 149-158, figs. 113-118.
 - Natural history, variation, parasites, ravages, and means against Isosoma horder; unity of habits in insects; generic classification of the joint-worm fly; figures male and female imago; figures 3 and 2 imagos of Semiotellus chalcidiphagus n. sp., Eurytoma sp., and Decatoma sp., and of 2 imago of Antigaster [= Eupelmus] mirabilis.
- 564. [Walsh, B. D., and C. V. Riley.] The wavy-striped flea-beetle. (Haltica [Phyllotreta] striclata Illiger). <Amer. Ent., April, 1869, v. 1, pp. 158-159, fig. 119.
 - Description, habits, and ravages of *Phyllotreta striolata* [=vittata]; figures larva, pupa, and imago of the same; quotes from H. Shimer's "The wavy-striped flea-beatle" (Amer. Nat., December, 1868, v. 2, pp. 514-517).
- 565. [Walsh, B. D., and C. V. Riley.] Concerning certain smart bugs. <Amer. Ent., April, 1869, v. 1, p. 160.
 - Preferences of Phylloxera vitifolia [=vastatrix] and Macrodactylus subspinosus for certain varieties of grape-vines, of Doryphora 10-lineata and Lema trilineata for certain varieties of potato-vines, and of Mytilaspis pomicorticis [=pomorum] and Carpocapsa pomonella for certain varieties of apple.
- 566. [Walsh, B. D., and C. V. Riley.] Curculio remedies. < Amer. Ent., April, 1869, v. 1, p. 161.
 - Comments on essay of L. C. Francis on the plum; coal oil not effective against Conotrachelus nenuphar.
- 567. [WALSE, B. D., and C. V. RILEY.] How great wits jump together. <Amer. Ent., April, 1869, v. 1, p. 161.

 Stopping holes with hard soap ineffective against borers.
- 568. [Walsh, B. D., and C. V. Riley.] On our table. <Amer. Ent., April, 1869, v. 1, p. 165.
 - Notices of Harris's Insects injurious to Vegetation, L'Insectologie agricole, and several other works not entomological.

- 569. [Walsh, B. D., and C. V. Riley.] Eggs of the oblong-winged katydid. <Amer. Ent., April, 1869, v. 1, p. 166, fig. 120.

 Appear to inquiring of T. A. Thon, I. J. Pice, R. J. Campbell, H. Cheener.
 - Answer to inquiries of T. A. Thop, J. L. Rice, B. J. Campbell, H. Cheeney, and A. McMoore; eggs of Phylloptera [= Amblycorypha] oblongifolia compared with those of Platyphyllum [= Cyrtophyllus] concavum; figures of the eggs of A. oblongifolia.
- 570. [Walsh, B. D., and C. V. Riley.] Mossy rose gall. <Amer. Ent., April, 1869, v. 1, p. 166.
 - Answer to inquiry of F. W. Collins; characters of Rhodites rose and its gall.
- 571. [Walsh, B. D., and C. V. Riley.] Caterpillars on lombardy poplars. <Amer. Ent., April, 1869, v. 1, p. 166.
 - Answer to inquiry of J. F. Gurley; characters of larva, habits, food-plants, and parasites of Acronycta acericola [= americana].
- 572. [Walsh, B. D., and C. V. Riley.] Museum pests. <Amer. Ent., April, 1269, v. 1, p. 166, fig. 121.
 - Answer to inquiries of F. W. Hoit, jr., G. M. L., and J. Huggins; figures larva, pupa, and image of Anthrenus sp.; A. varius compared with A. muse-orum.
- 573. [Walsh, B. D., and C. V. Riley.] Cockroach eggs. < Amer. Ent., April, 1869, v. 1, p. 166.
 - Answer to inquiry of H. C. Freeman; character of the egg-cases and adult of *Platamodes unicolor*; occurrence of *Ectobia germanica* in Illinois; ravages of species of *Blattidæ*.
- 574. [Walsh, B. D., and C. V. Riley.] Apple-tree leaf-crumplers. <Amer. Ent., April, 1869, v. 1, p. 166.
 - Answer to inquiry of J. F. Jones; result of the work of the larve of Physica nebulo [= Acrobasis indiginella] on the growth of apple-trees.
- 575. [Walsh, B. D., and C. V. Riley.] A most precious bug. <Amer. Ent., April, 1869, v. 1, p. 167.
 - Answer to inquiry of E. S. Holmes; characters and variety of Hylecatus lugubris; destructiveness of Lymexylon navale; use of the maxillary palpi in the 3 3 of Lymexylida.
- 576. [Walsh, B. D., and C. V. RILEY.] Blackberry-cane borers. <Amer. Eut., April, 1869, v. 1, p. 167.
 - Answer to inquiry of C. Parry; characters and habits of larva of unnamed borer [= Bembècia marginata]; plants affected by species of Ægeriadæ.
- 577. [Walsh, B. D., and C. V. Rilley.] Rows of eggs in pear-twigs. <Amer. Ent., April, 1869, v. 1, p. 167.
- Answer to inquiry of H. C. Freeman; characters of unknown eggs found in pear-twigs.
- 578. [Walsh, B. D., and C. V. Riley.] An orchard giving out. <Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. M. Clemens; habits and means against Saperda bivittata [= candida] and Chrysobothris femorata.
- 579. [Walsh, B. D., and C. V. Riley.] Plant-louse eggs on apple and mountain ash. <Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. Stewart; eggs of Aphis mali on apple-twigs and probably on those of mountain ash; Aspidiotus harrisii [= Chionaspis furfurus] infests both trees.

- 580. [Walsh, B. D., and C. V. RILEY.] Gigantic rhinoceros beetle. <Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of F. G. Smith; characters of Dynastes tityrus.
- 581. [WALSH, B. D., and C. V. RILEY.] Bee queries. < Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. R. Howard; Galleria cereana always injurious; a, new swarm composed of both old and new bees.
- 582. [Walsh, B. D., and C. V. Rilley.] Insects named. < Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. W. Butterfield; characters of Clytus [= Neoclytus] capræa. (See No. 560.)
- 583. [Walsh, B. D., and C. V. Riley.] Elm-tree borer. < Amer. Ent., April, 1869, v. 1, p. 168.
 - Answer to inquiry of W. M. Gregory; Saperda lateralis and Dryobius serfasciatus attack the elm in the larva state. (See No. 696.)
- 584. [Walsh, B. D., and C. V. Riley.] The chinch-bug (*Micropus leucopterus* Say). <Amer. Ent., 1869, v. 1, May, pp. 169-177, fig. 122; June, pp. 194-199, figs. 135-139. Reprint: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 15-37, figs. 1-10. See No. 1127 for synopsis of contents.
- 585. [Walsh, B. D., and C. V. Riley.] The grape-berry moth (*Penthina vitivorana*, Packard). <Amer. Ent., May, 1869, v. 1, pp. 177-179, figs. 123-125.
 - Natural history, ravages, and means against Penthina vitivorana [== Eudemis botrana]; figures, larva, pupa, cocoon, imago, and injured grape.
- 586. [Walsh, B. D., and C. V. Riley.] Poisonous flour. <Amer. Ent., May, 1869, v. 1, p. 179.
 - Application of the term weevil; vesicatory properties of Sitophilus [= Calandra] granaria; poisonous nature of flour made from wheat infested by these insects.
- 587. [WALSH, B. D., and C. V. RILEY.] Mounding peach-trees. < Amer. Ent., May, 1869, v. 1, pp. 180-181, fig. 126.

 Mounding as a means against Sannina exitiosa; figures 3 and 2 of the same.
- 588. [Walsh, B. D., and C. V. Riley.] Going it blind. <Amer. Ent., May, 1869, v. 1, pp. 182-183.
 - Remarks on the failure of persons to observe correctly.
- 589. [Walsh, B. D., and C. V. Riley.] Another new Curculio humbug. <Amer. Ent., May, 1869, v. 1, p. 183.
 - Extract from "agricultural paper," with comments on proposed means against Conotrachelus nenuphar.
- 590. [Walsh, B. D., and C. V. Riley.] Rear-horses vs. grasshoppers. <Amer. Ent., May, 1869, v. 1, p. 184, figs. 127-128.
 - Vernacular names of Mantis [= Phasmomantis] carolina; its usefulness; figures of eggs and 3 and 2 imago.
- 591. [Walsh, B. D., and C. V. Riley.] Apple tree plant-lice. (Aphis mali, Linn.) < Amer. Ent., May, 1869, v. 1, p. 184.
 - Abundance of Aphis mali in several localities; its comparative harmlessness; means against it.

- 592. [Walsh, B. D., and C. V. Riley.] Crack-jaw names. < Amer. Ent., May, 1869; v. 1, p. 184.
 - Combination of vernacular and technical names for the accommodation of different classes of readers.
- 593. [Walsh, B. D., and C. V. Riley.] Send plenty of specimens. <a href="mailto: Amer. Ent., May, 1869, v. 1, p. 185.
 - Reasons why several specimens of insects should be sent for examination.
- 594. [Walsh, B. D., and C. V. Rilley.] White grub fungus. < Amer. Ent., May, 1869, v. 1, p. 186, fig. 129.

 Answer to inquiries of J. Smith and of T. J. Freeman; larva of Lacknosterna
- fusca infested with Cordyceps ravenelii; figure of infested larva.

 595. [Walsh, B. D., and C. V. Rilley.] "Buck fly." < Amer. Ent.,
- May, 1869, v. 1, p. 186.

 Answer to inquiry of G. W. Copley; characters of larva and image of Satur-
 - Answer to inquiry of G. W. Copley; characters of larva and image of Saturnia [= Hemileuca] maia; food-plants of the larva.
- 596. [WALSH, B. D., and C. V. RILEY.] Swarms of minute flies in rooms. <Amer. Ent., May, 1869, v. 1, p. 186.</p>
 Answer to inquiry of S. S. Rathvon; habitat of larva of Sciara sp.
- 597. [Walsh, B. D., and C. V. RILEY.] Worms in osage orange seed. <Amer. Ent., May, 1869, v. 1, p. 186.
 - Answer to inquiry of A. Plant; habits of the larva of an undetermined dipteron found among osage orange seed.
- 598. [Walsh, B. D., and C. V. Riley.] Insects to be named. < Amer. Ent., May, 1869, v. 1, p. 186.
 - Answer to inquiry of X. Q. Z.; value of names to a collection of insects.
- 599. [WALSH, B. D., and C. V. RILEY.] The spotted lady bird. < Amer. Ent., May, 1869, v. 1, p. 186, fig. 130.</p>
 Answer to inquiry of R. Seevers; value of Hippodamia [= Megilla] maculata
- as a destroyer of noxious insects; figure of the same.

 600. [Walsh, B. D., and C. V. Riley.] Eggs of the white-marked tussock moth. <Amer. Ent., May, 1869, v. 1, p. 186.
- Answer to inquiry of S. G. Knight; characters of eggs of Orgyia leucostigma.

 601. [Walsh, B. D., and C. V. Riley.] Owl's pellets. <Amer. Ent.,

 May, 1869, v. 1, p. 187.
 - Answer to inquiry of C. H. G.; presence of injurious insects in the pellets disgorged by owls [= hawks]. See No. 643.
- 602. [Walsh, B. D., and C. V. Riley.] Crab-apple borers. <Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of J. Huggins; characters of undetermined lepidopterous borer infesting crab-apple trees.
- 603. [Walsh, B. D., and C. V. Rilley.] Mosquitoes. <Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of A. M. Abbott; life-habits of $Culicid\alpha$; benefits derived from the same.
- 604. [WALSH, B. D., and C. V. RILEY.] Large silken cocoon. < Amer. Ent., May, 1869, y. 1, p. 187.

 Answer to inquiry of W. W. Butterfield; food-habits of Attacus promethea.

- 695. [WALSH, B. D., and C. V. RILEY.] Bugs in alcohol. <Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of D. P. Smith; alcohol as a means of preserving insects.
- 606. [Walsh, B. D., and C. V. Riley.] Borer in plum-twig. < Amer. Ent., May, 1869, v. 1, p. 187.
 - Answer to inquiry of W. Colwell; Elaphidion parallelum [=villosum] bred from plum-twigs; is does not prune the twig. See No. 559.
- 607. [Walsh, B. D., and C. V. Riley.] Eggs of cut-worm moth. Amer. Ent., May, 1869, v. 1, p. 188, fig. 131.
 - Answer to inquiries of G. Pauls, T. A. Thorp, and E. S. Foster; characters and figure of eggs of *Agretis inermis* [=saucia]; characters and habits of the larva.
- 608. [Walsh, B. D., and C. V. Rilby.] Snow fleas. < Amer. Ent., May, 1869, v. 1, p. 188.
 - Answer to inquiry of H. H. G. Bradt; habits and food of Podura [= Achorutes] nivicola.
- 609. [WALSH, B. D., and C. V. BILEY.] Fuzzy galls on blackberry-twigs. <Amer. Ent., May, 1869, v. 1, p. 188.</p>
 Answer to inquiry of J. Huggins; characters of Diastrophus cuscutæformis and its gall.
- 610. [WALSH, B. D., and C. V. RILEY.] Pithy galls on blackberrytwigs. <Amer. Ent., May, 1869, v. 1, p. 188. Answer to inquiry of T. W. Gordon; characters of the gall made by Diastro-

phus nebulosus.

- 611. [WALSH, B. D., and C. V. RILEY.] Moth eggs. <Amer. Ent.,
 May, 1869, v. 1, p. 188.
 - Answer to inquiries of A. M. Shute and of J. Huggins; characters f the oggs of an undetermined moth.
- 612. [WALSH, B. D., and C. V. RILEY.] Horse-hair snakes. <Amer. Ent., May, 1869, v. 1, p. 188.

 Answer to inquiry of A. M. Abbott; characters and habitats of Gordiacaa.
- 613. [Walsh, B. D., and C. V. Riley.] Imitative butterflies. <Amer. Ent., June, 1869, v. 1, pp. 189-193, figs. 132-134.
 - Immunity of Danaida from and liability of Pierida to the attacks of predatory animals; mimicry of Danaida by Pierida and of Danais archippus by Limenitis disippus; hibernating habits and description of the larva of the latter, with figures of its larva, chrysalis, imago, and hibernaculum; figure of Danais archippus; theory of the origin of mimicry.
- 614. [WALSH, B. D., and C. V. RILEY.] Cabbage-worms upon gilly-flowers. <Amer. Ent., June. 1869, v. 1, p. 199.

 Habits and food-plants, seasons, and synonyms of Plutella cruoifergrum.
- 615. [Walsh, B. D., and C. V. Riley.] "Wasps and their habits." <Amer. Ent., June, 1869, v. 1, p. 200.
 - Comments on letter of S. S. Rathvon; species having essentially different habits though externally indistinguishable should be considered specifically distinct.
- 616. [WALSH, B. D., and C. V. RILEY.] The social wasps. <Amer. Ent., June, 1869, v. 1, p. 201.

 Comments on paper by D. A. A. Nichols; habits of Vespa crabro.

- 617. [WALSH, B. D., and C. V. RILEY.] Mounding peach-trees. < Amer. Ent., June, 1869, v. 1, pp. 201-202.
 - Comments on letter of R. L. Wells; success of the mounding system against **Egeria [=Sannina] exitiosa.
- 618. [WALSH, B. D., and C. V. RILEY.] Out of evil there cometh good. <Amer. Ent., June, 1869, v. 1, p. 202.
 - Probable abundance of the fruit crop in southern Illinois and in Missouri in 1869 due to the pruning of the trees by Tibicen septendecim in 1868.
- 619. [Walsh, B. D., and C. V. Riley.] The periodical Cicada. <Amer. Ent., June, 1869, v. 1, p. 202.
 - Extract from No. 474; request for information of the appearance of Cicada [= Tibicen] septendecim in any part of the United States in 1869.
- 620. [Walsh, B. D., and C. V. Riley.] The Curculio scarcer than last year. <Amer. Ent., June, 1869, v. 1, p. 202.
 - Extract from letter of A. M. Brown; comparative scarcity of Constructedus nenuphar in 1869.
- 621. [Walsh, B. D., and C. V. Riley.] The American Entomological Society. <Amer. Ent., June, 1869, v. 1, p. 203.
 - Notice of the formation, publications, aims, and needs of the American Entomological Society; proposition for the raising of a fund for the support of the society.
- 622. [Walsh, B. D., and C. V. Riley.] Remarkable peculiarity in the insect world. <Amer. Ent., June, 1869, v. 1, p. 204.
 - Insects contrasted with animals of other groups; as a rule they produce but one brood of offspring in the course of their lives.
- 623. [WALSH, B. D., and C. V. RILEY.] On our table. <Amer. Ent., June, 1869, v. 1, pp. 204-205.
 - Notices of: The Harris correspondence; The butterflies of North America, by W. H. Edwards; Guide to the study of insects, by A. S. Packard, jr.
- 624. [Walsh, B. D., and C. V. Riley.] Cannibal mites. < Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of C. S. Davis; Trombidium sp. found preying on grass-hopper eggs.
- 625. [Walsh, B. D., and C. V. Riley.] Gnats. < Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of W. O. Hiskey; distinctive characters of Culex and Chironomus; appearance of "clouds" of Chironomus.
- 626. [Walsh, B. D., and C. V. Riley.] Cut-worms severing cabbage plants. <Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of N. C. Burch; characters of the larva of Agrotis telifera [= ypsilon].
- 627. [Walsh, B. D., and C. V. Riley.] Beetle named. <Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of J. M. Shaffer; Anisodactylus baltimorensis flying in great numbers at Fairfield, Iowa.
- 628. [Walsh, B. D., and C. V. Riley.] Tiger beetles. < Amer. Ent., June, 1869, v. 1, p. 205.
 - Answer to inquiry of J. M. Shaffer; habitat of Cicindela vulgaris.

- 629. [WALSH, B. D., and C. V. RILEY.] Cocoons and chrysalids named. <Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of A. S. Fuller; characters of larva and image of Ceratocampa [= Citheronia] regalis; food-plants of its larva; characters of the cocoons of Attaous promethea, A. cecropia, A. [= Telea] polyphemus. and of Thyridopteryx ephemeræformis.
- 630. [WALSH, B. D., and C. V. RILEY.] White lined morning Sphinx. <Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of S. Blanchard; characters, transformations, habits, and distribution of *Deilephila lineata*; food-plants of its larva.
- 631. [Walsh, B. D., and C. V. Riley.] Insects named. <Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of W. W. Butterfield; identification of insects sent; differences between the spring and autumn broods of *Drasteria erechtea*.
- 632. [WALSH, B. D., and C. V. RILEY.] Peach-twig borer. <Amer. Ent., June, 1869, v. 1, p. 206, fig. 140.</p>
 Answer to inquiry of W. Muir; characters, habits, and figure of the larva of
 - Gortyna hitela.
- 633. [Walsh, B. D., and C. V. Riley.] The apple-twig borer. <Amer. Ent., June, 1869, v. 1, p. 206, fig. 141.
 - Answer to inquiry of A. Hinckley; habits, sexual differences, and figure of Bostrichus [= Amphicerus] bicaudatus.
- 634. [Walsh, B. D., and C. V. Riley.] Apple-tree borers on south side of trees. <Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of J. F. Wielandy; preference shown for the south and southwest sides of trees by the larva of Chrysobothris femorata.
- 635. [Walsh, B. D., and C. V. Riley.] Cocoons of the Cecropia moth. Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of T. W. Gordon; comparison between the cocoons of Telea polyphemus and Attacus cecropia.
- 636. [Walsh, B. D., and C. V. Biley.] Flea-beetles. <Amer. Ent., June, 1869, v. 1, p. 206.
 - Answer to inquiry of F. Hecker; characters of an undescribed Longitarsus infesting wheat fields.
- 637. [Walsh, B. D., and C. V. Riley.] Strawberry bugs. < Amer. Ent., June, 1869, v. 1, p. 207.
 - Answer to inquiry of J. M. Pearson; characters, ravages, and means against Corimelana pulicaria.
- 638. [Walsh, B. D., and C. V. Riley.] Eggs on apple-trees. < Amer. Ent., June, 1869, v. 1, p. 207, fig. 142.
 - Answer to inquiry of H. Compton; characters of eggs of Sinea diadema; figure of the imago; its value as a destroyer of canker-worms.
- 639. [Walsh, B. D., and C. V. Riley.] Lady-bird larvæ. <Amer. Ent., June, 1869, v. 1, p. 207, fig. 143.
 - Answer to inquiry of E. S. Foster; characters of the larva of Hippodamia convergens; figures its larva, pupa, and imago; usefulness of Cocinellidæ.

- 640. [Walsh, B. D., and C. V. Riley.] White-grub fungus, again. <Amer. Ent., June, 1869, v. 1, p. 207, fig. 144.
 - Answer to inquiry of W. C. Holmes; figures larva of Lachnosterna fusca attacked by Cordyceps ravenelii.
- 641. [WALSH, B. D., and C. V. RILEY.] Bag-worms. < Amer. Ent., June, 1869, v. 1, p. 207.
 - Answer to inquiry of C. Parry; means against Thyridopteryx sphemerasformis.
- 642. [Walsh, B. D., and C. V. Riley.] Tent-caterpillar. <Amer. Ent., June, 1869, v. 1, p. 208, fig. 145.
 - Answer to inquiry of S. Blanchard; characters of the image and figures of the early stages of Clisiocampa americana; means against the same.
- 643. [Walsh, B. D., and C. V. Riley.] Hawk's pellets. < Amer. Ent., June, 1869, v. 1, p. 208.
 - Answer to inquiry of C. H. G.; insectivorous habits of Buteo pennsylvanious. See No. 601.
- 614. [WALSH, B.D., and C.V.RILEY.] Plant lice on berberry. < Amer. Ent., June, 1869, v. 1, p. 208.

 Answer to inquiry of J. R. Preston; means against Aphidida.
- 645. [Walsh, B. D., and C. V. Riley.] Tent-caterpillar of the forest. <Amer. Ent., June, 1869, v. 1, p. 208, fig. 146.
 - Answer to inquiry of G. Whitcomb; characters, habits, parasites, means against, and figure of larva of Clisiocampa sylvatica [= disstria].
- 646. [WALSH, B. D., and C. V. RILEY.] Cotton insects. <Amer. Ent., July, 1869, v. 1, pp. 209-214, fig. 147-151.</p>
 Descriptions and figures of all stages of Aletia xylina and of Heliothis armigera; habits, seasons, ravages, food-plants of, and means against the same; figures larva of Clisiocampa sylvatica [= disstria].
- 647. [WALSH, B. D., and C. V. RILEY.] The true Army-worm (Leucania unipuncta Haworth). <Amer. Ent., July, 1869, v. 1, pp. 214-217, fig. 152-155.
 - Descriptions and figures of larva, pupa, and image of Leucania unipuncta; seasons, ravages, and enemies of the same; figure of Exerista militaris [= Nemoræa leucaniæ].
- 648. [Walsh, B. D., and C. V. Riley.] Belated individuals of the periodical Cicada. <Amer. Ent., July, 1869, v. 1, p. 217.

 Occurrence of scattering individuals of Tibicon septendecim in years before or after their regular period.
- 649. [WALSH, B. D., and C. V. RILEY.] Is the Curculio scarcer than it was last year? <Amer. Ent., July, 1869, v. 1, pp. 217, 218. Record of observations to prove the comparative scarcity of Constructed nonuphar during 1869.
- 650. [Walsh, B. D., and C. V. Riley.] To destroy Colorado potato bugs. <a href="https://doi.org/10.1007/j.jps.1019.0007/j.jps.
- Reprint of communication of G. Liddle, with comments; Paris green as a means against *Doryphora* 10 lineata.

 651. [Walsh, B. D., and C. V. Riley.] Dr. Hull's Curculio catcher.
- 651. [WALSH, B. D., and C. V. KILEY.] Dr. Hull's Curculo-catcher.
 <Amer. Ent., June, 1869, v. 1, pp. 220–221, fig. 156.</p>
 Description and figure of Hull's Curculio-catcher.

- 652. [WALSH, B. D., and C. V. RILEY.] The New York weevil (Ithycerus noveboracensis, Forster). <Amer. Ent., July, 1869, v. 1, pp. 221-222, fig. 157.
 - Habits, food-plants, distribution, description, and figure of larva and image of Ithycerus noveboracensis; means against the same.
- 653. [WALSH, B. D., and C. V. RILEY.] Mounding peach-trees again. <Amer. Ent., July, 1869, v. 1, p. 223.

 Letter of A. Dean, with comments; characters and habitat of Mycetophila persion.
- 654. [WALSH, B. D., and C. V. RILEY.] Overcrowded. <Amer. Ent., July, 1869, v. 1, p. 223.

 Duties of a State entomologist; impossibility of answering inquiries received during the past month.
- 655. [WALSH, B. D., and C. V. RILEY.] No air holes needed in sending insects. < Amer. Ent., July, 1869, v. 1, p. 223.</p>
 Directions for sending living insects.
- 656. [WALSH, B. D., and C. V. RILEY.] Plum-leaf worms. < Amer. Ent., July, 1869, v. 1, p. 223.
 Answer to inquiry of W. D. Hiskey; characters of larva of Lyda sp.
- 657. [WALSH, B. D., and C. V. RILEY.] Seed-corn maggot. <Amer. Ent., July, 1869, v. 1, p. 224, figs. 138–139.</p>
 Answer to inquiry of G. Pauls; characters, ravages, means against, and figure of larva of Anthomyia zeas; characters of the imago; figure of the puparium.
- 658. [WALSH, B. D., and C. V. RILEY.] Cut-worms. <Amer. Ent., July, 1869, v. 1, p. 224.

 Answer to inquiry of N. C. Birch; ravages of Agrotis telifera [= ypsilon].
- 659. [WALSH, B. D., and C. V. RILEY.] New York weevil. <Amer. Ent., July, 1869, v. 1, p. 224.</p>
 Answer to inquiries of D. H. Kauffman and of W. D. Turrill; ravages of Ithycerus noveboracensis.
- 660. [WALSH, B. D., and C. V. RILEY.] Insects around peach-trees. Answer to inquiry of G. C. Brodhead; larvæ of Asilus sp. and of Myostophila persicæ found around roots of peach-trees.
- 661. [WALSH, B. D., and C. V. RILEY.] Large green worm in a peach.

 Amer. Ent., July, 1869, v. 1, p. 224.
 Answer to inquiry of G. Wilgus; characters and food-habits of an undetermined larva found in a peach.
- 662. [WALSH, B. D., and C. V. RILEY.] Ichneumon flies. < Amer. Ent., July, 1869, v. 1, p. 224.</p>
 Answer to inquiry of H. Klinehaus; characters and figure of cocoons of Microgaster sp.
- 663. [WALSH, B. D., and C. V. RILEY.] Raspberry worms. < Amer. Ent., July, 1869, v. 1, p. 224.

 Answer to inquiry of B. Borden; characters and ravages of Sciandria [= Mo-

nophadnus] rubi.

- 664. [Walsh, B. D., and C. V. Rilley.] Hairy grape-leaf folders. <Amer. Ent., July, 1869, v. 1, p. 224.
 - Answer to inquiry of A. C. Davis; characters of larva and image of Pterophorus [= Oxyptilus] periscelidactylus.
- 665. [Walsh, B. D., and C. V. RILEY.] Row of eggs in maple-twigs. <Amer. Ent., July, 1869, v. 1, p. 224.

Answer to inquiry of J. Bower; characters of eggs of undetermined katydid.

- 666. [Walsh, B. D., and C. V. Riley.] Butterfly named. < Amer. Ent., July, 1869, v. 1, p. 224.
 - Answer to inquiry of A. R. Bodley; characters of Papilio marcellus; food-plants of its larva.
- 667. [Walsh, B. D., and C. V. Riley.] Grasshopper's eggs. < Amer. Ent., July, 1869, v. 1, p. 224.
 - Answer to inquiry of E. P. Burlingame; characters of the eggs of an undetermined grasshopper and of the image of *Edipoda* [= Discosteria] carolina.
- 668. [Walsh, B. D., and C. V. Riley.] Asilus fly larvæ. <Amer. Ent., July, 1869, v. 1, p. 225, figs. 161-162.
 - Answer to inquiry of G. Pauls; characters, habits, and figure of larva of Asilus sp.; food-habits of larva and image of A. sericeus; figure of the image of the same; Trupanea [= Promachus] apivorus as a destroyer of bees.
- 669. [Walsh, B. D., and C. V. RILEY.] New insect-foe of the black-berry. <Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiry of C. Parry; comparative characters of Aphidida and Psyllida; habits of Psylla rubi [= Trioza tripunctata].
- 670. [WALSH, B. D., and C. V. RILEY.| Army-worm. <Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiry of J. H. Butts; ravages and food-plauts of Leucania unipuncta.
- 671. [Walsh, B. D., and C. V. Riley.] Green grape-vine worm. Amer. Ent., July, 1869, v. 1, p. 225, fig. 163.
 - Answer to inquiry of G. Pauls; characters, food-plants, and figure of larva of Pyrophila pyramidoides.
- 672. [Walsh, B. D., and C. V. Riley.] Rose slug. <Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiries of G. W. Copley and B. S. Morris; characters, ravages, and means against Selandria [= Monostegia] rosæ.
- 673. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiry of J. Weed; food-habits of Leptostylus aculiferus and of Podabrus modestus.
- 674. [WALSH, B. D., and C. V. RILEY.] Chrysalis of the virgin tiger moth. <Amer. Ent., July, 1869, v. 1, p. 225.
 - Answer to inquiry of C. Mallinckrodt; characters of Arctia virgo; supposed food-plants of its larva.
- 675. [Walsh, B. D., and C. V. Rilley.] Clover-worms. < Amer. Ent., July, 1869, v. 1, p. 226, fig. 164.
 - Answer to inquiry of E. P. Flanders; characters of Asopia costalis; habits of its larva; figures larva, pupa, cocoon, and imago.

- 676. [WALSH, B. D., and C. V. RILEY.] Raspberry brand. < Amer. Ent., July, 1869, v. 1, p. 226.
 - Answer to inquiry of J. M. Beecher; larva of small gnat feeding on raspberry-brand.
- 677. [WALSH, B. D., and C. V. RILEY.] Rose-worms. < Amer. Ent., July, 1869, v. 1, p. 226.
 - Answer to inquiry of G. S. Grover; character, ravages, food-plants, and means against Heliothis margidens [= Pyrrhia exprimens].
- 678. [Walsh, B. D., and C. V. Biley.] Plum-tree plant-lice. <Amer. Ent., July, 1869, v. 1, p. 226, figs. 165-167.
 - Answer to inquiry of T. W. Gordon; characters of Aphis prunifolia and A. [=Myzus] cerasi; ravages, enemies, and means against Aphidida; figures larva of coccinellid, syrphid and hemerobi?.
- 679. [Walsh, B. D., and C. V. Riley.] 1chneumon-flies. <Amer. Ent., July, 1869, v. 1, p. 226.
 - Answer to inquiries of S. J. Throp and J. E. Trabue; life-habits of Micro-gaster sp. ?
- 680. [Walsh, B. D., and C. V. Rilley.] A new Curculio humbug. <Amer. Ent., July, 1869, v. 1, p. 226.
 - Answer to inquiry of E. P. Flanders; uselessness of a patent lamp for destroying Constructed us nenuphar.
- 681. [Walsh, B. D., and C. V. Riley.] Sweet-potato beetles. < Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiries of Subscriber and A. E. Trabue; food habits and characters of larva and image of Coptocycla aurichalcea and of C. [= Cassida] bivittata.
- 682. [Walsh, B. D., and C. V. Riley.] Injured strawberry and grapevines. <Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of W. P. Pierson; ravages of Capsus oblineatus [=Lyguspratensis].
- 683. [Walsh, B. D., and C. V. Riley.] Plum-tree insects. < Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of J. F. Waters; characters of Bibio albipennis; food-habits of its larva.
- 684. [Walsh, B. D., and C. V. Rilley.] Apple-tree bugs. <Amer. Ent., July, 1869, v. 1, p. 227.

 Answer to inquiry of W. L. Youse; character of Brochymena annulata.
- 685. [Walsh, B. D., and C. V. Riley.] Hickory-stem gall-louse. <Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of B. F. Long; characters of galls of Phylloxera caryacaulis; food-habits of Thrips.
- 686. [Walsh, B. D., and C. V. Riley.] Grape-vine leaf-hopper. < Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of R. M. Copeland; means against Tettigonia [= Typhlocyba] vitis.
- 687. [WALSH, B. D., and C. V. RILEY.] Twelve-spotted Diabrotica. <Amer. Ent., July, 1869, v. 1, p. 227, fig. 168.
 - Answer to inquiries of E. S. Foster and R. D. Parker; characters and means against *Diabrotica* 12-punctata; figure of the same.

- 688. [Walsh, B. D., and C. V. Riley.] The caterpillar of the forest. <Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of M. McKenzie; food-habits and means against Clisio-campa sylvatica [=disstria].
- 689. [Walsh, B. D., and C. V. Rilby.] Strawberry destroyer. < Amer. Ent., July, 1869, v. 1, p. 227.

Answer to inquiry of G. W. Copley; habits of unknown strawberry destroyer.

- 690. [WALSH, B. D., and C. V. RILEY.] Eggs of bugs on strawberry. <Amer. Ent., July, 1869, v. 1, p. 227.
 - Answer to inquiry of A. S. Fuller; characters of eggs of unknown reduvid attacked by parasites.
- 691. [Walsh, B. D., and C. V. Riley.] Frog-spittle insects. < Amer. Ent., July, 1869, v. 1, p. 228. Reprint: < Cultivator and Country Gentleman, 29 July, 1869, v. 34, p. 82.
 - Answer to inquiry of J. B. Hartwell:; habits, characters, and injuries of Aphrophora quadrangularis.
- 692. [Walsh, B. D., and C. V. Riley.] Eggs of ground-beetle. <Amer. Ent., July, 1869, v. 1, p. 228.

Answer to inquiry of E. J. Ayres; characters of eggs of undetermined groundbeetle found under bark of pear-twigs.

- 693. [Walsh, B. D., and C. V. Riley.] New insect-foe of the potato. <Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of I. Hicks; food-habits of Cassida [= Coptocycla] clavata.
- 694. [Walsh, B. D., and C. V. RILEY.] Tomato-stalk borer. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of E. J. Ayres; Gortyna nitela injurious to tomato-stalks.
- 695. [Walsh, B. D., and C. V. Riley.] Breeding cages. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of H. S. Redney; description of cage for breeding insects.
- 696. [Walsh, B. D., and C. V. Rilley.] Elm-tree borer. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of W. M. Gregory; characters and food-habits of Physocnemum brevilineum. See No. 583.
- 697. [WALSH, B. D., and C. V. RILEY.] Peach-twig borer. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of G. Fisher; means against undetermined lepidopterous borer in twigs of peach.
- 698. [Walsh, B. D., and C. V. Rilley.] Eggs of periodical Cicada in savin-twig. Amer. Ent., July 1809, v. 1, p. 228.
 - Answer to inquiry of J. A. Greason; Tibicen septendecim ovipositing in twigs of Juniperus sabina.
- 699. [Walsh, B. D., and C. V. Riley.] Elm-tree saw-fly. < Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of A. R. Whitney; characters of larva and image of Cimbex laportei [= americana]; food-plants of its larva.
- 700. [Walsh, B. D., and C. V. RILEY.] Snout-beetle. <Amer. Ent., July, 1869, v. 1, p. 228.
 - Answer to inquiry of H. Kleinhaus; supposed food-habits of Hylobius confusus.

- 701. [Walsh, B. D., and C. V. Riley.] The close of the first volume. <a href=
 - Prospectus of the second volume of the American Entomologist.
- 702. [WALSH, B. D., and C. V. RILEY.] The Royal horned caterpillar. (Ceratocampa (Citheronia) regalis, Fabr.). <Amer. Ent., August, 1869, v. 1, pp. 230-231, pl. 1.
 - Habits, seasons, sexual characters, food-plants, and vernacular names of *Citheronia regalis*; descriptions and figures of larva and pupa; figure of larva, pupa, and imago.
- 703. [WALSH, B. D., and C. V. RILEY.] Comparative scarcity of the Curculio again. < Amer. Ent., August, 1869, v. 1, p. 241.</p>
 Observations on the comparative abundance of Constructedus nenuphar during the early summer of 1869.
- 704. [WALSH, B. D., and C. V. RILEY.] *A possible cause of the bee disease. <Amer. Ent., August, 1869, v. 1, pp. 241-242.</p>
 Reprint of article by P. H. Philbrook (Amer. Bee Journal, May, 1869), with comments; dipterous enemies of the honey-bee.
- 705. [WALSH, B. D., and C. V. RILEY.] Ash and mountain ash. < Amer. Ent., August, 1869, v. 1, pp. 243–244.</p>
 Criticisms of paper of H. Shimer (Trans. Ill. State Hortic. Soc., 1868).
- 706. [WALSH, B. D., and C. V. RILEY.] How the Curculio flies by night. <Amer. Ent., August, 1869, v. 1, p. 244.

 Extract from lecture by I. P. Trimble, with comment.
- 707. [Walsh, B. D., and C. V. Riley.] The periodical Cicada; our first brood established. <Amer. Ent., August, 1869, v. 1, p. 244.

 Appearance in Connecticut of a brood of Tibicen septendooim in 1869.
- 708. [WALSH, B. D., and C. V. RILEY.] Be on the guard. Amer. August, 1869, v. 1, p. 244.
 Need of care in the transportation of living insects; accidental introduction of Doryphora 10-lineals.
- 709. [Walsh, B. D., and C. V. Riley.] Prophecy fulfilled. <Amer. Ent., August, 1869, v. 1, p. 244.

 Spread of *Doryphora* 10-lineats through Michigan.
- 710. [Walsh, B. D., and C. V. Riley.] A poisonous worm. Amer.
 Ent., August, 1869, v. 1, p. 245.
 Extract from exchange; larva of *Protoparce celeus* not poisonous.
- 711. [WALSH, B. D., and C. V. RILEY.] Wheat midge, alias milk weevil, alias red weevil. <Amer. Ent., August, 1869, v. 1, p. 245.
 - Answer to inquiry of C. Corbit; characters of larva of Cocidomyia [= Diplosis] tritio; change in heads of wheat.
- 712. [Walsh, B. D., and C. V. Rilley.] Large fish-fly. < Amer. Ent., August, 1869, v. 1, p. 245.
 - Answer to inquiry of A. R. McCutchen; characters of Chauliodes pectinicornis; habits of the larva of C. rastricornis.

6 ENT

- 713. [Walsh, B. D., and C. V. Riley.] Cottonwood leaf-galls. < Amer. Ent., August, 1869, v. 1, p. 245.
 - Answer to inquiry of J. B. Taylor; habits of Pemphigus populicaulis; characters of its gall.
- 714. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 245.
 - Answer to inquiry of J. G. Goodrich; characters of Saperda bivittata [= candida].
- 715. [Walsh, B. D., and C. V. Riley.] Four-lined leaf-bug on currant. <Amer. Ent., August, 1869, v. 1, p. 246.
 - Answer to inquiry of M. B. Bateman; characters, habits, and means against Capsus 4-vittatus [= Pæcilocapsus lineatus].
- 716. [Walsh, B. D., and G. V. Rilley.] Bee moth. < Amer. Ent., August, 1869, v. 1, p. 246, fig. 182.
 - Answer to inquiry of S. Blanchard; ravages and means against Galleria ocreana; figures larva, pupa, cocoon, and image of the same.
- 717. [WALSH, B. D., and C. V. RILEY.] Canker-worm parasites. <Amer. Ent., August, 1869, v. 1, p. 246.
 - Answer to inquiry of J. Petit; mention of Microgaster sp., parasitic on Anisopteryx.
- 718. [WALSH, B. D., and C. V. RILEY.] Beetles swarming about the lawn. <Amer. Ent., August, 1869, v. 1, p. 246.
 - Answer to inquiry of 8. Thompson; characters of Gymnetis [=Allorhina] nitida; habits of its larva.
- 719. [Walsh, B. D., and C. V. Riley.] Tiger-beetle larva. < Amer. Ent., August, 1869, v. 1, p. 246.
 - Answer to inquiry of R. J. Dodge; characters and habits of larva of Cicindel-idæ.
- 720. [Walsh, B. D., and C. V. Riley.] Large compound gall on grapevine. <Amer. Ent., August, 1869, v. 1, p. 247, fig. 183.
 - Answer to inquiries of A. S. Fuller and D. W. Kauffman; characters and figure of gall of *Lasioptera vitis*; habits of its larva; enemy and parasite of the same.
- 721. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 247.
 - Answer to inquiry of M. Treat; habits of larva of Temnochila [= Trogosita] virescens, of Aplodes [= Synchlora] rubivora, and of Calosoma calidum.
- 722. [Walsh, B. D., and C. V. Riley.] Destructive larvæ. <Amer. Ent., August, 1869, v. 1, p. 247.
 - Answer to inquiry of B. F. Lee; ravages of an undetermined larva; parasites of Saturnia [= Hemileuca] maia.
- 723. [Walsh, B. D., and C. V. Riley.] Eggs of tree-cricket on grape-vine. <Amer. Ent., August. 1869, v. 1, p. 247.
 - Answer to inquiry of B. F. Lee; characters of eggs of *Œcanthus niveus*; carnivorous habits of the larva of the same.

- 724. [Walsh, B. D., and C. V. Riley.] Grape-vine leaf-gall. < Amer. Ent, August, 1869, v. 1, p. 248, fig. 184.
 - Answer to inquiry of A. A. Hilliard; characters and figure of the leaf-galls of *Phylloxera vitifolia* [= vastatrix]; its enemies and means against them; varieties of grape infested; identity of the leaf- and root-galls; criticism of the new genera and families of H. Shimer.
- 725. [Walsh, B. D., and C. V. RILEY.] Grape-vine insects. < Amer. Ent., August, 1869, v. 1, p. 248.
 - Answer to inquiry of G. Pauls; characters of Pelidnota punctata; its larval habits; characters, habits, and food-plants of Enchophyllum [= Enchenopa] binotata.
- 726. [WALSH, B. D., and C. V. RILEY.] Museum pests. < Amer. Ent., August, 1869, v. 1, p. 248.

 Answer to inquiry of C. P. Faulkner; ravages of Dermestes lardarius.
- 727. [WALSH, B. D., and C. V. RILEY.] Wheat maggets. < Amer. Ent., August, 1869, v. 1, p. 248.</p>
 Answer to inquiry of S. K. Faulkner; characters of larva of Meromyza americana injurious to heads of wheat.
- 728. [WALSH, B. D., and C. V. RILEY.] Parasites on "hateful grass-hopper." < Amer. Ent., August, 1869, v. 1, p. 249.

 Answer to inquiry of S. K. Faulkner; characters and habits of Astoma [= Trombidium] locustarum.
- 729. [WALSH, B. D., and C. V. RILEY.] Crippled moths. < Amer. Ent., August, 1869, v. 1, p. 248.</p>
 Answer to inquiry of C. P. Faulkner; conditions needed to enable moths to expand their wings; means by which insects walk on smooth surfaces.
- 730. [WALSH, B. D., and C. V. RILEY.] Insects on the oleander. Amer. Ent., August, 1869, v. 1, p. 249, fig. 185.
 Answer to inquiry of T. W. Gordon; means against undetermined Coccid on cleander; habits and figure of Chilocorus bivulnerus.
- 731. [WALSH, B. D., and C. V. RILEY.] Insects found on apple trees. <a h
- 732. [WALSH, B. D., and C. V. BILEY.] Beetles named. <Amer. Ent., August, 1869, v. 1, p. 249.

 Answer to inquiry of J. M. Shaffer; Lytta atrata [= Epicauta pennsylvanica] caught on rag-weed.
- 733. [Walsh, B. D., and C. V. Riley.] Gigantic water-bug. < Amer. Ent., August, 1869, v. 1, p. 249, fig. 186.

 Answer to inquiry of E. M. Downing; habits and figure of Belostoma grandis [= americanum].
- 734. [WALSH, B. D., and C. V. RILEY.] Worm eating into green tomatoes. <Amer. Ent., August, 1869, v. 1, p. 249. Answer to inquiry of D. L. Hall; food-habits of Gortyna nitela.

- 735. [Walsh, B. D., and C. V. Riley.] Miscellaneous. < Amer. Ent., August, 1869, v. 1, p. 249.
 - Answer to inquiry of G. W. Copley; food-habits of Chrysochus auratus, Hemileuca maia, and Aphis [= Myzus] ribis; habits of Lozotania [= Cacacia] rosaceana.
- 736. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of H. A. Munger; food-habits of Cassida pallida [= Coptocycla aurichalcea], Lytta murina [= Macrobasis unicolor], and of the larvæ of the species of Prionus.
- 737. [Walsh, B. D., and C. V. Rilley.] Lightning-hoppers. < Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of T. W. Gordon; characters and habits of Pæciloptera pruinosa; vernacular names of Fulgoridæ, Membracidæ, and Jassidæ.
- 738. [WALSH, B. D., and C. V. RILEY.] Bag-worms. < Amer. Ent., August, 1869, v. 1, p. 250.</p>
 Answer to inquiries of T. W. Gordon and S. Thompson; food-habits of Thy-
 - Answer to inquiries of T. W. Gordon and S. Thompson; food-habits of Thyridopteryx ephemeræformis.
- 739. [Walsh, B. D., and C. V. Riley.] Woolly gall on white oak. <Amer. Ent., August, 1869, v. 1, p. 250, fig. 187.
 - Answer to inquiry of A. S. Fuller; characters and figure of the gall of Cynips [=Andricus] seminator.
- 740. [Walsh, B. D., and C. V. Riley.] Cabbage pests. < Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of J. A. Williams; characters and habits of an undetermined elaterid larva; means against the larvæ of *Elateridæ*.
- 741. [Walsh, B. D., and C. V. Riley.] Bugs gathering on pear shoots. <a href="
 - Answer to inquiry of E. J. Ayres; characters and habits of Corimelena pulicaria.
- 742. [Walsh, B. D., and C. V. Riley.] Potato-bug. <Amer. Ent., August, 1869, v. 1, p. 250.
 - Answer to inquiry of J. B. Cartwell; food-habits of Cassida [= Coptocycla] clavata.
- 743. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of H. T. Birch; Trochilium [= Egeria] acerni bred from maple.
- 744. [Walsh, B. D., and C. V. Riley.] Apple-tree worms. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of C. Waters; abundance of undetermined moth (Corycia vestaliata?); larva of the same on apple-trees.
- 745. [Walsh, B. D., and C. V. Riley.] Oak-fig gall. <Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of A. Fendler; characters of the gall of Cynips [= Bio-rhiza] forticornis and of a guest-fly, Ceroptres ficus, inhabiting the same.

- 746. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of D. L. Phares; irregular appearance of Cicada tredecim [= Tibicen septendecim]; food-habits of Oncideres cingulata.
- 747. [WALSH, B.D., and C. V. RILEY.] Small apple-leaf worms. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of H. Compton; means against Spilonota oculana [== Tmetocera ocellana].
- 748. [Walsh, B. D., and C. V. Riley.] Rose bug on apples. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of A. Dean; food-plants and means against Macrodactylus subspinosus.
- 749. [WALSH, B. D., and C. V. RILEY.] Unicorn apple-tree caterpillar. <Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of G. C. Brodhead; characters of the larva and imago of Notodonta [= Cœlodasys] unicornis; food-plants of the larva of the same.
- 750. [WALSH, B. D., and C. V. RILEY.] Large water beetle. < Amer. Ent., August, 1869, v. 1, p. 251.
 - Answer to inquiry of S. E. Mumford; characters of Cybister fimbriolatus. See No. 816.
- 751. [WALSH, B. D., and C. V. RILEY.] Beetles around peach-trees. <Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of E. Hollister, jr.; characters and habits of Helope pullus [= areus].
- 752. [WALSH, B. D., and C. V. RILEY.] A quick traveler. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of G. C. Brodhead; characters and harmlessness of Cormatia forceps.
- 753. [Walsh, B. D., and C. V. RILEY.] Unknown moth. < Amer. Ent., August, 1869, v. 1, p. 252.
- Answer to inquiry of W.G. Barton; distribution of Junonia lavinia in Illinois. 754. [Walsh, B. D., and C. V. Riley.] Corn-borer. < Amer. Ent.,
- August, 1869, v. 1, p. 252. Answer to inquiry of F. M. N.; identification of the larva of Gortyna nitela
- injurious to corn. 755. [WALSH, B. D., and C. V. RILEY.] Horns of stag-beetle. < Amer.
 - Ent., August, 1869, v. 1, p. 252. Answer to inquiry of C. R. Edwards; larval habits of Lucanus elaphus. .
- 756. [WALSH, B. D., and C. V. RILEY.] Imported gooseberry worms. <Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of C. P. Faulkner; means against Nematus ribesii.
- 757. [WALSH, B. D., and C. V. RILEY.] Worm on bark of walnut-tree. Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of F. S. Fuller; undetermined notodontoid larva on the bark of walnut.

- 758. [WALSH, B. D., and C. V. RILEY.] Rotten root. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of J. M. Beecher; scavenger habits of undetermined larva feeding on dead roots of apple-trees.
- 759. [Walsh, B. D., and C. V. Rilley.] Large dragon-fly. Amer. Ent., August, 1869, v. 1, p. 252.

 Answer to inquiry of G. S. Grover: characters and neefulness of Recket con-
 - Answer to inquiry of G. S. Grover; characters and usefulness of *Eschna constricta*.
- 760. [Walsh, B. D., and C. V. Riley.] Stinging larvæ. < Amer. Ent., August, 1869, v. 1, p. 252.
 - Answer to inquiry of E. H. King; characters and urticating properties of Saturnia [= Hemileuca] maia; peach-blow potatoes avoided by Doryphora 10-lineata.
- 761. [WALSH, B. D., and C. V. RILEY.] Raspberry worms. < Amer. Ent., August, 1869, v. 1, p. 252.</p>
 Answer to inquiry of E. H. Beebe; characters of larva of Sclandria [= Monophadnus] rubi.
- 762. [WALSH, B. D., and C. V. RILEY.] Leaf-galls and caterpillars on the sugarberry. <Amer. Ent., August, 1869, v. 1, p. 252.</p>
 Answer to inquiry of S. L. Scofield; undetermined gall on leaves of Celtis occidentalis; Orgyia leucostigma feeding on the leaves of the same.
- 763. [Walsh, B. D., and C. V. Rilley.] Dark grape-worm. < Amer. Ent., August, 1869, v. 1, p. 252.
- Answer to inquiry of T. W.G.; characters and food-plants of *Thyreus abbotii*.

 764. [Walsh, B. D., and C. V. Riley.] Too fond of honey. < Amer.
- Ent., August, 1869, v. 1, p. 252.

 Answer to inquiry of T. W. Gordon; Cermatia forceps found in a jar of honey.
- 765. [WALSH, B. D., and C. V. RILEY.] Caterpillar of polyphemus moth. <Amer. Ent., August, 1869, v. 1, p. 252.

 Answer to inquiry of A. De Wyl; larva of Telea polyphemus feeding on plums.
- 766. [Walsh, B. D., and C. V. Riley.] Why noxious insects increase upon us. < Amer. Ent., September-October, 1869, v. 2, pp. 1-2. Reasons why noxious insects increase; review of articles by E. S. Hull, H. W. Beecher, and Puritan.
- 767. [WALSH, B. D., and C. V. RILEY.] Tortoise-beetles. <Amer. Ent., September-October, 1869, v. 2, pp. 2-5, figs. 1-3. Resemblances of insects to other animals or to their surroundings; distribution, food-plants, and figure of Deloyala [= Coptocycla] clavata; description and figure of Physonota quinque-punctata n. sp. [= unipunctata]; figure of</p>
 - and figure of *Physonota quinque-punctata* n. sp. [= unipunctata]; figure of the larva of the same; correspondence of structural differences in larva with those in images of *Cassidida*, and with differences in food-plants; list of insects injurious to *Solanum*; figure of pupa and image of *Chelymorpha cribraria* [= argus].
- 768. [WALSH, B. D., and C. V. RILEY.] Scientific nomenclature. Amer. Ent., September—October, 1869, v. 2, pp. 5–8. Rules observed in giving specific names to animals and plants; the law of priority; nature and extent of generic subdivisions.

- 769. [WALSH, B. D., and C. V. RILEY.] Killing apple-worms by machinery. <Amer. Ent., September-October, 1869, v. 2, p. 9.
 - Extract from article of L. P. Haskell, with comment; use of rags in place of hay-bands as traps for *Carpocapsa pomonella*; destruction of the insects caught by means of a clothes-wringer.
- 770. [Walsh, B. D., and C. V. Riley.] A potter wasp (Odynerus flavipes f Fabr.). < Amer. Ent., September-October, 1869, v. 2, p. 10, fig. 4.
 - Method employed by *Bumenidæ* to provision their nests; construction of the same by species of *Odynerus*; habits of *O. flavipes* in provisioning a nest with several species of larvæ; figure of the image of the same and of the nest of a species of *Odynerus*; structure of wings and habits of *Vespidæ*, *Eumenidæ*, and fossorial wasps.
- 771. [WALSH, B. D., and C. V. RILEY.] Tomato-worms not poisonous. <Amer. Ent., September-October, 1869, v. 2, p. 11.
 - Prejudices in regard to certain animals; presence of the horn on the larvæ of almost all Sphingidæ; larva of Protoparce celeus not poisonous.
- 772. [WALSH, B. D., and C. V. RILEY.] Gooseberry and currant worms. <Amer. Ent., September-October, 1869, v. 2, pp. 12-22, figs. 5-11.
 - Need of precision in nomenclature; relations and distribution of the North American species of Ribes; insect enemies of the same; natural history and description of Ellopia [= Eufitchia] ribearia, Nematus ventricosus [= ribesii], and Pristiphora grossulariæ; figures larvæ and imagos of the three species and the pupa of the Eufitchia.
- 773. [WALSH, B. D., and C. V. RILEY.] Striped cucumber beetle. .

 <Amer. Ent., September-October, 1869, v. 2, p. 24, figs. 17-19.

 Answer to inquiry of M. M. Gray; description of larva, and habits, ravages of and means against the larva and image of Diabrotica vittata; figures larva, pupa, and image of the same.
- 774. [WALSH, B. D., and C. V. RILEY.] Leafy oak gall. < Amer. Ent., September-October, 1869, v. 2, p. 25, fig. 20.

 Answer to inquiry of B. H. B.; description and figure of the gall of Cynips q.-frondosa.
- 775 [Walsh, B. D., and C. V. Riley.] Drop of gold. <Amer. Ent., September-October, 1869, v. 2, p. 25.

 Answer to inquiry of B. H. B.; egg of Citheronia regalis? on shellbark-hick-

ory; characters of eggs and oviposition of Metapodius nasulus [femoratus].

776. [WALSH, B. D., and C. V. RILEY.] The luna moth. < Amer. Ent., September-October, 1869, v. 2, p. 25.

Answer to inquiry of G. W. Kinney; characters of Attacus [= Actias] luna;

food-plants of the larva of the same.

- 777. [Walsh, B. D., and C. V. Riley.] Hag-moth larva. < Amer. Ent., September-October, 1869, v. 2, p. 25, fig. 21.
 - Answer to inquiries of C. T. Farrell and M. B. Baldwin; characters of cocoon and imago of Limacodes [= Phobetron] pithecium; number of broods of the moth in the year; figure of the larva; occurrence of Harpactor [= Milyas] cinctus in Illinois.

- 778. [Walsh, B. D., and C. V. Riley.] Stinging bug. < Amer. Ent., September-October, 1869, v. 2, p. 25.
 - Answer to inquiry of J. M. Shaffer; habits and characters of Phymata erosa.
- 779. [Walsh, B. D., and C. V. Rilley.] Pear-tree worms. < Amer. Ent., September-October, 1869, v. 2, p. 25.
 - Answer to inquiry of B. Hathaway; larva of Notodonta [= Œdemasia] concinna found on pear-tree leaves.
- 780. [WALSH, B. D., and C. V. RILEY.] "Dobson." < Amer. Ent., September-October, 1869, v. 2, p. 25.
 - Answer to inquiry of Fisherman; ignorance as to what the larva called "Dobson" is.
 - 781. [Walsh, B. D., and C. V. Riley.] White-pine weevil. <Amer. Ent., September-October, 1869, v. 2, p. 26, fig. 22.
 - Answer to inquiry of A. S. Fuller; seasons, ravages, and means against Pissodes strobi; figures larva, pupa, and image of the same.
 - 782. [Walsh, B. D., and C. V. RILEY.] Unnatural secretion of wax. <a href="#"
 - Answer to inquiry of F. Brewer; description of a case of excessive secretion of wax by Apis mellifica.
- 783. [Walsh, B. D., and C. V. Riley.] Raspberry borer. <Amer. Ent., September-October, 1869, v. 2, p. 26.
 - Answer to inquiry of F. A. Gates; ravages of the larva of Oberea perspicillata [=bimaculata] in blackberry and raspberry bushes; characters of the image of Dryocampa senatoria; food-plants of the larva of the same.
- 784. [Walsh, B. D., and C. V. Riley.] Cocoon of horn-bug. < Amer. Ent., September-October, 1869, v. 2, p. 26.
 - Answer to inquiry of A. R. McClutchen; characters of cocoon of Lnoamus dama?
- 785. [Walsh, B. D., and C. V. Rilley.] Insects named. < Amer. Ent., September-October, 1869, v. 2, p. 26, fig. 23.
 - Answer to inquiry of J. R. Muhleman; characters and figure of Amphipyra [= Pyrophila] pyramidoides; food-plants and larva of the same and of A. pyramidea of Europe; characters of the larva and image of Agnomonia anilis; supposed food-plants of its larva.
- 786. [Walsh, B. D., and C. V. Riley.] Cecropia moth caterpillar.

 Amer. Ent., September-October, 1869, v. 2, p. 26.

 Answer to inquiries of H. G. Lewelling and S. H. I. Green; characters and
 - Answer to inquiries of H. G. Lewelling and S. H. I. Green; characters and food-plants of the larva of Attacus cocropia.
- 787. [Walsh, B. D., and C. V. RILEY.] How cut worms originate. <Amer. Ent., September-October, 1869, v. 2, p. 26.
 - Answer to inquiry of T. W. Gordon; cut-worms are larvæ produced from eggs of certain *Noctuidæ*.
- 788. [Walsh, B. D., and C. V. Riley.] Red-humped caterpillar. < Amer. Ent., September-October, 1869, v. 2, p. 27, figs. 24-26.
 - Answer to inquiry of D. W. Kauffman; habits, characters, food-plants, and means against larva of *Notodonta* [= *Œdomasia*] *concinna*; figures larva, pupa, and image of the same; poisonousness of the fluids of certain insects.

- 789. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., September-October, 1869, v. 2, p. 27.
 - Answer to inquiry of T. W. G[ordon]; characters and food-plants of Thelia bimaculata and of larva of Procris [= Harrisina] americana and Eudamus tityrus; characters of the imago of the last.
- 790. [Walsh, B. D., and C. V. Riley.] Gilt gold-beetle. < Amer. Ent.; September-October, 1869, v. 2, p. 27.
 - Answer to inquiry of W. H. Martin; characters and food-plants of Chrysochus auratus.
- 791. [Walsh, B. D., and C. V. Riley.] The trumpet grape-gall. Amer. Ent., September—October, 1869, v. 2, p. 28, fig. 27.

 Answer to inquiry of D. McClaine; description and figure of galls of Cecidomyte vitielituus [= viticola]; occurrence of similar galls on several varieties
- 792. [Walsh, B. D., and C. V. Riley.] Grape-berry moth. < Amer. Ent., September-October, 1869, v. 2, p. 28.

of grape-vines and on leaves of hickory and hackberry.

- Answer to inquiry of H. C. Barnard; ravages of Penthina vitivorana [= Eudemis botrana].
- 793. [Walsh, B. D., and C. V. Riley.] Oak pruner. < Amer. Ent., September-October, 1869, v. 2, p. 28.
 - Answer to inquiry of T. J. Plumb; occurrence of *Elaphidion putator* [=rillosum] at Madison, Wis.
- 794. [WALSH, B. D., and C. V. RILEY.] Potato-bugs. < Amer. Ent., September-October, 1869, v. 2, p. 28.
 - Answer to inquiry of W. R. Shelmire; ravages and food-plants of Lytta [= Epicauta] vittata; means against potato-eating Meloidæ; characters of an unknown lepidopterous larva boring in a potato-stalk; directions for packing insects.
- 795. [WALSH, B. D., and C. V. RILEY.] Blood-sucking cone-nose. <Amer. Ent., September-October, 1809, v. 2, p. 28.

 Answer to inquiry of G. W. C.; effect of the "bite" of Conordinus sanguisuga; food-habits of the same.
- 796. [WALSH, B. D., and C. V. RILEY.] Woolly slug-like worm on apple. <Amer. Ent., September—October, 1869, v. 2, p..29.</p>
 Answer to inquiry of H. A. Green; food-plants and characters of larva of Lagos opercularis; characters of the image of the same; improper method of packing living insects.
- 797. [WALSH, B. D., and C. V. RILEY.] A water-bug. <Amer. Ent., September-October, 1869, v. 2, p. 29.

 Answer to inquiry of W. V. Smith; characters of Ranatra fusca; habits and habitat of Nepidæ.
- 798. [Walsh, B. D., and C. V. Rilley.] Goldenrod galls. < Amer. Ent., September-October, 1869, v. 2, p. 29.

 Answer to inquiry of G. W. C.; characters of galls of Trypeta solidaginis.

and Cecidomyia solidaginis.

- 799. [Walsh, B. D., and C. V. Riley.] Oak-leaf gall. < Amer. Ent., September-October, 1869, v. 2, p. 29.
 - Answer to inquiry of B. H. Broadnox; description of galls of Cecidomyia quercus-pilulæ and C. q.-symmetrica; Cynipidæ inquilinous in galls of Cecidomyidæ; differences between larvæ of Cynipidæ and Cecidomyidæ; transformations of C. q.-pilulæ and of the Cynips sp., inquilinous in its gall; distinction between groups of oaks.
- 800. [Walsh, B. D., and C. V. Riley.] Humble bees. < Amer. Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiry of C. S. Davis; number of species and distribution of the genus Bombus in North America; habits of and differences between the several forms composing a society of social insects; habits of Bombus pennsylvanicus, Halictus sp., and Andrena sp.
- 801. [Walsh, B. D., and C. V. Riley.] Can land be insured against cut-worms and other insects? <Amer. Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiry of A. Willis; means against larvæ of cut-worms.
- 802. [Walsh, B. D., and C. V. RILEY.] Beetles named. < Amer. Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiry of T. W. Hoyt, jr.; characters of Cassida [= Coptocycla] aurichalcea and Brackinus americanus; effect of the discharge made by Brackinus sp. upon the inside of the human mouth.
- 803. [Walsh, B. D., and C. V. Riley.] Royal horned-caterpillar. Amer.Ent., September-October, 1869, v. 2, p. 30.
 - Answer to inquiries of W. C. Holmes and M. G. Kern; larva of Citheronia regalis found feeding on Syringa.
- 804. [Walsh, B. D., and C. V. Riley.] Parsnip caterpillar. <Amer.
 Ent., September-October, 1869, v. 2, p. 30.
 Answer to inquiry of T. W. Hoyt, jr.; characters of larva of Papilio asterias.
- 805. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., September-October, 1869, v. 2, p. 31.
 - Answer to inquiry of C. P. Faulkner; identification of several beetles; habits of Creophilus villosus, Listotrophus cingulatus, Scarites subterraneus, and Uloma impressa.
- Soc. [Walsh, B. D., and C. V. Riley.] Beetle named. < Amer. Ent., September-October, 1869, v. 2, p. 31.

 Answer to inquiry of W. Keyes; habits and characters of larva of Calopteron
- 807. [Walsh, B. D., and C. V. RILEY.] Moth named. < Amer. Ent., September-October, 1869, v. 2, p. 31.

terminale; distinctness of C. reticulatum from C. terminale.

- Answer to inquiry of W. G. Barton; food-plant of larva and characters of image of Alaria [= Rhodophora] florida.
- 808. [Walsh, B. D., and C. V. Riley.] Worm boring into cucumber. Amer.Ent., September-October, 1869, v. 2, p. 31.
 - Answer to inquiries of G. W. C., O. L. Barler, and E. S. Smith; characters of larva and image of *Phakellura* [=*Eudioptis*] *nitidalis*; food-plants of larva; characters of undetermined larva found boring in cucumbers.

- 809. [Walsh, B. D., and C. V. Riley.] Caterpillar of the io moth. <a href="mailto: Amer. Ent., September-October, 1869, v. 2, p. 31.
 - Answer to inquiry of Mrs. Tildesley; characters of larva and image of Saturnia [= Hyperchiria] io; food-plant and urticating properties of the larva.
- 810. [Walsh, B. D., and C. V. Rilley.] Apple-tree worms. < Amer. Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of H. K. Vickroy; habits and characters of larva of Acrobasis [= Pempelia] hammondi; habits of larva of Physita nebulo [= Acrobasis indiginella].
- 811. [WALSH, B. D., and C. V. RILEY.] Stinging larvæ. <Amer. Eut., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of J. C. Falls; meaning of the vernacular and technical names of *Empretia stimulea*; urticating properties of lepidopterous larvæ.
- 812. [WALSH, B. D., and C. V. RILEY.] Lappet caterpillar on appletree. <Amer. Ent., September-October, 1869, v. 2, p. 32.</p>
 Answer to inquiry of W. Stark; characters of larva of Gastropacha americana.
- 813. [Walsh, B. D., and C. V. Riley.] Spined spider. < Amer. Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiries of G. W. Kinney and T. W. Gordon; characters and synonymy of *Epeira* [= Acrosoma] spinea.
- 814. [Walsh, B. D., and C. V. RILEY.] Dangerous looking. < Amer Ent., September-October, 1869, v. 2, p. 32.
 - Answer to inquiry of M. M. Kenzie; characters and sting of Mutilla coccinea [= Spherophthalma occidentalis].
- 815. [Walsh, B. D., and C. V. Riley.] Bag-worms again. < Amer. Ent., September-October, 1869, v. 2, p. 32.

 Answer to inquiry of T. C. Tipton; ravages of Thyridopteryx ephemeræformis;
 - harmlessness of larva of Protoparce celeus; habitat of larva of Musca demestica.
- 816. [WALSH, B. D., and C. V. RILEY.] Large water-beetle. <Amer. Ent., September-October, 1869, v. 2, p. 32.

 Answer to inquiry of S. E. Munford; secondary sexual characters in elytra

of Cybister fimbriolatus. See No. 750.

[=cunea].

- 817. [WALSH, B. D., and C. V. RILEY.] Beetles under dead fish. <Amer. Ent., September-October, 1869, v. 2, p. 32.

 Answer to inquiry of T. Ferrell; food-habits and characters of Silpha peliata
 [=americana].
- 818. [Walsh, B. D., and C. V. Riley.] Universal remedies. < Amer. Ent., November, 1869, v. 2, pp. 33-35.

 Worthlessness of any one substance as a means against insects.
- 819. [WALSH, B. D., and C. V. RILEY.] Tent-caterpillars and fall-web-worms. Amer. Ent., November, 1869, v. 2, p. 39.

 Critical review of article in Western Rural, August 26, 1869; seasons, habits, food-plants, and characters of Clisicoampa americana and Hyphantria textor

- 820. [Walsh, B. D., and C. V. Riley.] The boll worm or corn-worm. 2d article. (*Heliothis armigera*, Hübner.) <Amer. Ent., November, 1869, v. 2, pp. 42-44, fig. 29.
 - Seasons, food-plants, ravages of, and means against *Heliothie armigera*; figures of larvæ, pupa, cocoon, and imago of the same; food-plants of *Gortyna nitela*.
- 821. [WALSH, B. D., and C. V. RILEY.] Galls and their architects. 2d article. <Amer. Ent., 1869–1870, v. 2: November, pp. 45–50, figs. 30–32; December–January, pp. 70–74, figs. 45–47; February, pp. 103–106, figs. 68–71.
 - See No. 518; definition and classification of galls; descriptions and figures of galls and larvæ of Nematus salicis-pomum, Euura s.-ovum, and E. s.-gemma [=orbitalis]; habits, seasons, and descriptions of the same; habits and seasons of Anthonomus sycophanta, Batrachedra salicipomonoila, and Nematus mendicus; differences between gall-makers and guest-flies; occurrence of distinct genera of gall-insects on plants of distinct genera; descriptions and figures of galls of Cynips [= Andricus] quercus-seminator and C. q.-frondosa; synoptic table of North American genera of Cynipidæ Psenides; description of Antistrophus n. g. and of A. lygodesmiæ-pisum n. sp. and its gall; description and figure of the larva and imago of Agrilus ruficollis and its gall; habits, food-plants of, and means against, the same; figure and description of Baridius [= Ampeloglypter] sesostris and its gall; habits, food-plants of, and means against, the same; characters of Madarus ampelopsidos [= Ampeloglypter ater] and its gall; habits of Buprestidæ.
- 822. [Walsh, B. D., and C. V. Riley.] Toads in gardens. < Amer. Ent., November, 1869, v. 2, p. 50.

 Value of toads as a means against noxious insects, etc., in gardens.
- 823. [WALSH, B. D., and C. V. RILEY.] Notes on the Tarantula-killer.
 <Amer. Ent., November, 1869, v. 2, p. 52.</p>
 Comments on note of C. Peabody; occurrence of Mygale hentzii and Pepsis formosa in Missouri.
- 824. [WALSH, B. D., and C. V. RILEY.] Swarms of lady-birds. < Amer. Ent., November, 1869, v. 2, p. 55.

 Occurrence of countless millions of Coccinellidæ in England; their origin and
- movements.

 825. [Walsh, B. D., and C. V. Riley.] The squash-bug does not touch the white bush scollop. <Amer. Ent., November, 1869, v. 2,
 - p. 55.
 Coreus [= Anasa] tristis does not attack the white bush scollop variety of the squash-vine; means against the same.
- 826. [WALSH, B. D., and C. V. RILEY.] [Scientific names.] < Amer. Ent., November, 1869, v. 2, p. 57.

 Use of English and scientific names of insects.
- 827. [Walsh, B. D., and C. V. Riley.] On our table. <Amer. Ent., November, 1869, v. 2, pp. 57-58.
 - Notices of: Record of American entomology for the year 1868.—The Canadian entomologist.—The butterflies of North America, by W. H. Edwards.—Guide to the study of insects, by A. S. Packard, jr., etc.

- 828. [Walsh, B. D., and C. V. Riley.] Locust borer. < Amer. Ent., November, 1869, v. 2, p. 58.
 - Answer to inquiry of J. Bagby; Arhopalus [=Cyllene] robinia Q undistinguishable from A. [=C.] pictus Q.
- 829. [Walsh, B. D., and C. V. Riley.] Saddle-back larva. < Amer. Ent., November, 1869, v. 2, p. 59, fig. 36.
 - Answer to inquiry of G. T. Cost; figure of the larva of Empretia stimulea found on Indian corn.
- 830. [Walsh, B. D., and C. V. Riley.] Silk spiders. < Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiries of G. Howe and C. W. Spaulding; characters of Epeira [= Argiope] riparia and Nephila plumipes.
- 831. [Walsh, B. D., and C. V. Riley.] Entomological works. < Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiry of S. W. Cowles; mention of works containing descriptions of North American Coleopters and Lepidopters; food-plants of the larva of Eudryas unio.
- 832. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiry of A. H. E. Bryant; characters of the cocoon of Attacus occropia and the image of Mutilla coccinea [=Sphærophthalma occidentalis].
- 833. [Walsh, B. D., and C. V. Riley.] A new bee enemy. < Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiry of F. Brewer; characters of an undetermined carabid larva found eating Apis mellifica.
- 834. [Walsh, B. D., and C. V. Riley.] Thousand-legged worms. <Amer. Ent., November, 1869, v. 2, p. 59.
 - Answer to inquiry of J. W. Merchant; characters and poisonousness of Scolopendra castanciceps; Julus sp. and Polydesmus sp. injurious to strawberries.
- 835. [Walsh, B. D., and C. V. Riley.] Cabbage-worms. < Amer. Ent., November, 1869, v. 2, p. 60, figs. 37-38.
 - Answer to inquiry of W. C. Holmes; figures larva, pupa, and image of *Pieris* protodice; means against cabbage-worms.
- 836. [Walsh, B. D., and C. V. Riley.] The rape butterfly. Amer. Ent., November, 1869, v. 2, p. 60.
 - Answer to inquiry of J. E. Chase; occurrence of Pieris rapæ in Bangor, Me.
- 837. [Walsh, B. D., and C. V. RILEY.] Bad packing. < Amer. Ent.,

 November, 1869, v. 2, p. 60.
 - Answer to inquiry of H. C. Beardslee; characters and food-plants of *Empretia stimulea*; food-plant of *Darapsa* [= Ampelophaga] myron; directions for sending larvæ by mail.
- 838. [Walsh, B.D., and C. V. Riley.] Granddaddy long-legs. < Amer. Ent., November, 1869, v. 2, p. 60.
 - Answer to inquiry of W. R. Howard; vernacular names and habits of Phalangida.
- 839. [Walsh, B. D., and C. V. Riley.] Borer in apple-twig. <Amer. Ent., November, 1869, v. 2, p. 60.
 - Answer to inquiry of G. C. Brackett; habits of Bostrichus [= Amphicerus] bicaudatus and Elaphidion parallelum [= villosum].

- 840. [Walsh, B.D., and C. V. Riley.] Grape-vine leaf-galls. < Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of W. T. Heildrup; habits and means against *Phyllexera* vastatrix; varieties of grape infested by the same.
- 841. [Walsh, B. D., and C. V. Rilley.] Maple-worms. < Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of H. K. Vickroy; characters and ravages of *Dryocampa* rubicunda and Acronycta americana; food-plants of Teles polyphemus and Attacus cecropia.
- 842. [Walsh, B. D., and C. V. Riley.] Melancholy chafer in apples. Amer. Ent., November, 1869, v. 2, p. 61, fig. 39.

 Answer to inquiry of J. F. Fulton; figure of Euryomia [=Euphoria] melan-
 - Answer to inquiry of J. F. Fulton; figure of Euryomia [=Euphoria] melan-cholica found boring in apples.
- 843. [Walsh, B. D., and C. V. Rilley.] Worms boring in cucumbers. <Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of W. B. Ramson; food-habits of *Phacellura* [= *Eudioptis*] nitidalis.
- 844. [Walsh, B. D., and C. V. Riley.] Lilac-borer. < Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of T. J. Freeman; characters, affinities, and means against *Egeria* [= Podosesia] syringæ.
- 845. [Walsh, B. D., and C. V. Riley.] Burying beetles. < Amer. Ent., November, 1869, v. 2, p. 61.
 - Answer to inquiry of J. H. Osborn; habits of Neorophorus marginatus and of Silphida generally; characters of Hylobius stupidus [= Pachylobius picivorus] found on plum-trees.
- 846. [Walsh, B. D., and C. V. Riley.] Wire-worms in potatoes. <Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of W. R. Shelmire; characters of larva (Melanotus incertus?) boring in potatoes; food-habits and means against larvæ of Elateridæ.
- 847. [Walsh, B. D., and C. V. Rilev.] Insects named. < Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of J. F. Waters; unknown tortricid and Limacodes sp., found on apple-tree; characters of Charlesterus antennator.
- 848. [Walsh, B. D., and C. V. Riley.] Girdled pear twigs. < Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of T. A. Thorp; pear twigs girdled by Oncideres cingulata.
- 849. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of E. T. Dale; food-habits of Hippodamia glacialis, Strachia [=Murgantia] histrionica and Blepharida rhois; figure of Hippodamia glacialis.
- 850. [Walsh, B. D., and C. V. Riley.] Tomato-feeding worm. < Amer. Ent., November, 1869, v. 2, p. 62, fig. 41.
 - Answer to inquiry of A. C. Davis; description, seasons, and food-plants of larva of *Prodenia commelinæ*; figure of dorsal surface of a segment of the same.

- 851. [Walsh, B. D., and C. V. Rilley.] Cocoons of Ichneumon flies. <Amer. Ent., November, 1869, v. 2, p. 62.
 - Answer to inquiry of C. Mitchell; cocoons of Microgaster sp. found on larva of Protoparce celeus.
- 852. [Walsh, B. D., and C. V. Riley.] Gall on spotted touch-me-not. Amer. Ent., November, 1869, v. 2, p. 63, fig. 42.
 - Answer to inquiry of A. N. Prentiss; description and figures of the gall of Cecidomyia impatientis found on Impatient fulva.
- 853. [Walsh, B. D., and C. V. RILEY.] Unknown larvæ. < Amer. Ent., November, 1869, v. 2, p. 63.

 Answer to inquiry of J. M. Harold; characters of unknown larva allied to Lagoa.
- 854. [Walsh, B. D., and C. V. Riley.] Blood sucking cone nose. <Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of D. B. Watson; occurrence of Conorhinus sanguisugus at Saint Louis, Mo.
- 855. [Walsh, B. D., and C. V. Riley.] Eggs on a grape-cane. < Amer. Ent., November, 1869, v. 2, p. 63.

 Answer to inquiry of J. Cochrane; difficulty of identifying eggs of insects.
- 856. [Walsh, B. D., and C. V. Riley.] Gregarious willow-worms. , November, 1869, v. 2, p. 63.
 - Answer to inquiry of G. C. Brackett; characters, habits, and food-plants of the larva of Clostera americana [= Ichthyura inclusa].
- 857. [Walsh, B. D., and C. V. Riley.] Caterpillars named. <Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiries of G. W. Copley and W. D. Butler; food-habits of larves of Eudamus tityrus and Papilio troilus.
- 858. [Walsh, B. D., and C. V. Riley.] Insects named. < Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of L. G. Saffer; vernacular name of Mutilla coccinea [= Sphærophthalma occidentalis]; characters of Stizus [= Spheoius] speciosus.
- 859. [Walsh, B. D., and C. V. Riley.] Range of the rear-horse. <Amer. Ent., November, 1869, v. 2, p. 63.
 - Answer to inquiry of V. T. Chambers; northern range of Mantis [= Phasmomantis] carolina.
- 860. [WALSH, B. D., and C. V. RILEY.] Royal horned-caterpillar.
 - Answer to inquiry of D. L. Phares; Citheronia regalis more abundant in the South than in the North.
 - 861. [Walsh, B. D., and C. V. Riley.] Hair-worm or hair-snake. <Amer. Ent., November, 1869, v. 2, p. 64.
 - Answer to inquiry of E. H. King; Gordius sp. parasitic in the pupa of Or-chelimum sp.
 - 862. [Walsh, B. D., and C. V. Rilley.] Dahlia-stalk borer. < Amer. Ent., November, 1869, v. 2, p. 64.
 - Answer to inquiry of G. C. Broadhead; larva of Gortyna nitela boring in stalk of dahlia.
 - 863. [Walsh, B. D., and C. V. Riley.] Parsnip worm. < Amer. Ent., November, 1869, v. 2, p. 64.
 - Answer to inquiry of J. Adams; larva of Papilio asterias feeds on paranip.



U. S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

OF

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE.

BY

SAMUEL HENSHAW.

PART III.


THE MORE IMPORTANT WRITINGS

OF

CHARLES VALENTINE RILEY.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1889.

7 ENT



BIBLIOGRAPHY OF THE MORE IMPORTANT CONTRIBUTIONS TO AMERICAN ECONOMIC ENTOMOLOGY.

PART III.

864. R[ILEY], C. [V.] The house-fly. < Prairie Farmer, 2 May, 1863, [v. 27], n. s., v. 11, pp. 276-277. S.-b. No. 1, p. 9.

Stomoxys calcitrans distinguished from Musca domestica; habits of each; transformations of the latter: habit of larve of collecting underneath

transformations of the latter; habit of larvæ of collecting underneath boards.

- 865. R[ILEY], C. [V.] The May-beetle. <Prairie Farmer, 6 June, 1863, [v. 27], n. s., v. 11, p. 356. S.-b. No. 1, p. 7.</p>
 Ravages of Phyllophaga quercina [= Lachnosterna fusca]; description and habits of larvæ and imagos; remedies.
- 866. R[ILEY], C. V. Larvæ of the ten-striped spearman. < Prairie Farmer, 8 August, 1863, [v. 28], n. s., v. 12, pp. 85-86, fig. S.-b. No. 1, p. 11.

Description of eggs; figure and first description of larva of Doryphora 10-lineata; habits of larva; seasons; remedies; enemies of larva.

- 867. R[ILEY], C. V. The squash-bug. <Prairie Farmer, 8 August, 1863, [v. 28], n. s., v. 12, p. 86. S.-b. No. 1, p. 11.</p>
 Description of imagos and larvæ of Corcus [= Anasa] tristis; habits, seasons, remedies.
- 868. RILEY, C. V. The cut worm. < Prairie Farmer, 12 March, 1864, [v. 29], n. s., v. 13, p. 169.</p>
 Answer to communication of G. R. Huffman; habits and transformations of
- Agrotidida; means against them.

 869. [RILEY, C. V.] Entomological. < Prairie Farmer, 21 May, 1864,
 [v. 29], n. s., v. 13, p. 361.

Answer to communication of J. S. Lawver; ravages of unknown caterpillars, probably cut-worms, and of *Ithycerus noveboracensis* on young appletrees; means against these insects.

- 870. [RILEY, C. V.] Apple-borers. < Prairie Farmer, 14 January, 1865, [v. 31], n. s., v. 15, p. 21, 5 figs. S.-b. No. 1, p. 30.
 - Description and figure of larva and imago of Buprestis [= Chrysobothris] femorata; injury done by the larva; remedies. Figure of imago of Saperda bivittata [= candida].

- 871. [RILEY, C. V.] Peach-tree borers. < Prairie Farmer, 25 February, 1865, [v. 31], n. s., v. 15, pp. 122-123, 6 figs. S.-b. No. 1, pp. 30-31.
 - Descriptions and figures of larva, pupa, cocoon, and 3 and 2 images of *Ege-ria* [= Sannina] exitiosa; description of egg; seasons and habits of larva and image; remedies. Figure of image of Dicerca divarioata.
- 872. [RILEY, C. V.] Entomology. <Prairie Farmer, 22 April, 1865, [v. 31], n. s., v. 15, p. 306.
 S. b. No. 1, pp. 32-33.
 Coming forth of insects in spring; desirability of observing noxious insects and of recording observations; abundance of eggs of Orgyia this season; methods of destroying them.
- 873. [RILEY, C. V.] Flea-beetles and Curculio. <Prairie Farmer, 27 May, 1865, [v. 31], n. s., v. 15, p. 418, fig. S.-b. No. 1, p. 33. Prevalence of and means against *Halticida*; habits of and means against *Conotrachelus nenuphar*; figure of pupa.
- 874. [RILEY, C. V.] Apple-tree caterpillars. < Prairie Farmer, 3 June, 1865, [v. 31], n. s., v. 15, pp. 437-438. S.-b. No. 1, p. 34.

 Answer to communication from J. C. Brown; descriptions of larva, pupa, and imago, of Chatochilus pometellus; habits of the same; other caterpillars injurious to the leaves of apple trees and means against them.
- 875. [RILEY, C. V.] Curculio catcher. < Prairie Farmer, 10 June, 1865 [v. 31], n. s., v. 15, p. 457, fig.

 Figure of E. S. Hull's Curculio catcher, with explanatory text.
- 876. [RILEY, C. V.] The army-worm. < Prairie Farmer, 8 July, 1865, [v. 32], n. s., v. 16, pp. 3-4, 3 figs. S. b. No. 1, pp. 36-37.

 Answer to communication from W. R.; descriptions and figures of larva, pupa, and image of Leucania unipuncta; number of broods unknown; remedies.
- 877. [RILEY, C. V.] The currant-worm. <Prairie Farmer, 15 July, 1865,
 [v. 32], n. s., v. 16, p. 27, 4 figs. S.-b. No. 1, p. 36.
 Description of eggs; descriptions and figures of larva, pupa, and image of Eufitchia ribearia; habits, seasons, remedies.
- 878. [RILEY, C. V.] Swallows. < Prairie Farmer, 15 July, 1865, [v. 32], n. s., v. 16, p. 27. S.-b. No. 1, p. 36.
 Usefulness of swallows in the destruction of noxious insects.
- 879. [RILEY, C. V.] Another insect friend. <Prairie Farmer, 22 July, 1865, [v. 32], n. s., v. 16, p. 50. S.-b. No. 1, p. 35.

 Rogas n. sp. parasitic on larva of Sesia pelasgus? [= Hemaris thysbe]; economic benefits of parasitism.
- 880. [RILEY, C. V.] Singular caterpillar. <Prairie Farmer, 22 July, 1865, [v. 32], n. s., v. 16, p. 50. S.-b. No. 1, p. 35.

 Descriptions of larva, pupa, and image of Notodonta [= Colodasys] unicornis; habits; food-plants.
- 881. [RILEY, C. V.] Collecting and preserving insects. < Prairie Farmer, 5 August, 1865, [v. 32], n. s., v. 16, pp. 101-102, fig. Interest and importance of entomology; directions for the construction of nets and cabinets, and for the capture, killing, preparation, and preservation of insects.

- 822. RILEY, C. V. The grass-bug and its habits: Currant worms. <Cultivator and Country Gentleman, 10 August, 1865, v. 26, p. 98. Reprint: <Boston Cultivator, 19 August, 1865, v. 37, p. 259. S.-b. No. 1, p. 60.
 - Habits of Cercopididæ; nature of the image of current-worms.
- 883. [RILEY, C. V.] Apple plant-louse. <Prairie Farmer, 19 August, 1865, [v. 32], n. s., v. 16, p. 127. S.-b. No. 1, p. 40.
 - Answer to inquiry of J. Doron; habits and fecundity of Aphie mali; means against it.
- 884. R[ILEY], C. [V.] Seventeen-year locust. < Prairie Farmer, 19 August, 1865, [v. 32.], n. s., v. 16, p. 127. S.-b. No. 1, pp. 40-41. Agrees with 8. P. G. in doubting that Cicada [= Tibicen] septendecim lives seventeen years immature, and gives reasons for his doubt.
- 885. [RILEY, C. V.] Tobacco-worm. <Prairie Farmer, 2 September, 1865, [v. 32], n. s., v. 16, p. 165. S.-b. No. 1, p. 41.

 Answer to inquiry of A. B. Knowlton; habits of Macrosila quinquemaculata [= Protoparce celeus].
- 886. [RILEY, C. V.] The chinch-bug. <Prairie Farmer, 9 September, 1865, [v. 32], n. s., v. 16, p. 190. S.-b. No. 1, pp. 50-51.

 Criticism of communication by D. H. Sherman; Blissus leucopterus does not always deposit its eggs on grain of wheat, if it ever does so; proposed remedy therefore inefficient.
- 887. RILEY, C. V. The sheep gad-fly. <Prairie Farmer, 14 October, 1865, [v. 32], n. s., v. 16, pp. 288-289, figs. 1-6. S.-b. No. 1, pp. 48-49.
 - Answer to inquiries of a subscriber; figures and descriptions of larva, pupacase, and imago of Cephalemyia [= Estrus] ovis; habits of and means against it; imago viviparous.
- 888. [RILEY, C. V.] Chinch-bug not in seed grain. <Prairie Farmer, 21 October, 1865, [v. 32], n. s., v. 16, p. 308. S.-b. No. 1, p. 4. Supposed evidence that the eggs of Blissus leucopterus are not deposited and do not winter in seed wheat.
- 889. [RILEY, C. V.] Lice on calves. <Prairie Farmer, 13 January, 1866, [v. 33], n. s., v. 17, p. 24.

 Directions, on the authority of Robert Jennings, for freeing calves of lice.
- 890. [RILEY, C. V.] The chinch-bug once more. A reply to D. H. Sherman. <Waukegan [Ill.] Gazette, 20 January, 1866, v. 16, No. 18, p. 4.
 - Critical review of article of D. H. Sherman.
- 891. RILEY, C. V. The chinch-bug. <Prairie Farmer, 3 March, 1866,
 [v. 33], n. s., v. 17, p. 133. S. b. No. 1, p. 71.
 Answer to inquiry of W. R. Everett; means against Blissus leucopterus; habits, hibernation, and oviposition.
- 822. [RILEY, C. V.] Wire-worms. < Prairie Farmer, 3 March, 1866, [v. 33], n. s., v. 17, p. 133, figs. 4-5. S.-b. No. 1, p. 71.

 Answer to inquiry of Creswell; means against larve of Elaterida.

- 893. R[ILEY], C. V. Entomological. < Prairie Farmer, 24 March, 1866, [v. 33], n. s., v. 17, p. 192, figs. 5-6. S.-b. No. 1, p. 68.
 - Answers to inquiries of E. T. Nelson, J. Doron, and H. W. Shore; habits and figures of *Amphicorus bicaudatus*; remedies. Apple-twigs pierced by one of the *Uroceridæ*.
- 894. RILEY, C. V. The chinch-bug. <Pract. Ent., 26 March, 1866, v. 1, pp. 47-48.
 - Criticism of D. H. Sherman's communication on this insect; Blissus leucopterus does not oviposit in the fuzzy end of wheat-grain; brief life-history.
- 895. RILEY, C. V. The chinch-bug once more. < Waukegan [Ill.] Gazette, 31 March, 1866, v. 16, No. 28, p. 4. S.-b. No. 1, p. 70. Reply to D. H. Sherman's second communication; Blissus leucopterus does not deposit its eggs in the grain of wheat; means against it.
- 896. [RILEY, C. V.] Bark-lice remedy. < Prairie Farmer, 7 April, 1866, [v. 33], n. s., v. 17, p. 229. S.-b. No. 1, p. 79.
 Answers to inquiries of a subscriber and of S. J. B.; means against Mytilespis pomicorticis [= pomorum].
- 897. [RILEY, C. V.] "Bug" on melon, etc. < Prairie Farmer, 7 April, 1866, [v. 33], n. s., v. 17, p. 229. S. b. No. 1, p. 79.

 Answer to inquiry of M. E. W.; means against Diabrotica vittata.
- 898. [RILEY, C. V.] Warbles. < Prairie Farmer, 21 April, 1866, [v. 33], n. s., v. 17, p. 276.
 - The occurrence of lumps on the back of cattle caused by larvæ of Hypoderma bovis is not unusual or alarming.
- 899. RILEY, C. V. The ailanthus silk-worm. <Prairie Farmer, 28
 April, 1866, [v. 33], n. s., v. 17, p. 289, fig. S.-b. No. 1, p. 93.
 Partial reprint: <New York Tribune, 29 May, 1866, v. 26, No. 7844, p. 7. S.-b. No. 1, p. 92.
 - Describes attempts to introduce Bombyx arrindia and Samia [=Attacus] cynthia into Europe for commercial purposes. Describes and figures egg, larva, and imago, and figures pupa of Samia [=Attacus] cynthia; habits, seasons, and hardiness, and disadvantages for purposes of silk culture of this species; superiority of native species.
- 900. [RILEY, C. V.] Lice on pigs. < Prairie Farmer, 28 April, 1866, [v. 33], n. s., v. 17, p. 292.

 Means against.
- 901. RILEY, C. V. Novel facts about cut-worms. <Prairie Farmer, 2 June, 1866, [v. 33], n. s., v. 17, pp. 371, 372, figs. 3-8. S. b. No. 1, p. 89.
 - Climbing habits of the larvæ of Agrotididæ; figures and descriptions of the larvæ of Agrotis cochranii [= A. messoria], A. clandestina, and f Hadena subjuncta; enemies of cut-worms.
- 902. [RILEY, C. V.] The canker-worm. <Prairie Farmer, 16 June, 1866, [v. 33], n. s., v. 17, p. 412. S. b. No. 1, p. 96.

 Answer to inquiry of a subscriber; means against Anisopteryx.
- 903. R[ILEY], C. V. Large fish-fly. < Prairie Farmer, 16 June, 1866, [v. 33], n. s., v. 17, p. 412. S. b. No. 1, p. 96.

Answer to inquiry of A. B. Knowlton; habits and distribution of Perla dorsata.

- 904. RILEY, C. V. The bee-moth. <Cultivator and Country Gentleman, 21 June, 1866, v. 27, p. 399. S.-b. No. 1, p. 97.

 Answer to inquiry of E. S. Fowler; habits of Galleria cereana.
- 905. RILEY, C. V. The potato-bug. <Prairie Farmer, 23 June, 1866, [v. 33], n. s., v. 17, p. 432. S.-b. No. 1, p. 97.</p>
 Answer to inquiry of J. D. Ellington; eastward spreading of and means against Doryphora decemlineata.
- 906. [RILEY, C. V.] Army-worm. < Prairie Farmer, 23 June, 1866, [v. 33], n. s., v. 17, p. 432. S.-b. No. 1, p. 97.

 Occurrence of Lewania unipuncta in various parts of Illinois.
- 907. RILEY, C. V. White willow insects. < Prairie Farmer, 30 June, 1866, [v. 33], n. s., v. 17, p. 452. S.-b. No. 1, p. 101.

 Answer to inquiry of L. G. H.; descriptions of larva, pupa, and image of Vancesa antiopa; description of larva of Nematus ventralis; habits of and means against these.
- 908. RILEY, C. V. Still they come. < Prairie Farmer, 30 June, 1866, [v. 33], n. s., v. 17, p. 452. S.-b. No. 1, p. 101.

 Answer to inquiry of S. D.; means against Doryphora decemlineata.
- 909. RILEY, C. V. The worm question. < Ohio Farmer, 7 July, 1866,
 v. 15, p. 209. S.-b. No. 2, p. 17.
 - Comments on a controversy between J. K. and J. Brocket concerning *Pyrrharctia isabella*; molting, pupation, and cocoon-making of this and other caterpillars.
- 910. RILEY, C. V. Army-worm and canker-worm wisdom. < Prairie Farmer, 21 July, 1866, [v. 34], n. s., v. 18, p. 38. S.-b. No. 1, p. 101; No. 2, p. 40.
 - Comments on entomological ignorance amongst members of the Fruit Grower's Society of Western New York; efficacy of means against Anisoptoryx.
- 911. R[ILEY], C. V. [Attacus cecropia.] < Prairie Farmer, 21 July, 1866, [v. 34], n. s., v. 18, p. 40. S.-b. No. 1, p. 104; No. 2, p. 40. Answer to inquiry of F. T.; habits of Attacus cecropia; its usefulness for silk.
- 912. R[ILEY], C. V. [Cantharis cinerea.] < Prairie Farmer, 21 July, 1866, [v. 34], n. s., v. 18, p. 40. S.-b. No. 1, p. 104; No. 2, p. 40. Answer to inquiry of W. D. Hoord; criticizes the figure of Lytta fabricii [=Macrobasis unicolor] in Harris' Insects; usefulness of the Meloida as vesicants; habits of and means against M. unicolor.
- 913. RILEY, C. V. [Edema albifrons.] < Prairie Farmer, 1 August 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.

 Answer to inquiry of H. Kennedy; probability of unknown notodontoid proving Edema albifrons.
- 914. RILEY, C. V. Practical entomology in reality. < Maine Farmer, 2 August, 1866. S.-b. No. 2, p. 21.

 Critical review of Breekett's Practical Entomology No. 9: some cut-worms.
 - Critical review of Brackett's Practical Entomology No. 9; some cut-worms do climb trees; *Œstrus ovis* is viviparons.
- 915. RILEY, C. V. [Clytus speciosus.] < Prairie Farmer, 4 August, 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.

 Answer to inquiry of C. L. Edwards; habits of and means against Clytus

[=Plagionolus] speciosus injuring maple-trees; description of larva of Anisota [= Dryocampa] rubicunda which feeds upon maple leaves.

- 916. RILEY, C. V. Grape-leaf louse. < Prairie Farmer, 4 August, 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.

 Answer to inquiries of "Subscriber"; description, habits and development of Pemphigus vitifolia [= Phylloxera vastatrix].
- 917. RILEY, C. V. White willow worm. < Prairie Farmer, 4 August, 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.

 Descriptions of larva and image of Nematus ventralis; food-plants; ravages and number of broods.
- 918. RILEY, C. V. Insects in timber. < Prairie Farmer, 4 August, 1866, [v. 34], n. s., v. 18, p. 73. S.-b. No. 1, p. 110.

 Unknown species of *Tortricidæ* injurious to forests in Brown and Schuyler Counties, Illinois, in 1865.
- 919. RILEY, C. V. Der neue Kartoffel Käfer. < Deutsche Prairie Farmer, August, 1866, 2 figs. S.-b. No. 2, p. 14.

 Descriptions and figures of larva and image of Doryphora decemlineats; its eastern migrations; habits; means against it:
- 920. RILEY, C. V. Cicadas and walking-sticks. <Prairie Farmer, 1 September, 1866, [v. 34], n. s., v. 18, p. 136. S.-b. No. 2, p. 41. Answer to inquiry of J. D. Swain; condensed account of Cicada canicularis [=tibicen]; comparison with C. [= Tibicen] septendecim; mentions Spectrum [= Diapheromera] femoratum.
- 921. [RILEY, C. V.] Locust-borer. <Prairie Farmer, 1 September, 1866, [v. 34], n. s., v. 18, p. 140.</p>
 Answer to inquiry of O. H. S.; nearly all of the locust groves in the West have been destroyed by Clytus [= Cyllene] robiniæ; no remedy known.
- 922. RILEY, C. V. Apple-tree caterpillars. < Prairie Farmer, 8 September, 1866, [v. 34], n. s., v. 18, p. 152. S.-b. No. 2, p. 41.

 Answer to inquiries of S. Caverno and H. M. Lyman; habits, descriptions of larvæ, and images of Datana ministra and Notodonta [= Œdemasia] concinna; means against them; description of pupa of Datana ministra.
- 923. RILEY, C. V. Joint-worm. <Prairie Farmer, 8 September, 1866, [v. 34], n. s., v. 18, p. 152. S.-b. No. 2, p. 41.</p>
 Answer to inquiry of G. W. Conklin; habits of and means against Isosoma hordei.
- 924. RILEY, C. V. [Nematus ventralis.] < Prairie Farmer, 8 September, 1866, [v. 34], n. s., v. 18, p. 152.

 Answer to inquiry of G. E. W.; ravages of Nematus ventralis in Kalmar, Minn.
- 925. [RILEY, C. V.] Ten-lined potato-beetle. < Prairie Farmer, 8 September, 1866, [v. 34], n. s., v. 18, p. 152. S.-b. No. 2, p. 42.

 Larva of Loma trilineata on potato plants in Maine mistaken for that of Doryphora decembineata.
- 926. [RILEY, C. V.] Locusts. < Prairie Farmer, 3 November, 1366, [v. 34], n. s., v. 18, p. 290.</p>
 Occurrence and ravages of Calopterius spretus in portions of Kansas and western United States in 1866.
- 927. RILEY, C. V. Caterpillars on the pine. <Prairie Farmer, 10 November, 1866, [v. 34], n. s., v. 18, p. 301. S. b. No. 2, p. 42.

 Answer to inquiry of N. R. Strong and A. L. Jones; detailed description of larva of Lophyrus abbotii; habits of larva; means against the insect.

- 928. RILEY, C. V. Elm- and pear-tree borer. <Prairie Farmer, 10 November, 1866, [v. 34], n. s., v. 18, p. 301. S.-b. No. 2, p. 42.</p>
 Answer to inquiry of James Matteson; description and habits of Tremex columbs.
- 929. RILEY, C. V. Grasshoppers and locusts. < Prairie Farmer, 24
 November, 1866, [v. 34], n. s., v. 18, p. 333. S.-b. No. 2, pp. 4344.
 - Answer to inquiry of J. N.; distinguishes grasshoppers from locusts; characterizes Achetadæ [= Gryllidæ], Gryllidæ [= Locustidæ], and Locustidæ [= Acrididæ]; oviposition, transformations, ravages, and migrations of Acrididæ; recognition of Caloptenus spratus.
- 930. RILEY, C. V. Black-knot once more. <Gardeners' Mo. and Hortic., November, 1866, v. 8, pp. 331-332. S.-b. No. 2, p. 31. Fungoid origin of black-knot [Sphæria morbosa]; larvæ of Construchelus nenuphar present in the swellings.
- 931. R[ILEY], C. V. Brimstone for borers. <Prairie Farmer, 8 December, 1866, [v. 34], n. s., v. 18, p. 365. S.-b. No. 2, p. 44.

 Criticism of a communication by W.; sulphur inserted in trunks of trees ineffectual as a means against insects.
- 932. RILEY, C. V. The wire-worm. < Cultivator and Country Gentleman, 27 December, 1866, v. 28, p. 414. S.-b. No. 2, p. 34. Critical review of J. D. Gros' "The wire-worm;" myriapods and the larve of Elaterides are called wire-worms; separable by the number of their legs; means against them.
- 933. [RILEY, C. V.] Tilden tomato and the tobacco-worm. < Prairie Farmer, 5 January, 1867, [v. 35], n. s., v. 19, p. 5.

 Remarks on communication of T. B.; tomato plants eaten by tobacco-worms [Protoparos carolina] in preference to tobacco plants; possibly the worms observed were tomato-worms [P. oeleus].
- 934. [RILEY, C. V.] Remarks on Saperda, Chrysobothris, Carpocapsa, and Conotrachelus. < Prairie Farmer, 12 January, 1867, [v. 35], n. s., v. 19, p. 23.
 - Report of remarks made at the annual meeting of Southern Illinois Fruit Growers' Association.
- 935. [RILEY, C. V.] Bark-lice. < Prairie Farmer, 12 January, 1867, [v. 35], n. s., v. 19, p. 24.

 Washing apple-trees with lye effective against bark-lice.
- 936. RILEY, C. V. Insects in the flower garden. A troublous time. <Prairie Farmer, 19 January, 1867, [v. 35], n. s., v. 19, p. 37. S.-b. No. 2, p. 45.
 - Answer to communication by Kate Sherman; describes larva of [Heliothis phlogophagus] injuring Phlox.
- 937. [RILEY, C. V.] Salt and vinegar for insects. < Prairie Farmer, 19 January, 1867, [v. 35], n. s., v. 19, p. 37. S.-b. No. 2, p. 45. Review of T. Glover's Report of the U. S. Entomologist for 1865, T. Glover's Entomological exhibition in Paris, and W. C. Lodge's Fruits and fruit trees of the Middle States; use of salt and vinegar as means against insects; criticism of errors. See Nos. 222, 939, 942.</p>

. • • -

U. S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

OF

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE.

BY

SAMUEL HENSHAW.

PART III.

THE MORE IMPORTANT WRITINGS

OF

CHARLES VALENTINE RILEY.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1889.

7 ENT

- 958. [RILEY, C. V.] The Curculio. < Prairie Farmer, 1 June, 1867, [v. 35], n. s., v. 19, p. 368. S. b. No. 2, p. 73.
 - Successful results from the use of machines for destroying Conotrachelus nenuphar; this insect hibernates in the image state.
- 959. RILEY, C. V. Fifteen-spotted lady-bird. <Prairie Farmer, 8_ June, 1867, [v. 35], n. s., v. 19, p. 381.
 - Answer to inquiry of A. B. Knowlton; food, variable coloration of imago, and usefulness of Mysia [Anatis] 15-punctata.
- 960. RILEY, C. V. Cherry Aphis. < Prairie Farmer, 8 June, 1867, [v. 35], n. s., v. 19, p. 381.
 - Answer to inquiry of G. Lee; habits, ravages of and means against Mysus cerasi.
- 961. RILEY, C. V. Tree-cricket. < Prairie Farmer, 8 June, 1867, [v. 35], n. s., v. 19, p. 381.
 - Eggs of *Ecanthus niveus* deposited in raspberry canes cause the death of the wood above them.
- 962. RILEY, C. V. Apple-tree borer. < Prairie Farmer, 8 June, 1867, [v. 35], n. s., v. 19, p. 381.
 - Answer to inquiry of A. B. Campbell; commends D. B. Wier's remedy against the apple-tree borer; Saperda bivittata [= candida] goes through its transformations in two years, though commonly believed to require three years.
- 963. RILEY, C. V. Insects affecting apple-tree roots. < Prairie Farmer, 15 June, 1867, [v. 35], n. s., v. 19, p. 397. S.-b. No. 2, p. 100.
 - Letter from O. B. Galusha, with answer; descriptions of young and mature wingless individuals of the root-inhabiting form, Pemphigus pyri, of Schizoneura lanigera; habits, seasons, and reproduction of this form; its gallmaking and means against it; description of larva of Cecidomyia species found with the lice; and of larva and image of Mycetophila persica [n. sp. ?]; seasons of the latter; larva of Saperda calcarata bores in roots of apple; larva of Helops micans feeds on decaying apples; Julus [Spirobolus] marginatus infested with Gamasus juloides.
- 964. RILEY, C. V. A chapter on cut worms. <Prairie Farmer, 22 June, 1867, [v. 35], n. s., v. 19, pp. 413-414, 7 figs. S.-b. No. 2, pp. 79-80.
 - Extract from J. Townley's "Do cut-worms destroy tree buds?". Buds of fruit trees destroyed by the larvæ of Agrotididæ; other ravages and means against the same; descriptions of the larvæ of Agrotis subgothica [= A. herilis and A. tricosa], A. telifera [= A. ypsilon], and Celæna [= Hadena] renigera; description of A. cochranis n. sp. [= A. messoria]; figures larva and imago of A. cochranis, A. telifera, and Celæna renigera; and imago of A. subgothica; habits, seasons, and vernacular names of Agrotididæ.
- 965. [RILEY, C. V.] Strawberry-worm. < Prairie Farmer, 22 June, 1867, [v. 35], n. s., v. 19, p. 414. S. b. No. 2, p. 80.
 - Answer to inquiry of A. R. Whitney; geographical distribution of *Emphytus* [= Harpiphorus] maculatus.
- 966. RILEY, C. V. Insects stripping the bur-oak. <Prairie Farmer, 13 July, 1867, [v. 36], n. s., v. 20, p. 21. S.-b. No. 2, p. 81.

 Answer to inquiry of A. B. Price; habits of Lacknosterna pilosicollis [= tristis.]

- 967. RILEY, C. V. Borers. < Prairie Farmer, 13 July, 1867, [v. 36], n. s., v. 20, p. 21. S.-b. No. 2, p. 81.
 - Answer to inquiry of J. Wentworth; means against Clytus [= Cyllene] robinia and Xyleutes [= Cossus] robinia.
- 968. RILEY, C. V. Potato-beetle. < Prairie Farmer, 13 July, 1867, [v. 36], n. s., v. 20, p. 21. S.-b. No. 2, pp. 81-82.
 - Answer to inquiries of J. L. W. and G. L. Merriwether; Doryphora 10-lineata three-brooded; its occurrence at Shipman, Ill.
- 969. [RILEY, C. V.] Smith's patent Curculio trap. <Prairie Farmer, 13 July, 1867, [v. 36], n. s., v. 20, p. 21. S.-b. No. 2, p. 82. Communication from N. C. Coffman, with reply; condemns J. Smith's Curculio trap.
- 970. RILEY, C. V. [Aphis ribis.] < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81.

 Answer to inquiry of C. J. Eckhart; means against Mysus ribis.
- 971. RILEY, C. V. Mantis carolina. < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81.

 Answer to communication of J. H. Graves; eggs of Mantis [= Phasmomantis] carolina found in Ogle County, Ill.
- 972. RILEY, C. V. Lappet caterpillars on the apple. <Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81.

 Answer to communication of Perkins and Congdon; description of the larva, pupa, cocoon, and image of Gastropacha [= Tolype] velleda.
- 973. RILEY, C. V. Apple-bark lice on pears. < Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81.

 Mytilaspis pomicorticis [= pomorum] on the fruit of pear-trees; said to occur on currents, plum, and black oak.
- 974. RILEY, C. V. Currant bush borer. <Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81.

 Answer to inquiry of H. B. Kinne; habits, description, and means against Egeria tipuliformis.
- 975. RILEY, C. V. Apple-leaf crumpler. <Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S. b. No. 2, p. 81.

 Answer to inquiry of H. C. Clock; habits and means against Physica nebulo [= Aorobasis indiginella].
- 976. RILEY, C. V. Tomato-stalk borer. <Prairie Farmer, 3 August, 1867, [v. 36], n. s., v. 20, p. 69. S.-b. No. 2, p. 81.

 Answer to inquiry of O. G. Nevins; food-plants of Gortyna nitela.
- 977. RILEY, C. V. A few errors corrected. <Wisc. Farmer, 17 August, 1867. S.-b. No. 2, p. 83.

 Habits of Doryphora 10-lineata and of Carpocapsa pomonella.
- 978. RILEY, C. V. Curculio. < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.

 Answer to inquiry of J. Shearer; means against Conotrachelus nenuphar.
- 979. RILEY, C. V. Hop-vine caterpillars. < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.

 Communication from J. H. Graves, with answer; Mantis [= Phasmomantis]

carolina breeds as far north as Ogle County, Ill. Description of larva and imago of Hypena humuli [= scabra]; means against it.

- 980. RILEY, C. V. Wheat-worms. < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.
 - Communication from C. Campbell, with answer; description of larva and image and habits of *Pyralis* [= Asopia] farinalis; ravages of Tenebrio molitor; means against both species.
- 981. RILEY, C. V. [Root-borer.] < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.

 Communication from S. T. Kelsey, with answer; ravages of an unknown cerambyoid larva which destroys the roots of various plants.
- 982. RILEY, C. V. Bark-lice on the pear. <Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.

 Letter from T. D. Plumb, with answer; Mytilaspis pomicorticis [= pomorum] lives on fruit of pear and crab-apple; the scale a secretion; extent of injuries caused by this insect.
- 983. RILEY, C. V. A nuisance made useful. < Prairie Farmer, 7 September, 1867, [v. 36], n. s., v. 20, p. 148. S.-b. No. 2, p. 82.

 Means against Doryphora 10-lineata discredited.
- 984. RILEY, C. V. Strawberry leaf-roller. <Prairie Farmer, 5 October, 1867, [v. 36], n. s., v. 20, p. 212. S.-b. No. 2, p. 100.</p>
 Answer to inquiry of N. R. Strong; Anchylopera [= Phoxopterie] fragariæ injures strawberry leaves; larvæ of Selandria [= Monostegia] rosæ also eat leaves of the strawberry.
- 985. RILEY, C. V. False caterpillars on the pine. <Prairie Farmer, 5 October, 1867, [v. 36], n. s., v. 20, p. 212. S.-b. No. 2, p. 101. Answer to inquiry of A. S. Fuller; habits and description of Lophyrus locontei.
- 986. RILEY, C. V. An unknown worm. < Prairie Farmer, 5 October, 1867, [v. 36], n. s., v. 20, p. 212. S.-b. No. 2, p. 101. Extract from Beaver Dam Citizen, with comments; habits and description of Existalis sp.
- 987. [RILEY, C. V.] Bark-louse. <Prairie Farmer, 19 October, 1867, [v. 36], n. s., v. 20, p. —. S.-b. No. 2, p. 75.

 Benzine and soap as a means against bark-lice.
- 988. [RILEY, C. V.] Editorial excursion to the Rocky Mountains. <Prairie Farmer, 7 December, 1867, [v. 36], n. s., v. 20, pp. 353–354. S.-b. No. 2, pp. 112–113.

 List of plants, insects, and mammals observed.
- 989. RILEY, C. V. Bark-lice. < Prairie Farmer, 21 December, 1867, [v. 36], n. s., v. 20, p. 389. S. b. No. 2, p. 112.

 Answer to inquiries of G. Backster and J. C. Cobbey; means against Mytilaspis pomicorticis [= pomorum].
- 990. RILEY, C. V. Root Aphis. < Prairie Farmer, 21 December, 1867, [v. 36], n. s., v. 20, p. 389. S.-b. No. 2, p. 112.

 Answer to inquiry of J. M. Jordan; means against the root and stem forms of
- Schizoneura lanigera.

 991. [RILEY, C. V.] The Colorado potato-beetle. < Prairie Farmer, 21
 December, 1867, [v. 36], n. s., v. 20, p. 389. S.-b. No. 2, p. 112.
 Communications from C. W. Murtfeldt asserting the possession of poisonous

qualities by the larvæ of Doryphora 10-lineata,

- 992. RILEY, C. V. Cut-worm. Fall and spring plowing. <Moore's Rural New Yorker, ———, 1867. S.-b. No. 2, p. 94. Criticism of article by Agricola.
- 993. RILEY, C. V. Entomology. <Prairie Farmer Annual [No. 1 for 1868], 1867, pp. 53-59, 6 figs. S.-b. No. 2, pp. 125-126; No. 14, pp. 216-219.
 - Descriptions and figures of larva, pupa, and imago of Coptocycla [= Cassida] bivittata; habits, seasons, and means against it; list of Cassidida injurious to sweet-potato. Descriptions of larva, nidus, and imago of Desmia maculalis, figures pupa; habits and seasons of and means against it. Figures Colaspis flavida; describes its supposed habits. Food-plants of Heliothis armigera. Figures larva and imago of Gortyna nitela; its food-plants, seasons, and hibernation. Figures larva and imago of Heliothis phlogophagus; synonymy, geographical distribution, and means against it; remarks on the balance of nature. Figures larva, pupa, cocoon, and imago of Asopia costalis; its synonymy and ravages.
- 994. RILEY, C. V. Bark-lice. <Prairie Farmer, 15 February, 1868, [v. 37], n. s., v. 21, p. 100. S.-b. No. 2, p. 118.
 - Answer to inquiry of J. Hawkins; natural limitation and parasites of *Mytilaspis pomicorticis* [=pomorum]; it attacks healthy and unhealthy trees; means against it; experience with washes.
- 995. RILEY, C. V. Potato beetle. <Prairie Farmer, 22 February, 1868, [v. 37], n. s., v. 21, p. 117.
 S. b. No. 2, p. 119; No. 3, p. 58.
 Answer to inquiry of F. T. Moore; invention of a machine for killing Doryphora 10-lineata; hand-picking and heavy mulching the most available remedies.
 - 996. RILEY, C. V. Apple-tree plant-lice. <Prairie Farmer, 22 February, 1868, [v. 37], n. s., v. 21, p. 117. S.-b. No. 2, p. 119; No. 3, p. 58.
 - Letter from L. M., with answer; habits of and means against Aphie mali.
 - 997. RILEY, C. V. Apple-root blight. <Prairie Farmer, 22 February, 1868, [v. 37], n. s., v. 21, p. 117. S.-b. No. 2, p. 119; No. 3, p. 58.
 - Answer to inquiry of C. S. J.; Schizoneura lanigera not the cause of "rotten root."
 - 998. RILEY, C. V. Apple leaf-crumpler. <Prairie Farmer, 22 February, 1868, [v. 37], n. s., v. 21, p. 117. S.-p. No. 2, p. 119; No. 3, p. 58.
 - Answer to inquiry of J. M. Pearson; means against Physica nebulo [= Acrobasis indiginella].
 - 999. RILEY, C. V. Tree-cricket. <Prairie Farmer, 14 March, 1868, [v. 37], n. s., v. 21, p. 164. S.-b. No. 2, p. 120; No. 3, p. 58.

 Letter from J. J. Huggins, with answer; place of oviposition of *Ecanthus niveus*; list of plants in which this species oviposits.
- 1000. RILEY, C. V. Egg-masses and cocoons on apple-trees. < Prairie Farmer, 14 March, 1868, [v. 37], n. s., v. 21, p. 164. S.-b. No. 2, p. 120; No. 3, p. 58.
 - Answer to inquiry of a correspondent; description of egg-masses and of larva of Orgyia leucostiqma; beneficial influence and means of encouraging parasites.

- 1001. RILEY, C. V. Hop insects; Hop Aphis. <Prairie Farmer, 21 March, 1868, [v. 37], n. s., v. 21, p. 184. S.-b. No. 3, p. 58.</p>
 Brief account of and means against Hypena humuli [= scabra] and Phorodon humuli.
- 1002. RILEY, C. V. Supposed eggs of the preying Mantis. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Answer to inquiry of J. H. Graves; cocoons of *Rhogas* sp. attached in a mass to twigs of an apple-tree.
- 1003. RILEY, C. V. Oak-tree borer. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Answer to inquiries of E. G. Mygatt; habits and food-plants of Xylentes [= Cosens] robinia; means against it.
- 1004. RILEY, C. V. Maple-bark lice. <Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.

 Lecanium sp. found on twigs of sugar-maple.
- 1005. RILEY, C. V. Eggs of the katydid. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Answer to inquiry of W. Colwell; habits and food-plants of Platyphyllum [= Cyrtophyllus] concavus.
- 1006. RILEY, C. V. Eggs of tree-cricket in raspberry canes. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 - Letter from E. T. Nelson, with answer; oviposition of *Coanthus niveus* in raspberry canes; means against this insect.
- 1007. RILEY, C. V. Bark-lice. <Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.</p>
 Answer to inquiry of C. D. Robinson; means against Mytilaspis pomicorticis [=pomorum].
- 1008. RILEY, C. V. Bark-lice again; the native species. <Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.</p>
 Letter from N. Coleman, with answer; comparative characters of Mytilaspis
 - Letter from N. Coleman, with answer; comparative characters of Mytilaspis pomicorticis [= pomorum] and Diaspis harristi [= Chionaspis furfurus]; food-plants of the latter.
- 1009. RILEY, C. V. Dahlia-stalk borer. <Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59. Answer to inquiry of S. M. Wierman; means against Gortyna nitela.
- 1010. RILEY, C. V. Apple-worm. < Prairie Farmer, 28 March, 1868, [v. 37], n. s., v. 21, p. 201. S.-b. No. 2, p. 121; No. 3, p. 59.
 Answer to communication of S. J. C.; Carpocapsa pomonella hibernates as a larva within the cocoon.

- 1011. RILEY, C. V. False caterpillars on the Scotch and Austrian pines. <Prairie Farmer, 2 May, 1868, [v. 37], n. s., 21, p. 285, fig.
 - Description and figures of larva, cocoon and imago, and of male antenna of Lophyrus leconici; food-plants and habits of larva; value of technical science.
- 1012. BILEY, C. V. Prevention of bark-lice. <Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60.</p>
 Means against Mytilaspis pomicorticis [= pomorum] and Physita nebulo [= Acro-

basis indiginella]; figure of the latter.

- 1013. RILEY, C. V. Beetles in stomach of meadow-lark. < Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60.
 - Fragments of Nottiglossa [= Oncometopia] undata and of Platynus excavatus or P. æruginosus found in the stomach of Sturnella magna.
- 1014. RILEY, C. V. Beetle on sugar-maple. <Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60.</p>
 Eburia quadrigeminata found under bark of a felled sugar-maple, also on honey-locust.
- 1015. RILEY, C. V. White worms in wells. <Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S. b. No. 3, p. 60.

 Fish or frogs placed in wells will free them from worms.
- 1016. RILEY, C. V. Tanzy for borers. <Prairie Farmer, 9 May, 1868,
 [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60.
 Means against borers in apple- and peach-trees.
- 1017. RILEY, C. V. Peach-borer. < Prairie Farmer, 9 May, 1868, [v. 37], n, s., v. 21, p. 301. S.-b. No. 3, p. 60.

 Refers to articles on means against *Ageria* [= Sannina] exitiosa.
- 1018. RILEY, C. V. Black grape-vine caterpillars. <Prairie Farmer, 9 May, 1868, [v. 37], n. s., v. 21, p. 301. S.-b. No. 3, p. 60. Descriptions of larva and image of Thyreus abbotic; the abundance of parasites prevents serious injury by the larva.</p>
- 1019. RILEY, C. V. The potato-beetle. <Cultivator and Country Gentleman, 21 May, 1868, v. 31, p. 378.</p>
 Criticism of article on the "Colorado potato-bug" (Cultivator and Country Gentleman, 23 April, 1868, p. 309); use of hellebore against Doryphora 10-lineata.
- 1020. RILEY, C. V. The May-beetle; white grub. <Journ. of Agric., 1868, 4 figs. S.-b. No. 3, p. 10.</p>
 Habits, food-plants, enemies, and means against Lacknosterna quercina [= fusca]; figures the larva, pupa, and image of the same.
- 1021. RILEY, C. V. Canker-worm. <Colman's Rural World, 1868.
 S.-b. No. 3, p. 16.
 Habits of and means against Anisopteryx [= Paleacrita] vernata.
- 1022. RILEY, C. V. The apple-worm or codling-moth. <Colman's Rural World, 1868, fig. S. b. No. 3, p. 16.
 Seasons of and means against Carpocapsa pomonella; figures all stages except egg.

8 ENT

- 1023. RILEY, C. V. Remedy for the apple-borer. <Colman's Rural World, 1868. S. b. No. 3, p. 16.

 Means against Superda bivittata [= candida].
- 1024. RILEY, C. V. Grape-vine hoppers. <Colman's Rural World, 1868. S. b. No. 3, p. 16.
 Tettigonia coagulata injurious to leaves of grape-vines.
- 1025. RILEY, C. V. Thousand legged worm. <Colman's Rural World, 1868. S.-b. No. 3, p. 16.

 Habits of Julus [= Spirobolus] marginatus.
- 1026. RILEY, C, V. Honey-locust seed-weevil. < Prairie Farmer, 20 June, 1868, [v. 37], n. s., v. 21, p. 397. S.-b. No. 3, p. 59. Habits, description of, and means against Spermophagus robinics.
- 1027. RILEY, C. V. Potato-beetle. <Prairie Farmer, 20 June, 1868, [v. 37], n. s., v. 21, p. 397. S.-b. No. 3, p. 59.

 Natural history of and means against Doryphora 10-lineata.
- 1028. RILEY, C. V. Large moth on apple-tree. < Prairie Farmer, 27 June, 1868, [v. 37], n. s., v. 21, p. 410. S.-b. No. 3, p. 60. The larva of Attacus cecropia spins its cocoon on apple and other trees.
- 1029. RILEY, C. V. Will unimpregnated eggs hatch? < Prairie Farmer, 27 June, 1868, [v. 37], n. s., v. 21, p. 410. S.-b. No. 3, p. 60.
 - Remarks upon the statement by H. D., of Champaign, Ill., concerning the hatching of unimpregnated eggs of *Atlacus cecropia*; mentions the families and genera of insects in which parthenogenesis occurs.
- 1030. [RILEY, C. V.] Wire-worms: Experiments in killing. < Prairie Farmer, 27 June, 1868, [v. 37], n. s., v. 21, p. 410. S. b. No. 3, p. 60.
 - Effect of several liquid preparations upon the larvæ of *Elateridæ* immersed in them; corn soaked in turpentine grew unmolested by wire-worms.
- 1031. [RILEY, C. V.] [Pear-slug and current-worm.] < Prairie Farmer, 27 June, 1868, [v. 37], n. s., v. 21, p. 410. S.-b., No. 3, p. 60. Means against Selandria [= Eriocampa] corast and Nematus ventricosus [=ri-besii].
- 1032. RILEY, C. V. What becomes of bumble-bees? <Cultivator and Country Gentleman, 2 July, 1868, v. 32, p. 18.

 Answer to Anna's "Bumble-bees;" only the queens of Bombus survive the winter; new colonies formed by hibernated queens.
- 1033. RILEY, C. V. Large gray straight-horned snout-beetle. < Prairie Farmer, 4 July, 1868, [v. 38], n. s., v. 22, p. 2-3, figs. S.-b. No. 3, pp. 31, 56.
 - Descriptions and figures of larva and image of Ithyocrus noveboracensis; method of oviposition; injuries to apple, oak, and other trees.
- 1034. RILEY, C. V. The seventeen-year Cicada. < Prairie Farmer, 4

 July, 1868, [v. 38], n. s., v. 22, p. 2. S.-b. No. 3, pp. 31, 56.

 Dates and localities of occurrences of Cicada [= Tibicen] septendecim; occurrence of C. [= T.] tredecim in Missouri.

- 1065. RILEY, C. V. Raspberry canes dying. <Prairie Farmer, 11 July, 1868, [v. 38], n. s., v. 22, p. 10, 1 fig. S.-b. No. 3, pp. 31, 56.

 Answer to inquiry of J. A. Jackson; death of raspberry canes discussed; in this case not the result of injury by insects.
- 1036. RILEY, C. V. Bag-worms. < Prairie Farmer, 11 July, 1868, [v. 38], n. s., v. 22, p. 10. S.-b. No. 3, pp. 31, 56.

 Letter from B. B. Warfield, with answer; habits, food-plants, and means against Thyridopteryx ephemeraformis; description of the female.
- 1037. RILEY, C. V. Oak and rose galls. <Prairie Farmer, 11 July, 1868, [v. 38], n. s., v. 22, p. 10. S. b. No. 3, pp. 31, 32, 56. Answer to inquiry of F. H. G.; descriptions of galls and images of Cynips [= Rhodites] bicolor on rose and of Cynips [= Andrices] seminator on white oak; occurrence of Cicada [= Tibicon] septendecim in Michigan.</p>
- 1038. RILEY, C. V. Apple-borer and root Aphis. <Prairie Farmer, 11
 July, 1868, [v. 38], n. s., v. 22, p. 10. S.-b. No. 3, pp. 32, 56.
 Answer to inquiry of W. colwell; ravages of Saperda bivittata [=candida]
 and of Schisoneura langera.
- 1039. RILEY, C. V. Evergreen plant-lice. <Prairie Farmer, 11 July, 1868, [v. 38], n. s., v. 22, p. 10. S.-b. No. 3, pp. 32, 56.

 Answer to inquiry of J. Cochrane; balsam-fir injured by Lachaus strobif; means against it.
- 1040. RILEY, C. V. Bark-lice again. < Prairie Farmer, 18 July, 1868, [v. 38], n. s., v. 22, p. 18, 2 figs. S.-b. No. 3, pp. 34, 54.</p>
 Answer to inquiries of J. Hodgeson, S. D., J. Sutherland, and J. H. G.; descriptions and figures of the scales of Mytilaspis pomicorticis [= pomorum] and Diaspis harrisii [= Chionaspis furfurus].
- 1041. RILEY, C. V. Larvæ of grape-vine flea-beetle. <Prairie Farmer, 18 July, 1868, [v. 38], n. s., v. 22, p. 18. S.-b. No. 3, pp. 34, 54. Answer to inquiry of J. A. Pettingil; leaves of grape-vine eaten by larvæ of Haltica chalybea; description of larva; habite and means against it.
- 1042. [RILEY, C. V.] A corn Curculio < Prairie Farmer, 25 July, 1868, [v. 38], n. s., v. 22, p. 26. S.-b. No. 3, p. 33.</p>
 Communication from L. V. Smith, with answer; seasons and ravages of Sphenophorus sculptilis.
- 1043. RILEY, C. V. Ephemera flies; a hard story. < Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, pp. 36, 57. Extract from Peoria (Ill.) Transcript, with letter of J. Cochrane, of Havana, Ill.; notes on the above; appearance of swarms of Palingenia [= Hexagenia] bilineals in Illinois and of other may flies in Europe; life-history of Ephemeridæ.
- 1044. RILEY, C. V. Driving potato-beetles. < Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, pp. 36.

 Doryphora 10-lineata can not be driven; Macrobasis unicolor can; care should be taken in sending the former through the mail, so as not to increase its distribution.
- 1045. RILEY, C. V. Gregarious walnut caterpillar. <Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, pp. 36, 57. Answer to inquiry of "a subscriber;" leaves of walnut destroyed by larve of Datana ministra; habits and means against the same.

- 1046. RILEY, C. V. Bugs on grape-vines. < Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S. b. No. 3, p. 36.

 Letter from J. H. Bingham, with answer; new growth of stems and leaves of grape-vines injured by Corimelana pulloaria; means against it.
- 1047. RILEY, C. V. Corn-worms. <Prairie Farmer, 15 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, p. 36.</p>
 Letter from E. Daggy, with answer; injury to maize by an undescribed species of Hadena?.
- 1048. RILEY, C. V. Large worm on apple-tree. <Prairie Farmer, 15
 August, 1868, [v. 38], n. s., v. 22, p. 50. S.-b. No. 3, p. 36.
 Answer to inquiry of J. M. Devore; description of Papilio turnus, 2, form glaucus, which pupated on trunk of apple-tree.
- 1049. RILEY, C. V. White-pine trees killed by borers. < Prairie Farmer, 26 September, 1868, [v. 38], n. s., v. 22, p. 98.
- 1050. RILEY, C. V. Swarms of butterflies. < Prairie Farmer, 26 September, 1868, [v. 38], n. s., v. 22, p. 98.
- 1051. RILEY, C. V. Worms feeding on the hawthorn. < Prairie Farmer, 26 September, 1868, [v. 38], n. s., v. 22, p. 98.
- 1052. RILEY, C. V. Twig borers, sack-bearers, etc. < Prairie Farmer. 19 December, 1868, [v. 38], n. s., v. 22, p. 194.
- 1053. RILEY, C. V. Oil beetles. < Prairie Farmer, 19 December, 1868, [v. 38], n. s., v. 22, p. 194.
- 1054. RILEY, C. V. Apple-tree caterpillars. < Prairie Farmer, 19 December, 1868, [v. 38], n. s., v. 22, p. 194.
- 1055. RILEY, C. V. Twigs punctured by periodical Cicada. < Prairie Farmer, 19 December, 1868, [v. 38], n. s., v. 22, p. 194.
- 1056. RILEY, C. V. Report of committee on entomology. <Trans. Ill. Hortic. Soc. for 1867, 1868, n. s., v. 1, pp. 105-107, figs. 1-8. Criticism of previous reports; effect of cold on Aphis mali; distribution, food-plants, and means against Doryphora 10 lineata; distinctions between D. 10-lineata and D. juncta; ravages of Lytta [= Epicauta] vittata, L. [=E.] oinerea, and L. atrata [=E. penneylvanica] on potato, and L. [=Pomphopæa] whea on fruit-trees; means against them; ravages of Hallica [= Crepidodera] cucumeris on potato; habits, ravages, and specific characters of Cassididæ found on sweet potato; Saperda bivittata [= candidu] biennial; history and development of Mytilaspis pomicorticis [=pomorum]; its food-plants, supposed hermaphroditism [parthenogenesis] and mode of production of scale of this species; difficulty of explaining the production of galls; habits, characters, food-plants, and ravages of Conotrachelus nenuphar, Carpocapsa pomonella, Hiliothie armigera, Gortyna nitela, Mameetra picta, Procris [= Harrisina] americana, Selandria vitis [= Blennocampa pygmæa], Darapsa [= Ampelophaga myron, and Desmia maculalis; notes on other insects injurious to grape, apple, and wheat, and on Mycetophila persica, Emphytus [= Harpiphorus] maculatus, Agrotis [= Hadena] devastatrix,

and Hadena juncta; note on recent writings by the author.

3

7

7

1057. RILEY, C. V. Entomology. < Prairie Farmer Annual [No. 2 for 1869], 1868, pp. 30-41, 6 figs. S.-b. No. 14, pp. 220-226.

Directions for collecting and preserving insects; descriptions of apparatus and cabinets; means against cabinet pests; descriptions and figures of all stages, except egg, of Lophyrus abbotii and L. lecontoi; habits and foodplants of and means against both species; figures of eggs, pupe, and image of Cicada [= Tibioon] septendecim and of twig punctured by image; dates and localities of occurrence of this species and of C. [= T.] tredecim; C. [= T.] cassinii a form of this insect; figures larva, puparium, and image of Cephalemyia [= Estrue] evis; habits and ravages of and means against it; image viviparous in the nostrils of sheep.

1058. RILEY, C. V. The American Meromyza, Meromyza americana, Fitch. Attacking wheat just before it ripeus. < Moore's Rural New Yorker, 30 January, 1869, v. 20, p. 71, fig. S.-b. No. 3, p. 83.

Habits and ravages of *Meromysa americana*; figures larva, pupa, and injured stalk of wheat; related European species have similar habits; remedies and natural enemies.

1059. RILEY, C. V. First annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <4th Ann. Rept. State Board of Agric. for 1868, March, 1869, pp. 182–187, 98 figs. Separate: <Jefferson City, Mo., March, 1869, pp. 182–187, 2 pl., 98 figs. Notice: <Amer. Ent., April, 1869, v. 1, p. 165.

CONTENTS.

TRODUCTORY
INODUCIÓNI
NOXIOUS INSECTS.
Two species known to occur in the United States, 7—Harris's barklouse not numerous enough to do material damage, 7. The oyster-shell bark-louse, Mytilaspis pomorum

days of the year only, 18.

1059. RILEY, C. V.—Continued.

THE PERIODICAL CICADA, Tibicen septendecim	18
Its singular life history, 18—Seventeen and thirteen year races, 19—The two races not distinct species, 19—Two distinct forms	
occurring in both broods, 20—Season of their appearance and disappearance, 22—Natural history and transformations, 22—Elevated chambers of the pupa, 22—The larvæ frequently occurring at great depth in the ground, 24—The operation of emerging from the pupa, 24—Only the males are capable of singing, 24—Trees in which the females deposit their eggs, 24—Mode of eviposition, 24—The newly hatched larva, 25—The W on the wings of the Cicada, 25—Enemics, 26—Fungus infesting the imago, 26—The sting of the Cicada, 26—Wide-spread fear of the insect on account of its supposed stinging powers, 26—Explanations of the sting, 27—Injury caused by the insect, 29—by the larva, 29—by the imago, 29—Fruitless attempts to stop the injury, 30—Chronological table of all well-ascertained broods in the United States, 30—The insect will appear during the next 17 years somewhere in the United States every year except in 1878, 41—Number of broods that will appear in the next 17 years in the different States, 42.	•
APPLE-TREE BORERS	42
The round-headed apple-tree borer, Saperda candida	42
Differences between it and the foregoing species, 46—Habits of the beetle, 47—Amount of injury caused by it, 47—Parasite attacking it, 47—Remedies, 47.	••
THE PEACH BORER, Sannina exitiosa Its nature, 47—Differences in the sexes, 48—Remedies, 48—The mounding system the best remedy, 48—Testimony as to the value of the mounding system, 48—Other remedies, 49.	47
THE PLUM CURCULIO, Conotrackelus nenuphar Difference of opinion among authors on some points in its natural history, 50—Reasons for this difference of opinion, 51—Facts in its natural history, 52—It causes the spread of the peach-rot, 52—Fruit trees attacked and those not attacked by it, 53—It may hibernate as larva or pupa, but does generally as imago, 53—Mode of egg-laying, 54—It has one annual brood, 55—Walsh's experiments to show that it is two-brooded, 55—Natural remedies, 56—No parasites known to infest it, 56—Enemies, 57; The Pennsylvania soldier-beetle and its larva, 57; Lacewing-larva, 57; The subangular ground-beetle, 58; Ground-beetle larva, probably of the Pennsylvania ground-beetle, 59—Hogs as Curculio destroyers, 59—Artificial remedies, 60—Jarring the trees the most effectual method, 60—Dr. Hull's Curculio catcher, 60—Lessons for the fruit-grower from the account of the Curculio, 62.	50

1059. RILEY, C. V.—Continued.	
THE CODLING MOTH OR APPLE WORM, Carpocapsa pomonella	62
It is common wherever apples are grown, 62—Description of the insect in its different states, 63—Its life-history, 63—Other fruits attacked by it, 64—Remedies, 65—Picking up the fallen fruit, 65—Entrapping the worms the best remedy, 66—Trimble's hay band system and how to apply it, 66—Attracting the moth by	
fires, 67.	
CUT-WORMS The natural history of twelve distinct species, 67—Definition of the term "cut-worm," 67—Habits of cut-worms, 67—Their natural history briefly given, 68—Difficulty of breeding them in captivity, 69—Climbing cut-worms, 69—Injury done by them to orchards, 69—Fruit trees and shrubs they attack, 70—They attack large trees, 71.	67
The variegated cut-worm, Agrotis saucia	72
The full-grown larva, 72—The eggs, 72—Habits of the larva, 72—Cut-worm moths deposit their eggs on the leaves and not on the ground, 73—The imago, 73—Description of the insect as larva, pups, and imago, 74.	
The dark-sided cut-worm, Agrotis messoria	74
General characters of the larva, 74—Habits of, and injury done by	13
it, 75—Description of image, 75; of the larva and chrysalis, 76.	
The climbing cut-worm, Agrotis scandens	76
Injury done by the larva, 77—General characters of the larva, 77, of the moth, 78—Description of the larva, 78; of the imago, 78.	
The w-marked cut-worm, Agrotis clandestina	79
General characters of the larva, 79—Plants it attacks, 79—Char-	
acters of the moth, 79—Description of the larva and chrysalis, 79.	
The greasy cut-worm, Agrotic ypeilon	80
The larva very variable in coloration, 80—Its injury to tomato and tobacco plants, 80—General characters of the moth, 80—Description of larva characters and images 81	
tion of larva, chrysalis, and imago, 81. The western striped cut-worm, Agrotis herilis	81
Resemblance of its larva to that of the corn rustic, 81—General characters of the worm and moth, 82—Description of the larva, 82.	01
The dingy cut-worm, Agrotis subgothica	82
The glassy cut-worm, Hadena devastatrix. Habits and general characteristics of the larva, 83—Characteristics of the moth, 84—Description of larva and chrysalis, 84.	83
The speckled cut-worm, Hadena subjuncta. Characteristics and habits of the insect, 84—Description of larva, chrysalis, and imago. 85.	84
The small white bristly cut-worm, Hadena renigera	86
Other cut-worms	87
worm, 87. The wheat cut-worm	87

120 BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY. 1059. RILEY, C. V.—Continued. CUT-WORMS-Continued. Remedies against cut-worms 89 Natural enemies, 89; Microgaster militaris, 89; Paniscus geminatus, 89; The spined soldier-bug, 89; The cut-worm lion, 89-Other enemies, 90-Artificial remedies for climbing cut-worms, 90; for common field cut-worms, 91. Insects infesting the potato 91 General remarks, 91—Number of species affecting the potato, 92. The stalk-borer, Gortyna nitela 92 Habits of the larva and imago, 92-Remedy, 93. The potato stalk-weevil, Trichobaris trinotata 93 Its geographical distribution, 93—Its habits, 93—Remedy, 95. The potato- or tomato-worm, Protoparce celeus..... 95 It can not sting with its horn, 95-Its chrysalis, 95-How the imago differs from the tobacco-worm moth, 95-Remedies and parasites, 96. Blister-beetles, Meloidæ 97 The striped blister-beetle, 96-The ash-gray blister-beetle, 97-The black-rat blister-beetle, 98—The black blister-beetle, 98— The margined blister-beetle, 98-Synonymical remarks, 98-Remedies for blister-beetles, 99. The three-lined leaf-beetle, Lema trilineata..... 99 Merdigerous habit of the larva, 99-It has two annual broods, 100—Other notes on the habits of the insect, 100. The cucumber flea-beetle, Crepidodera cucumeris 101 The Colorado potato-beetle, Doryphora 10-lineata..... 101 Its past history and future progress, 101—Its native home, 101—Its gradual spread eastward, 102-Its confusion with the bogus Colorado potato-beetle, 103-How the two species differ in habits, 104; in their larval states, 104; in the egg state, 105-Description of the larva of Doryphora juncta, 106-Differences in the imagos of the two species, 106-Habits of the Colorado potatobeetle, 107-When it appears and disappears, 107-Number of eggs laid by each female, 107-Food-plants, 107-Singular fact that D. juncta has not acquired the habit of attacking the potato, 108-Natural remedies, 109-Complicated economy of nature, 109-Decrease in the number of potato-beetles on account of increase in the number of parasites, 109-The Colorado potato-beetle parasite, 111-Its general character and habits, 111-Description of Lydella doryphoræ, 111-Lady-birds and their larvæ, 112-The spined soldier-bug, 113-The common squash-bug, erroneously considered an enemy of the potato-beetle, 113-The bordered soldier-bug, 114—The many banded robber, 114—The rapacious soldier-bug, 114-The Virginia tiger-beetle, 115-The flery ground-beetle, 115-Blister-beetles, 115-The larvæ not touched by fowl, 115-Artificial remedies, 116-Ineffectiveness

of mixtures tried, 116—Killing the beetle early in spring, 116—Pincers for crushing the insect, 116—Benson's machine, 116—Proper choice of varieties of potatoes, 117—The pest will overrun the Eastern States, 117—Carelessness in transmitting speci-

mens of the beetle, 117.

1059. RILEY, C. V.—Continued.	
The apple-root plant-louse, Schisoneura lanigera Three distinct kinds of rots affecting the roots of apple-trees, 118—The root-louse the cause of one of these rots, 118—The cause of the other rots still hidden, 119—The root-louse especially in- jurious in southerly latitudes, 119—It occurs also on other parts of the tree besides the root, 120—Description of the winged louse, 120—Fitch's description of the winged form refers to another species, 120—The root-louse belongs to the genus Eriosoma, 121—Natural enemies, 121; Chalcis-fly, 121; the root-louse Syr- phus-fly, 121; Scymnus cervicalis, 122—Artificial remedies, 123.	118
THE WOOLLY ELM TREE LOUSE, Schizoneura rileyi Its general appearance and habits, 123—Description of the winged form, 124.	123
Insects injurious to the grape-vine	124
The new grape-root borer, Prionus laticollis Reports on the damage caused by it, 124—Description of the larva, 126—It belongs probably to the cylindrical Orthosomas 126— Former accounts of the natural history of this beetle, 127—Its injury known for several years, 127—Remedies, 128.	124
The grape curculio, Craponius inæqualis	128
The grape-seed Curculio, Isosoma vitis	129
The grape-cane gall Curculio, Ampeloglypter sessetris	131
The grape-vine Fidia, Fidia viticida	132
The grape fruit-worm, Eudemis botrana Amount and extent of the injury caused by it, 133—Characteristics of the larva, 134—Transformations, 134—Description of larva, chrysalis, and imago, 135—Remedies, 135.	133
The eight-spotted forester, Alypia 8-maculata	136
The grape-vine plume, Oxyptilus periscelidactylus	137
The snowy tree-cricket, <i>Ecanthus niveus</i> Characteristics of the insects, 138—It is injurious, 138—Nature of the injury caused by it, 138—Remedy, 139.	138
THE RASPBERRY GEOMETER, Synchlora rubivoraria	139
140.	
THE GOOSEBERRY FRUIT-WORM, Dakruma convolutella	140

1059.	RILEY, C. V.—Continued.	
	THE STRAWBERRY LEAF-ROLLER, Phoxopteris fragaria	142
	Extent and nature of the damage caused by it, 142—Habits of the	
	insect, 142—Accounts of its injury in Indiana and Illinois, 142—	
	Remedy, 143—Description of the image and larva, 143.	
	THE WHITE-MARKED TUSSOCK-MOTH, * Orgyia leucostigma	144
	The egg-mass, 144—The larva and larval changes, 144—The full-	
	grown larva, 145—Habits of the larva, 145—Mode of casting off	
	the larval skin, 145—The cocoon, 146—The image, 146—Two an-	
	nual broods, 146—Food-plants, 146—Remedies, 147.	
	THE BAG-WORM, alias BASKET-WORM, alias DROP-WORM, Thyridoptoryx	1 400
	ephemoriformis	147
	Its geographical distribution, 148—Injury caused by it, 148—The egg, 148—The larva and its growth, 148—Habits of the larva,	
	149—The chrysalis, 149—The sex distinguishable in the chrysa-	
	lis state, 149—The imago, 149—Food-plants, 150—Parasites, 150;	
	Cryptus inquisitor, 150; Hemiteles thyridoptorygis, n. sp., 150—	
	• Remedies, 151.	
	THE AILANTHUS-WORM, Œta punotella	151
	Injury done to the Ailanthus tree, 151-Habits of the larva, 151-	
	The chrysalis, 151-The image, 152-Geographical distribution,	
	152-Remedy, 152-Description of larva and chrysalis, 152; of	
	the imago, 153.	
	THE WALNUT TORTRIX, Cacceoia rileyana	153
	Habits of the larva, 153—General appearance of the moth, 153—	
•	Phytophagic form of the insect on snowberry, 153—Description	
	of larva, chrysalis, and imago, 154; of the variety symphoricarpi,	
	. '154.	
	THE SHED-CORN MAGGOT, Anthomyta zeas	·154
	Accounts of damage caused by it, 154—The maggot, 155—Trans-	
	formation, 155—Description of the imago, 155—Remedy, 155—	
	Habits of Anthonyia larvæ, 156.	150
	THE WHITE GRUB, Lacknosterna fueca	156
	fect insect, 157—Résumé of its life-history, 157—Remedies, 157—	
	Regularity in the appearance of the beetle, 158—Accounts of the	
	fungus infesting the white grub, 158.	
	THE AMERICAN MEROMYZA, Meromyza americana	159
	Nature of the damage caused by it, 159—Characteristics of larva.	
	chrysalis, and imago, 160—European Diptera with similar habits.	
	160—Remedies, 161.	
	THE SHEEP BOT-FLY OF HEAD-MAGGOT, Estrus ovis	161
	The insect in its different states, 161—Its larva, 162—Pupa, 162—	
	Characteristics of the imago, 162—Fatal results of the presence	
٠	of the magget in the head of the sheep, 163—Rabbits attacked	
	by gad-fly, 164—Testimony regarding the viviparous habits of	
	the bot-fly, 164—Remedies, 165.	
	INSECT ENEMIRS OF THE HONRY-BEE	166
	The bee-moth or wax-worm, Galleria cereana	166
	General appearance of the moth, 166—There are no moth-proof	
	bee-hives, 166—Habits of the worm, 167—How its presence in the hive may be recognized, 167—Prevention and remedy, 167.	
	mive may be recognized, 10/-1 revenued and remetly, 10/.	

^{*} Reprint: <Cultivator and Country Gentleman, 14 May, 1874, v. 39, pp. 310-311, 3 figs.

BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	123
1059. RILEY, C. V.—Continued. INSECT ENEMIES OF THE HONEY-BEE—Continued. The bee-killer, Promachus apivorus. It is an Asilid fly, 168—Mr. Thompson's account of the fly, 168—How it captures and kills bees, 168—No remedy known, 168.	168
BENEFICIAL INSECTS.	
THE REAR-HORSE, alias CAMEL-CRICKET, alias DEVIL'S RIDING-HORSE, Phasmomantis carolina Its food, 169—How it grasps its prey, 169—Difference in the sexes, 170—The larva, 170—The egg-mass, 170—The mode of egg-laying, 170—Voracious disposition of the Mantis, 171—Its beneficial influence, 171—Trachina parasite of the Mantis, 171.	169
INNOXIOUS INSECTS.	
THE SOLIDAGO GALL-MOTH, Gelechia gallæsolidaginis	173

THE CHICKWEED [=KNOTWEED] GEOMETER, Hamatopis grataria.... 179

Its natural history, 179—Description of larva and chrysalis, 179.

180

1060. RILEY, C. V. Eggs of the Mantis or rear-horse < Moore's Rural New Yorker, 10 April, 1869, v. 20, p. 234, 2 figs. S.-b. No. 3, p. 52.

Answer to inquiry of T. C. Bartle; description of Mantis [= Phasmomantis] carolina; its food, habits, oviposition, and parasites; figures its egg-masses and those of Phylloptera [Amblycorypha] oblongifolia.

 RILEY, C. V. Native bark-lice on apple-trees. < Prairie Farmer, 17 April, 1869, [v. 39], n. s., v. 23, p. 122, 1 fig. S.-b. No. 3, p. 57.

Answer to inquiry of T. B. Gardner; life history of and means against Diaspis harrisis [= Chionaspis furfurus]; figure of the same.

1062. RILEY, C. V. Curculio. <Prairie Farmer, 17 April, 1867, [v. 39],
 n. s., v. 23, p. 122. S.-b. No. 3, p. 57.

Answer to inquiry of James Weed; habits of and means against Constructedus nemuphar.

1063. RILEY, C. V. Cherry-tree borers. < Prairie Farmer, 17 April, 1869, [v. 39], n. s., v. 23, p. 122. S.-b. No. 3, p. 57.

Answer to inquiry of Onargo Horticultural Society; larve of Buprestis [= Directa] divarioata and of Trockilium sp., injurious to cherry-trees; softmaple attacked by T. [= Egeria] accrni; means against these insects.

- 1064. RILEY, C. V. White-grub fungus. <Prairie Farmer, 15 May, 1869, [v. 39], n. s., v. 23, p. 154. S.-b. No. 3, p. 76.</p>
 Letter from D. W. Tindall, with answer; ravages of Lachnosterna queroina [=fusca] in Clinton County, Mo., in 1868; Torrubia growing on the larve in 1869.
- 1065. RILEY, C. V. The canker-worm, Anisopteryx vernata Peck. <Moore's Rural New Yorker, 29 May, 1869, v. 20, p. 345, figs. S.-b. No. 3, p. 73.
 - Description, figures, and natural history of canker-worms; the two species are confounded.
- 1066. RILEY, C. V. The seed-corn maggot, Anthomyia zeas, Riley. Destroying the seed after it is planted. <Moore's Rural New Yorker, June, 1869, fig. S.-b. No. 3, p. 81. Advance print, with changes: <1st Ann. Rept. State Ent. Mo., March, 1869, pp. 154-156, figs. 86-87.
 - See No. 1059 for synopsis of contents.
- 1067. RILEY, C. V. Cherry-tree plant-lice. <Moore's Rural New Yorker, 10 July, 1869, v. 20, p. 443. S.-b. No. 3, p. 83.

 Answer to inquiry of G. J. Magee; means against Aphis [= Myzus] cerasi.
- 1068. RILEY, C. V. Gooseberry span-worms. < Moore's Rural New Yorker, 10 July, 1869, v. 20, p. 443. S.-b. No. 3, p. 83.

 Letter from A. Yancey, with answer; occurrence of Euftchia ribearia in Iowa; habits and means against it; habits of and influence of weather on Blissus leucopterus; scarcity of Doryphora 10-lineata in Iowa.
- 1069. RILEY, C. V. Apple-leaf crumpler mistaken for Curculio. < Prairie Farmer, 10 July, 1869, [v. 40], n. s., v. 24, p. 218, fig. S.-b. No. 3, p. 80.
 - Letter from B. T. Taylor; improper use of the word Curculio; figures larvacase and image of *Phycita nebulo* [= Acrobasis indiginella]; means against it; outline figure of Constrachelus nenuphar.
- 1070. RILEY, C. V. Peach-tree borer. <Prairie Farmer, 10 July, 1869, [v. 40], n. s., v. 24, p. 218. S.-b. No. 3, p. 80. Answer to inquiry of C. Allen; habits of and means against Egeria [=Sannina] existosa.</p>
- 1071. RILEY, C. V. To protect plums from Curculio. Prairie Farmer, 10 July, 1869, [v. 40], n. s., v. 24, p. 218. S.-b. No. 3, p. 80. Answer to inquiry of H. H. S.; recommends the collection of the image of Conotrachelus nenuphar by jarring.
- 1072. RILEY, C. V. White-grub; information wanted. < Prairie Farmer, 28 August, 1869, [v. 40], n. s., v. 24, p. 274, 4 figs. S.-b. No. 3, p. 95.
 - Letter from A. T. V., with answer; habits of and means against Lacknosterna queroina [= fueca]; figures of larva and imago.
- 1073. RILEY, C. V. Unknown corn pest. <Prairie Farmer, 28 August, 1869, [v. 40], n. s., v. 24, p. 274. S.-b. No. 3, p. 95.</p>
 Letter from E. B. Hickey, with answer; ravages of cut-worms and of an unknown pest on maize; brief description and habits of the latter.

- 1074. RILEY, C. V. Larva of the grape vine flea-beetle. <Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S. b. No. 3, pp. 100, 103.
 - Answer to inquiry of D. D. Vosburgh; habits, ravages, and means against Graptodera [= Haltica] chalybea.
- 1075. RILEY, C. V. Rose bug. < Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, pp. 100, 103.
 - Answer to inquiry of a subscriber; description of imago, transformations of, and means against *Macrodactylus subspinosus*.
- 1076. RILEY, C. V. Large green caterpillar on the apple. <Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, p. 100.
 - Answer to inquiry of S. E. A. Palmer; descriptions of larva and image of *Attacus cecropia*; its habits and seasons.
- 1077. RILEY, C. V. Conical galls on leaves of wild grape-vine. <Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S. b. No. 3, p. 100.
 - Answer to inquiry of D. McClaine; description of galls of Cecidomyta vitisviticola; characters of the larvæ of Cecidomyta.
- 1078. RILEY, C. V. A strange bug. <Moore's Rural New Yorker, 28
 August, 1869, v. 20, p. 555. S. b. No. 3, p. 100; No. 4, p. 3.
 The insects described in North Carolina's "A strange bug," were probably

 Procus venosus; habits of the genus Procuss
- 1079. RILEY, C. V. Currant-worms and black-currants. < Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, p. 100; No. 4, p. 3.
 - Critical review of Addi on currant-worms; in North America three species of larvæ feed on the leaves of currants, and two species of borers live within the stem.
- 1080. RILEY, C. V. Curculio. < Moore's Rural New Yorker, 28 August, 1869, v. 20, p. 555. S.-b. No. 3, p. 100; No. 4, p. 3. Commendation of an editorial criticism of Northwest's article on the Curculio.
- 1081. [RILEY, C. V.] Insects injurious to the grape-vine. [No. 1.] <Amer. Ent., August, 1869, v. 1, pp. 231-234, figs. 169-173. Reprint, with slight changes: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 87-91, figs. 60-63.
 - Treats of Priorus laticollis and P. imbricornis. See No. 1127 for synopsis of contents.
- 1082. [RILEY, C. V.] Insects infesting the sweet potato. <Amer. Ent., August, 1869, v. 1, pp. 234-238, figs. 174-181. Reprint, with slight changes: <2d Ann. Rept. State Ent. Mo., March; 1870, pp. 56-64, figs. 26-38.
 - Treats of tortoise-beetles, Cassididæ. See No. 1127 for synopsis of contents.
- 1083. RILEY, C. V. The borers. < Western Rural, September, 1869. S.-b. No. 4, p. 2.
 - Means against Ægeria cucurbitæ [= Melittia ceto].

- 1084. RILEY, C. V. Supposed bark-lice eggs in Missouri. < Prairie Farmer, 4 September, 1869, [v. 40], n. s., v. 24, p. 282. S.-b. No. 3, p. 95.
 - Letter from J. Reed, with answer; eggs of a moth mistaken for those of Mytilaspis pomicorticis [= pomorum]; the latter unknown in Missouri.
- 1085. RILEY, C. V. New York weevil on apple-trees. < Prairie Farmer, 4 September, 1869, [v. 40], n. s., v. 24, p. 282, 3 figs. S_{\$\foralleq\$} b. No. 3, p. 95.
 - Letter from R. M. Guy, with answer; figures of larva and imago, description of imago, geographical distribution, methods of oviposition, food-plants, and means against *Ithycerus noveboracensis*.
- 1086. [RILEY, C. V.] Insects injurious to the grape-vine. No. 2. < Amer. Ent., September-October, 1869, v. 2, pp. 22-24, figs. 12-16. Reprint, with slight changes. <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 71-73, figs. 44-48.
 - Treats of Charocampa pampinatrix [= Ampelophaga myron]; see No. 1127 for synopsis of contents.
- 1087. RILEY, C. V. Larva of the imperial moth. <Prairie Farmer, 9 October, 1869, [v. 40], n. s., v. 24, p. 322. S.-b. No. 3, p. 92. Letter of C. H. Thayer, with answer; description of larva, pupa, and image of Coratocampa [= Eacles] imperialis; food-plants of the same and of C.
- 1088. RILEY, C. V. Apple snout beetle or four-humped Curculio. < Prairie Farmer, 9 October, 1869, [v. 40], n. s., v. 24, p. 322. S.-b. No. 3, p. 92.

[=Citheronia] regalis.

- Letter from R. W. Gandy, with answer; habite and description of Anthonomus quadrigibbus; recommends jarring.
- 1089. RILEY, C. V. That venomous potato-worm! < Moore's Rural New Yorker, 20 November, 1869, v. 20, p. —. S.-b. No. 3, pp. 107; 112.
 - Critical review of several recent articles upon tomato-worms; structure and harmlessness of larvæ of Sphingidæ; geographical distribution of Sphinz [=Protoparce] carolina and S. quinquemaculata [=P. celeus].
- 1090. [RILEY, C. V.] The bag-worm, alias basket-worm, alias drop-worm, Thyridopteryx ephemeræformis, Haw. < Amer. Ent., November, 1869, v. 2, pp. 35-38, fig. 24.
 - Geographical distribution, food-plants, seasons, transformations, parasites, copulation, and oviposition of and means against *Thyridopteryx ophemera-formis*; figures and descriptions of larva, larva-cases, pupa, and imago; description of eggs; abodes constructed by insects.
- 1091. [RILLY, C. V.] Insects injurious to the grape vine. No. 3. < Amer. Ent., November, 1869, v. 2, pp. 54-55, figs. 33-35. Reprint, with slight changes. <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 74-76, figs. 49-51.
 - Treats of Philampelus achemon; see No. 1127 for synopsis of contents.
- 1092. RILEY, C. V. The saddle-back caterpillar. < Moore's Rural New Yorker, 4 December, 1869, v. 20, p. —, fig. S.-b. No. 3, p. 103.
 - Letters from A. W. Baker and G. T. Cost, with answer; figures and characterization of the larva of *Empretia stimulea*; food-plants, urticating properties, and transformations; description of the image.

- - Answer to communication of W. J. B., concerning Cicada [= Tibicen] septendecim.
- - Occurrence of Gonicootee hologaster in great numbers on a hen; means against the same; figure of it.
- - Habits and place of oviposition of Agrotis inermis [= saucia]; description of its eggs and larvæ.
- 1097. RILEY, C. V. That glow-worm. <Cultivator and Country Gentleman, 6 January, 1870, v. 35, p. 5, fig. S.-b. No. 4, p. 23. Reprint: <Amer. Ent., October, 1880, v. 3, n. s., v. 1, p. 254, fig.
 - Figures larva and image of Photuris pennsylvanica; Photinus pyralis also luminous in larval and adult stages; comparison with Lampyris noctiluca.
- 1098. [RILEY, C. V.] In memoriam. <Amer. Ent., December, 1869— January, 1870, v. 2, pp. 65–68.
 Biographical and obituary notice of B. D. Walsh.
- 1099. [RILEY, C. V.] The harlequin cabbage-bug, Strachia histrionica; Hahn. <Amer. Ent., December, 1869-January, 1870, v. 2, pp. 79-80, fig. 56.
 - Enumeration of the enemies of the cabbage in the United States; methods of their injury; importation and spread of some species; figures of the imago; description of eggs, habits, geographical distribution, seasons, and odors of Strachia [= Murgantia] histrionica; extract from G. Lincecum's "Texan cabbage-bug."
- 1100. [RILEY, C. V.] An entomologist caught napping. <Amer. Ent., December, 1869-January, 1870, v. 2, p. 84.
 - Criticism of the view that trees, grasses, or any other particular forms of vegetation are the natural coverings of the earth; this criticism applied especially to John Curtis.
- 1101. [RILEY, C. V.] Poisonous qualities of the Colorado potato-bug. Amer. Ent., December, 1869-January, 1870, v. 2, pp. 85-86. Extract from "Winona Republican," with remarks upon the poisonous nature of the blood of *Doryphora* 10-lineata.
- 1102. [RILEY, C. V.] Insects injurious to the grape-vine. No. 4. <Amer. Ent, December, 1869-January, 1870, v. 2, pp. 89-90, figs. 58-59. Reprint, with slight changes: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 76-78, figs. 52-53.
 - Treats of Philampelus satellitia [= pandorus]; see No. 1127 for synopsis of contents.

- 1103. [RILEY, C. V.] Toads vs. bugs. < Amer. Ent., December, 1869-January, 1870, v. 2, p. 91.
 - Translation of extracts from Fogt's "Noxious and beneficial animals;" existence of a commerce in toads between France and England; usefulness, tameness, and gratitude of toads.
- 1104. [RILEY, C. V.] The tomato-worm again. <Amer. Ent., December, 1869-January, 1870, v. 2, pp. 91-92.
 - Reprint, with comments, of article entitled "The tomato-worm," from Syracuse Standard; absurd nature of newspaper accounts of insects.
- 1105. [RILEY, C. V.] Mr. Walsh's successor. < Amer. Ent., December, 1869-January, 1870, v. 2, pp. 92-93.
 - Condition in which B. D. Walsh left the office of State Entomologist of Illinois at his death; plans of his intended second report; provisions for the preparation of the same and the filling of the office.
- 1106. [RILEY, C. V.] To our subscribers. Amer. Ent., December, 1860-January, 1870, v. 2, p. 93.

 Proposed chapge in the character of the American Entomologist.
- 1107. [RILEY, C. V.] The Walsh entomological collection. < Amer. Ent., December, 1869-January, 1870, v. 2, pp. 93-94.
 - Statement of the conditions under which the Walsh collection of insects is to be sold; expression of preferences in regard to its disposition; extent and method of preservation of the collection.
- 1108. [RILEY, C. V.] A State entomologist for Minnesota. < Amer. Ent., December, 1869-January, 1870, v. 2, p. 94.
 - Commendation of resolutions passed by the Minnesota State Horticultural · Society recommending the appointment of a state entomologist; promotion of entomological studies by appropriations from the several States.
- 1109. RILEY, C. V. [Field for the entomologist in the South.] < Amer. Ent., v. 2, December, 1869-January, 1870, p. 94; February, 1870, p. 121.
 - Extent and novelty of entomological work in the southern United States; J. P. Stelle at work in this field.
- 1110. [RILEY, C. V.] On our table. < Amer. Ent., December, 1869-January, 1870, v. 2, pp. 95, 96.
 - Notices of J. T. C. Ratzeburg's works on "Forest trees" and "Weeds of Germany and Switzerland."
- 1111. [RILEY, C. V.] Information wanted. <Amer. Ent., December, 1869-January, 1870, v. 2, p. 96.
 - Answer to inquiry of M. A. Kendall; habits of Xylocopa carolina [= virginica], Rhyssa [= Thalessa] lunator, and Spectrum [= Diapheromera] femorata.
- 1112. |RILEY, C. V.] The Cecropia moth, Attacus cecropia, Linn.

 <Amer. Ent., February, 1870, v. 2, pp. 97-102, fig. 59 [bis]-67.

 Descriptions and figures of larva, cocoon and image of Attacus cecropia; figure of 'pupa; descriptions of egg and of young larva at its several stages; nomenclature, food-plants, and parasites; its value as a silk-worm; figures and descriptions of Ophion macrurum, Excrista cecropia n. sp., and Chalcis [= Spilochalcis] maria n. sp.; figure of larva of Ophion macrurum and of cocoons of Cryptus nuncius; habits of these parasites; Excrista cecropia considered a variety of E. militaris [= Nemora a leucania"].

- 1113. RILEY, C. V. Report of the committee on entomology. Read
 ... before the Missouri State Horticultural Society. < Amer.
 Ent., February, 1870, v. 2, pp. 106-109. Reprint: <2d Ann.
 Rept. State Ent. Mo., March, 1870, pp. 5-8, 13-15.
 See No. 1127 for synopsis of contents.
- 1114. [RILEY, C. V.] Silk-worm eggs. <Amer. Ent., February, 1870, v. 2, p. 109.</p>
 Seventy-eight packages of silk-worm eggs, valued at \$800 per package and weighing two tons, shipped from Yokohama to France via California and the Pacific Railroad.
- 1115. RILEY, C. V. Imported insects and native American insects. <Amer. Ent., February, 1870, v. 2, pp. 110-112, fig. 72-75. Reprint: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 8-12.

See No. 1127 for synopsis of contents.

- 1116. [RILEY, C. V.] The trumpet grape-gall, Vitis viticola O. S. — Amer. Ent., February, 1870, v. 2, pp. 113-114, fig. 76.

 Figure of the gall of Cecidomyia vitis-viticola. C. v.-lituus a synonym; occurrence of similar gall in England on Tilia probably caused by mites.
- 1117. [RILEY, C. V.] The goat-weed butterfly, Paphia glycerium Doubleday. <Amer. Ent., February, 1870, v. 2, pp. 121-123, flgs. 81-83. Reprint, with slight changes. <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 124-128, flgs. 94-96. See No. 1127 for synopsis of contents.
- 1118. [RILEY, C. V.] Insects injurious to the grape-vine. No. 5. < Amer. Ent., February, 1870, v. 2, pp. 123-124, fig. 84. Reprint, with slight changes. <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 78-79, fig. 54.

 Treats of Thyreus abbotii; see No. 1127 for synopsis of contents.
- 1119. [RILEY, C. V.] Scorpion in Kansas. < Amer. Ent., February, 1870, v. 2, p. 126.

Answer to inquiry of K. Kelsey; occurrence of Buthus carolinianus in Texas, Missouri, and Kansas.

- 1120. [RILEY, C. V.] The grain Bruchus of Europe just imported. <Amer. Ent., February, 1870, v. 2, pp. 126-127, fig. 85.
 - Answer to inquiry of A. S. Fuller; notice of the recent importation into New York of the European Bruchus granarius; danger of and remedy for such importation; habits and ravages of and means against the beetle in Europe as described in J. Curtis's, "Farm Insects;" figure of the imago and of infested beans; figures of larva, pupa, and imago of B. pisi and of infested pea.
- 1121. [RILEY, C. V.] Locust borer. < Amer. Ent., February, 1870, v. 2, pp. 127-128, figs. 86-89.
 - Answer to inquiry of J. M. Shaffer; habits, food-plants, seasons, and ravages of and means against Xyleutes [= Cossus] robinia; figures larva, pupa, and male and female images.

- 1122. [RILEY, C. V.] Eggs of snowy tree-cricket on raspberry canes. <Amer. Eut., February, 1870, v. 2, p. 128.
 - Answer to inquiry of J. B. Root; oviposition of and means against *Counthus niveus*.
- 1123. [RILEY, C. V.] Spotted rove-beetle. <Amer. Ent., February, 1870, v. 2, p. 128.
 - Answer to inquiry of J. Huggins; habits of Staphylinus maculosus and of rovebeetles in general.
- 1124. [RILEY, C. V.] Raspberry gouty gall. <Amer. Ent., February, 1870, v. 2, p. 128, fig. 90.
 - Answer to inquiry of C. Carpenter; ravages of and means against Agrilus ruficollis; description and figure of larva.
- 1125. [RILEY, C. V.] Parasitic cocoons. < Amer. Ent., February, 1870, v. 2, p. 128, fig. 91.
 - Answer to inquiry of G. C. Brackett; figures cocoons and conjectures as to the habits of [Microplitis ceratomiæ var. actuosus].
- 1126. [RILEY, C. V.] The pigeon Tremex in apple. <Amer. Ent., February, 1870, v. 2, p. 128.
 - Answer to inquiry of J. Huggins; habits and food plants of Tremex columba; description of imago; Rhyssa [= Thalessa] lunator destroys the larva.
- 1127. RILEY, C. V. Second annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <5th Ann. Rept. State Board of Agric. for 1869, March, 1870, pp. 136 + 6, 99 figs. Separate: <Jefferson City, Mo., March, 1870, pp. 136+6, 99 figs.

CONTENTS.

Preface	3
NOXIOUS INSECTS.	
PRINCE OF THE CONSTITUTE ON ENTONOLOGY OF THE STATE HOPEL	

5

8

REPORT OF THE COMMITTEE ON ENTOMOLOGY OF THE STATE HORTI-CULTURAL SOCIETY

Noxious insects less injurious in Missouri in 1869 than usual, 5—The army-worm and the grain plant-louse considerably injurious in Missouri in 1869, 5—The chinch-bug and the codling-moth less injurious, 6—A species of Thrips destroying great numbers of the Curculio, 6—Eggs of the apple-tree plant-louse destroyed by insect foes and birds, 6—According to Dr. Hull the "scab" in apples is caused by the apple-tree plant-louse, 7—The pickle-worm doing great damage during 1869, 7—Importance of preventing the introduction of injurious insects, 7—Cultivation causes insects to multiply unduly, 8—More attention paid in Europe to injurious insects than in this country, 8.

1127. RILEY, C. V.—Continued.

flora not so highly improved and developed as in Europe, 12—Australian fauna still more "old-fashioned" than America, 12—The parasites of injurious insects are not imported with the insects themselves, 13—Government aid should be solicited to exterminate recently imported injurious insects, 13—But little attention given so far by our Government to assist the study of economic entomology, 14—Danger of introducing the oyster-shell bark-louse into Missouri, 15—Immunity of the Pacific States from many of our fruit insects, 15.

THE CHINCH-BUG,* Blissus leucopterus

15

37

37

37

It is the most injurious of all insects infesting grain, 16-Its past history, 17; it was known in South Carolina in Revolutionary times, 17; it was injurious in Missouri as early as 1854, 17; noticed in Illinois in 1840, 17; it was very injurious in Missouri in 1868, 17; but hardly noticed in 1869, 17—Probable reason why it was not noticed in Missouri in former times, 18-Why it is not injurious in Massachusette and New York, 18—Its natural history, 18—The pupa state in the different insect orders, 18—Time required for different insects to complete the cycle of development, 19-The chinch-bug is two-brooded in Missouri, 19-Its winter quarters, 20-Its rapid multiplication, 20-Dr. Shimer's account of its nuptial flights, 21-It deposits the eggs underground on the roots of the plant, 21—The egg, 22—Dimorphous forms of the chinch-bug, 22-Its destructive powers, 22-Account of its appearance in immense numbers, 23-Heading off the marching bugs by a barrier of pine boards, 23—Heavy rains destructive to the chinch-bug, 24-Moisture injurious to the egg, 24—The chinch-bug is always worse in a day season than in a wet one, 24-Dr. Shimer's theory on epidemic disease affecting the chinch-bug, 25-Cannibal foes of the chinch-bug, 25; several species of ladybirds, 25; the weeping lacewing, 26; how the lacewing larva seizes its prey, 26; the insidious flower-bug, 27; the common quail, 28-Amount of damage done by the chinchbug, 28-Remedies, 28-Burning in winter the old corn-stalks and other dead stuff on and near the fields, 29-Mixing winter rye among spring wheat, 29-Intercepting the marching bugs by fence-boards, 29—Sowing gas-lime, 30—Other remedies, 30— Bogus chinch-bugs, 31-Several species of Heteroptera confounded with the true chinch-bug, 31-The smell emitted by the half-winged bugs, 32-The insidious flower-bug, 32-The ashgray leaf-bug and its injury to grape-vines, 33-The flea-like negro-bug, 33; injury caused by it to raspberry, strawberry, and garden flowers, 34-Two other species of negro-bug, 35-Recapitulation of the natural history of the chinch-bug, 36.

The tent-caterpillar of the forest, Clisicoampa disetria.....

It can not properly be called an army-worm, 37.

^{*} Extract in < Rept. State Board Agric., Kansas for 1873-74, pp. 129-131.

1127.	RILEY, C. V.—Continued.	
	The Southern grass-worm, Laphygma frugiperda	41
	It resembles in habits the veritable army-worm, 41.	
	The true army-worm	41
	Its past history, 41—Rev. Powers's account of its invasion in the New	
	England States in 1770, 42—Accounts of later invasions previous	
	to 1861, 43—Years of its appearance in Illinois, 43—The invasion	
	of the year 1861, 44—Its appearance in Missburi in 1869, 44—Its	
	sudden appearance and disappearance, 45—Reason for the ap-	
	parently sudden appearance, 45—Army-worm years are wet with	
	the preceding year dry, 46—Reason for the increase and decrease	
	of the number of worms, 46—Its natural history, 47—Previous ac-	
	counts of its natural history, 47—When the eggs are laid, 47—Where they are laid, 48—Misdirected instinct in insects and	
	birds, 48—Exceptions to the normal habit of the worm, 48—Color	
	of the worm, 49—The chrysalis and image, 49—Parasites, 50;	
	the red-tailed Tachina-fly, 50; its beneficial work, 50; it in-	
	fests also other insects, 50; Walsh's description of the fly, 51; it	
	has been redescribed as Exorista osten sackenii, 51; the yellow-	
	tailed Tachina-fly, 51; description of the fly, 51; the glassy Me-	
	sochorus, 52; the diminished Pezomachus, 52; the military	
	Microgaster, 52; the purged Ophion, 53; the army-worm Ich-	
	neumon-fly, 53—Habits of the army-worm and suggestions for	
	its destruction, 53—Burning grass meadows in winter or early	
	spring, 54—Plowing late in the fall, 54—The marching of the	
	worms, 54—Plants they prefer, 54—They become beneficial by	
	devouring the chess in the fields, 55—Ditching, 55—Description	
	of the insect as larva and imago, 56.	
	INSECTS INFESTING THE SWEET-POTATO	56 57
	Tortoise-beetles	57
	The clubbed tortoise-beetle affects the Irish potato, 56—Its general appearance, 57—Characteristics of tortoise-beetles, 57—	
	Merdigerous habits of tortoise-beetles and others of the same	
	family, 58—General appearance of the larvæ, 58—Their dung	
	parasol, 59—Larval molts, 59—Egg of tortoise-beetles, 60—The	
	chrysalis, 60—Habits of and injury done by the beetles, 60—Rem-	
	edies, 61.	
	The two-striped sweet-potato beetle, Cassida bivittata	61
	It seems to be confined to that plant, 61—The larva and the use of	
	its fork, 61—Its pupa and imago, 61.	
	The golden tortoise-beetle, Coptocycla aurichalcea	62
	Food-plants and characteristics of the larva, 62—Brilliant color of	
	the beetle, 62.	-
	The pale-thighed tortoise-beetle, Coptocycla aurichalcea	62
	It is hardly distinguished from the foregoing species, 62.	63
	The mottled tortoise-beetle, Coptocycla guttata	03
	The black-legged tortoise-beetle, Cassida nigripes	63
	Characteristics of the image and larva, 63.	w
	THE PICKLE-WORM, Eudioptis nitidalis	64
	Other insects infesting cucurbitaceous vines	64
	The squash-borer, 64—It seems to be confined to the Eastern States,	
	64—The striped cucumber-beetle, 64—Injury done by the beetle,	
	64; by the larva, 65—The larva and pupa, 65—Number of annual	
	generations, 65—Remedies, 66—Extent of the injury caused by	
	it, 66—The 12-spotted Diabrotica, 66.	

1107	Denne O. V. Continued	
1127.	RILEY, C. V.—Continued.	
	The pickle-worm	67
	Characteristics of the moth, 68—Accounts of injury done by the	
	worm in Missouri and Illinois, 69; in other portions of the coun-	•
	try, 70—It was not known before as injurious, 70—Remedy, 70.	
	INSECTS INJURIOUS TO THE GRAPE-VINE	71
	The hog-caterpillar of the vine, Ampelophaga myron	71
	The egg, 71—Characteristics of the larva, 71—Its habits when	
	about to transform, 72—The chrysalis and imago, 72—It is, one-	
	brooded North and two-brooded farther South, 72—It is very in-	
	jurious, 73—The Microgaster parasite and its development, 73—	
	Habits of caterpillar infested with the parasite, 73.	
`	The achemon Sphinx, Philampelus achemon	74
	Appearance and habits of the larva, 74—The chrysalis, 75—The	
	insect is single-brooded, 75—The moth and its issuing from the	
	pupa-shell, 75—No parasites known, 76.	
	The satellite Sphinx, Philampelus pandorus	76
	How to distinguish its larva from that of the foregoing species, 76	
	-Development of the larva, 76-Variations in color of the larva,	
	77—Its position when at rest, 77—The moth, 78.	
	The Abbot Sphinx, Thyreus abbotii	78
	Its distribution, 78—The larva varies much in color, 78—The chrys-	
	alis and imago, 79. The blue caterpillars of the vine	79
	The eight-spotted forester, Alypia octomaculata, 80—Larva pre-	15
	viously mistaken for it, 80—Habits and characteristics of the	
	larva, 80—Harris's description of the larva, 81—The moth, 81—	
•	Mr. Andrews's account of its ravages, 81—Remedies, 82.	
	The beautiful wood nymph, Eudryas graia, 83—Characteristics of	
	the moth, 83—Close resemblance between the larva of this and	
	the foregoing species, 83—The differences pointed out, 83—De-	
	velopment of the insect, 83.	
	The pearl wood nymph, Eudryas unio, 83—It greatly resembles the	
	beautiful wood nymph, 83—Its probable larva, 84—Practical im-	
	portance of distinguishing these closely allied species, 84.	
•	The American Procris, Harrisina americana	85
	Work of its larva, 85—Description of full-grown larva, 86—The	
	moth, 86—It is not very destructive, 86—Two annual broods of	
	the insects, 86—Parasite of the American Procris, 87.	
	The new grape-root borer, Prionus laticollis	87
	Correction of opinion formerly expressed, 87.	
	The broad-necked Prionus, 87—Duration of the larva state, 87—	
	Its transformation, 88—It bores also in apple roots, 88—Great	
	damage done by the borer, 88—No good remedy known, 88.	
	The tile-horned Prionus, Prionus imbricornis, 89—How it differs from the foregoing, 89—Its occurrence on prairie land, 90—Small	
	dimorphous male form, 90—The larva subsists also upon the	
	roots of herbaceous plants, 90—Practical considerations, 91.	
	The grape-seed maggot, Isosoma vitis	92
	The grape-seed Curculio larva of the first report is that of a hymen-	
	opterous insect, 92-The perfect insect is closely allied to the	
	joint-worm fly, 92-Mr. Saunders' account and description of the	

1127. RILEY, C. V.—Continued.	
THE CANKER-WORM. Paleacrita vernata: Anisopteryx pometaria The eggs, 94—The larva and larval changes, 95—Importance for	94
the orchardists to recognize the true canker-worm, 95—The imported elm leaf-beetle mistaken for it, 95—Description of the	
larva, 96—Its food-plants, 96—The chrysalis, 96—Only one an-	
nual brood in the latitude of St. Louis, 37—The moth and its varieties, 97—It is less injurious in Missouri than in the Eastern	-
States, 97—Remedies, 98—Classification of remedies proposed,	
98—The trough and bandage systems, 99—Muriate of lime as remedy, 100—Jarring the tree, 101—Late fall plowing, 101—	
Summer plowing, 102—Efficiency of hogs, 102—Enemies, 102;	
birds, mites, and parasites, 102; ground-beetles, 103; the fraternal	
potter-wasp, 103. Cabbage-worms	104
The Southern cabbage butterfly, Pieris protodice	104
Its geographical range, 104—Injury caused by it in Missouri, 104 —Description of the larva, 105—The chrysalis and imago, 105— Habits and other food-plants, 105.	
The potherb butterfly, Pieris olearacea, 105—It is a Northern spe-	
cies, 105-It will very likely never occur in Missouri, 106-Geo-	
graphical range of insects principally influenced by temperature,	
106—Isentomic lines, 106—Southern insects found near St. Louis, 106.	
The imported cabbage butterfly, Pieris rapæ, 106—Amount of dam-	
age caused by it in Canada, 107—Its spread westward, 107—It	
will undoubtedly spread to St. Louis, 107—The insect in Europe, 107—History of its introduction, 107—The insect in its dif-	
ferent stages, 108—Its food, 108—Remedies, 109—Parasites, 109.	
The cabbage Plusia, Plusia brassica	110
Characteristics and habits of the larva, 110—Its transformations,	
111—Remedies, 111—Description of larva, chrysalis, and imago, 111—A similar worm occurring on thistles, 112.	
The zebra caterpillar, Mameetra picta	112
Habits and characteristics of the larva, 112—The chrysalis and	
the moth, 113—Two annual broods, 113—Food-plants, 113. THE TARNISHED PLANT-BUG, Lygus pratensis	113
Injury caused by it to various trees and plants, 114—It is a very	110
variable species, 114—Its development, 114—No effective remedy	
known, 115—Preventive measures, 115.	
THE PHILENOR SWALLOW-TAIL, Papilio philenor Its food-plant, 116—Damage done by it, 116—Characteristics and	116
development of the larva, 116—Description of the larva, 117—	
The pupa, 117—The image, 117—Prevention, 118.	
THE COTTONWOOD DAGGER, Aeronycta lepusculina	119
General appearance of the larva, 119—Two annual broods, 119—	•
Chrysalis and moth, 119—Larvæ of other species belonging to the genus Acronycta, 119—Parasites, 120—Description of larva	
and imago, 120—Characters and habits of other species of the same genus, 121.	
THE MISSOURI BEE-KILLER, Proctacanthus milbertii	121
The true scientific name of the Nebraska bee-killer, 121—Wing-veins	
of the genera Asilus, Promachus, and Erax, 122—Description of the Missouri bee-killer, 122—How to destroy the flies, 123—Habits	
and life-history of Asilus-flies, 123—Description of larva and pupa	
of Erax bastardi (?), 124—Synonymical notes on the imago, 124.	

1127. RILEY, C. V.—Continued.

INNOXIOUS INSECTS.

THE GOAT-WEED BUTTERFLY, Paphia glycerium	125
Its geographical distribution and position in classification, 125—Its	
food-plant, 125—Habits of the larva, 126—Larval changes, 126—	
Conformity in the color of the larva with that of the leaves,	
127—Description of the full-grown larva, 127—Transformation	
of the larva to chrysalis, 127—The two sexes of the imago, 127—	
Hibernation, 128.	
THE BLACK BREEZE-FLY, Tabanus atratus	128
Breeze flies beneficial in the larva state, 128-Tormenting power	
of breeze-flies, 128—Their mode of flight, 129—Our knowledge	
of their larval character and habits, 129—General characters of	
the larva of the black breeze-fly, 129-It is semi-aquatic, 129-	
Walsh's description of the larva, 130—Habits and food of the	
larva, 130-Its transformations, 131-Description of the pupa,	
131-Probable habits of breeze-fly larvæ on the Western prai-	
ries, 132.	
GALLS MADE BY MOTHS.	132
THE FALSE INDIGO GALL-MOTH, walshia amorphella	132
The gall and its structure, 132—General appearance of the larva	
and the moth, 133—Description of larva and imago, 133.	
THE MISNAMED GALL-MOTH, Euryptychia saligneana	134
Is it a true gall-maker or an inquiline?, 134—Walsh's description	
of the larva, 134-Description of the image, 134-Generic char-	
acters, 134—Reasons why the insect is an intruder and not a gall-	
maker, 134—Enumeration of the known gall-making moths, 135—	
How the gall is formed, 135.	
RILEY, C. V.) Mr. Walsh's nortrait. < Amer. Ent., March. 1	X7().

1128. [RILEY, C. V.] Mr. Walsh's portrait. <Amer. Ent., March, 1870, v. 2, p. 129.

Remarks accompanying the portrait of B. D. Walsh; resolutions passed on the death of Walsh by the London branch of the Entomological Society of Ontario; by the American Entomological Society, the Illinois State Horticultural Society, and by the Kansas State Horticultural Society.

- 1129. RILEY, C. V. The plum Curculio, Conotrachelus nenuphar Herbst. <Amer. Ent., March, 1870, v. 2, pp. 130-137, fig. 92.
 - Paper read before the Illinois State Horticultural Society at its 14th annual meeting; summary of established facts and discussion of mooted points in the life-history of Constructelus nenuphar; seasons, habits, transformations, food-plants, enemies of and means against the same; descriptions and figures of larva and imago; figure of pupa; hibernation and the effects of climate on the prolongation of the life of insects.
- 1130. RILEY, C. V. Insects injurious to the grape-vine. No. 6. <Amer. Ent., March, 1870, v. 2, pp. 150-153, figs. 100-102. Reprint, with slight changes: <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 80-82, fig. 55.

Treats of Alypia octomaculata; see No. 1127 for synopsis of contents.

- 1131. [RILEY, C. V.] Pithy blackberry gall. <Amer. Ent., March, 1870, v. 2, pp. 159-160, fig. 103.
 - Answer to inquiry of C. W. ["S. C."] Spaulding; descriptions and figures of larva and gall of Diastrophus nebulosus; figure of pupa; seasons, foodplant, guest-fly [Periclistus sylvestris], and parasite [Eurytoma diastrophi] and means against this species; the genus Diastrophus confined to Rosacea, Cynips to Cupulifera and Antistrophus to Composita.
- 1132. [RILEY, C. V.] Clover-worms. < Amer. Ent., March, 1870, v. 2, p. 160.
 - Answer to inquiry of G. Pauls; geographical distribution, food-habits, vernacular names and synonymy of and means against Asopia costalis.
- 1133. [RILEY, C. V.] Seed:ticks under bark of apple-trees. < Amer. Ent., March, 1870, v. 2, p. 160.
 - Answer to inquiry of O. B. Galusha; character of insects as regards the number of legs; occurrence of *lxodes unipunctata* under outer bark of apple-trees at Morris, Ill.
- 1134. [RILEY, C. V.] Parasitic cocoons. < Amer. Ent., March, 1870, v. 2, p. 160.
 - Answer to inquiry of S. W. Beckworth; occurrence of a mass of cocoons of [Microplitis ceratomia var. actuosus] under red-oak at South Pass, Ill.
- 1135. [RILEY, C. V.] Is any knowledge useless? <Amer. Ent. and Bot., April, 1870, v. 2, pp. 164-166.
 - keprint of article from Manufacturer and Builder, November, 1869; minute investigations in science may be of great practical importance; cases in which a knowledge of life-history of Galeruca calmariensis [=zanthomelæna], Constrachelus nenuphar, and Lymexylon navale was or might have been of great value.
- 1136. [RILEY, C. V.] Tomato fruit-worm. <Amer. Ent. and Bot.,
 April, 1870, v. 2, p. 172.

 Notice of statement by J. J. Weir that Heliothis arminera was bred from
 - Notice of statement by J. J. Weir that *Heliothis armigera* was bred from larvæ which fed on the fruit of tomato in England; food-plants of this insect.
- 1137. RILEY, C. V. Insects injurious to the grape-vine. No. 7. < Amer. Ent. and Bot., April, 1870, v. 2, pp. 173-174, figs. 107-108. Reprinted, with slight changes, from <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 85-87, figs. 58-59.
 - Treats of Procris [=Harrisina] americana; see No. 1127 for synopsis of contents.
- 1138. [RILEY, C. V.] The death web of young trout. <Amer. Ent. and Bot., April, 1870, v. 2, p. 174.
 - Reprint, with review, of Seth Green's "An enemy to young trout;" young trout and young white-fish killed by the web of an unknown aquatic larva, probably one of the caddis-flies; habits of the larvæ of *Phryganeidæ*. See No. 1160.
- 1139. [RILEY, C. V.] "Scab" in apple v. apple-tree plant-lice. <Amer. Ent. and Bot., April, 1870, v. 2, p. 178.
 - Notes the freedom of apple-trees in 1870 from the eggs of Aphis mali, and the opportunity to test the question of the connection of these insects with the "scab."

- 1140. [RILEY, C. V.] Insects named. < Amer. Ent. and Bot., April, 1870, v. 2, p. 179.
 - Answer to inquiry of M. Hobart; identification of several insects; foodplants of Grapta comma, Geometra [= Zerene] catenaria, Serica vespertina, and Tetraopes 5-maculatus; geographical distribution of Grapta comma; description of larva of Ecpantheria scribonia.
- 1141. [BILEY, C. V.] Supposed trout enemy. <Amer. Ent. and Bot., April, 1870, v. 2, pp. 179-180.
 - Answer to inquiry of F. Mather; identification of several insects; habite and occurrence of Capnia minima; occurrence of Piophila cases and Ptinus brunness at Honeoye Falls, N. Y.
- 1142. [RILEY, C. V.] Food for trout. < Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of Seth Green and Collins; with knowledge of the character of the worms which forms a desirable food for young trout, it may be possible to suggest some method of propagating the worms artificially.
- 1143. [RILEY, C. V.] Hair-snakes. < Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of E. W. M.; parasitic habits, abodes and prolificacy, and classificatory relations of Gordius varius and G. aquaticus.
- 1144. [RILEY, C. V.] Egg-sack of some unknown spider. <Amer. Ent. and Bot., April, 1870, v. 2, p. 180.

 Answer to inquiry of A. Engelmann; figure of egg-sack of *Epeira* sp. ?
- 1145. [RILEY, C. V.] Do worker bees sting the drones to death?

 <Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of M. W. V.; there is no good reason to doubt that worker bees sting the drones to death when the mission of the latter is ended.
- 1146. [RILEY, C. V.] Red spider. <Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of R. H. Warder; Trombidium [= Tetranychus] telarius imported from Europe; size, color, and abodes of and means against the same.
- 1147. [RILEY, C. V.] Insect named. <Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of M. Barrett; food-habits of Psocus venosus and other Psocidæ.
- 1148. [RILEY, C. V.] To destroy plant-lice. <Amer. Ent. and Bot., April, 1870, v. 2, p. 180.
 - Answer to inquiry of B. F. Lazear; means against Aphididæ on house plants.
- 1149. [RILEY, C. V.] Raspberry root gall. <Amer. Ent. and Bot., April, 1870, v. 2, p. 181, fig. 110.
 - Answer to inquiry of K. Parsons; description and figure of gall of *Rhodites* radicum occurring on roots of *Rosaceæ*; genera of parasites raised from it; interest of the question of the manner and extent of parasitization of this gall.
- 1150. [RILEY, U. V.] Spined slug-worm. <Amer. Ent. and Bot., April, 1870, v. 2, p. 181.
 - Answer to inquiry of L. G. Saffer and A. R. Bodley; brief description of the larva of Limacodes sp.

- 1151. [RILEY, C. V.] Apple-tree insects. < Amer. Ent. and Bot., April, 1870, v. 2, p. 181.
 - Answer to inquiry of L. Camfield; identification of specimens and directions for the destruction of Orgyia leucostigma and Physica nebulo [=Acrobasis indiginella].
- 1152. [RILEY, C. V.] Native apple-tree bark-lice. <Amer. Ent. and Bot., April, 1870, v. 2, p. 181, fig. 111.
 - Answer to inquiry of A. C. Hammond; food-plants of and means against Diaspis harrisii [= Chionaspis furfurus]; figure of infested twig of apple; the presence of enemies and parasites prevent this species becoming a formidable pest.
- 1153. [RILEY, C. V.] The hedge-hog caterpillar. <Amer. Ent. and Bot., April, 1870, v. 2, p. 182, fig. 112.
 - Answer to inquiry of H. Burt; descriptions and figures of larva and image of Arctia [= Pyrrharctia] isabella; figure of pupa and cocoon; habits and hibernation; its larva and that of Ecpantheria scribonia called "feverworm," and ignorantly supposed to cause malarial fevers; food-habits of Horinus [= Merinus] lævis.
- 1154. [RILEY, C. V.] Chick-weed Geometer. < Amer. Ent. and Bot., April, 1870, v. 2, p. 182.
 - Answer to inquiry of J. Huggins; identification of specimens; reference to account of *Hæmatopis grataria*; Cermatia forceps common in houses in the latitude of St. Louis, Mo.; the sulphur remedy for canker-worms absurd.
- 1155. [RILEY, C. V.] Bean-weevil. < Amer. Ent. and Bot., April, 1870, v. 2, p. 182.
 - Answer to inquiry of G. W. Copley; specimens identified; occurrence and ravages of *Bruchus obsoletus* in Illinois.
- 1156. [RILEY, C. V.] Bag-worm at South Pass, Ill. < Amer. Ent. and Bot., April, 1870, v. 2, p. 182.
 - Answer to inquiry of G. H. Baker; occurrence of Thyridopteryx ephemeraformis at South Pass, Ill.; importance of its destruction.
- 1157. [RILEY, C. V.] Eggs of oblong-winged katy-did. <Amer. Ent. and Bot., April, 1870, v. 2, p. 182.
 - Answer to inquiry of E. D. Ladd; differences between the eggs of Phylloptera [= Amblycorypha] oblongifolia and those of Platyphyllum [= Cyrtophyllus] concavus; the former occur on current and various trees.
- 1158. [RILEY, C. V.] Insects injurious to the grape vine. No. 8. <Amer. Ent. and Bot., May, 1870, v. 2, pp. 208-209, fig. 127. Reprint, with slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 61-63, fig. 24.
 - Treats of Desmia maculalis; see No. 1301 for synopsis of contents.
- 1159. [RILEY, C. V.] The periodical Cicada alias the 17-year and 13-year locust. <Amer. Ent. and Bot., May, 1870, p. 211.
 - Quotes, from the 1st Ann. Rept. State Ent. Mo., the localities in which Cicada [= Tibicen] septendecim and C. [= T.] tredecim will appear in 1870, with request for reports of the occurrence of these insects.
- 1160. [RILEY, C. V.] The death-web of young trout. < Amer. Ent. and Bot., May, 1870, v. 2, p. 211.
 - Supplementary to No. 1135; the larvæ mentioned belong to the genus Simulium.

- 1161. [RILEY, C. V.] Worms under mulch hay. <Amer. Ent. and Bot., May, 1870, v. 2, p. 212.
 - Answer to inquiry of J. F. Flagg; description, season, habits, and ravages of and means against the larve of Tipula sp.
- 1162. [RILEY, C. V.] A new pear-tree insect. <Amer. Ent. and Bot., May, 1870, v. 2, p. 212, fig. 129.
 - Answer to inquiry of E. J. Ayres; food-habits and ravages of and means against *Platycerus quercus*; figure of imago.
- 1163. [RILEY, C. V.] Apple-twig borer. < Amer. Ent. and Bot., May, 1870, v. 2, p. 212.
 - Answer to inquiry of J. B. Myers; Bostrichus [= Amphicerus] bicaudatus bores into the axil of limbs of pear-trees.
- 1164. [RILEY, C. V.] Cocoons of polyphemus moth. < Amer. Eut. and Bot., May, 1870, v. 2, 212.
 - Answer to inquiry of H. J. Dunlap; cocoon of Attacus [= Telea] polyphomus found on Morello cherry-tree.
- 1165. [RILEY, C. V.] Galls on supposed dock. <Amer. Ent. and Bot., May, 1870, v. 2, p. 212.
 - Answer to inquiry of S. V. Summers; Gelechia galla-solidaginis forms galls on stems of Solidago; Gastrophysa [= Gastroidea] cyanea breeds on Rumex.
- 1166. [RILEY, C. V.] Mossy rose-gall. <Amer. Ent. and Bot., May, 1870, v. 2, p. 213, fig. 130.
 - Answer to inquiry of W. M. Locke; description of gall, larva, and imago of *Rhodites rosæ*; figure of the gall; description of a parasitic larva.
- 1167. [BILEY, C. V.] Punctures on the rose twig. <Amer. Ent. and Bot., May, 1870, v. 2, p. 213, fig. 131.
 - Answer to inquiry of G. W. Copley; punctures in rose-stem; eggs, probably of a cricket, imbedded in pith of the same; description of the egg and figure of injured stem.
- 1168. [RILEY, C. V.] Snout-beetle. < Amer. Ent. and Bot., May, 1870, v. 2, p. 213.
 - Answer to inquiry of Mary Treat; Hylobius confusus a timber borer and usually in pine.
- 1169. [RILEY, C. V.] The oyster-shell bark-louse in Missouri. < Amer. Ent. and Bot., May, 1870, v. 2, pp. 213-214, fig. 132.
 - Answer to inquiry of B. P. Hanan; occurrence of Aspidiotus conchiformis [=Mytilaspis pomorum] in Missouri; importance of the thorough extirpation of the insect; figure of a piece of bark infested by the same.
- 1170. [BILEY, C. V.] The pod-like willow-gall. <Amer. Ent. and Bot., May, 1870, v. 2, p. 214, fig. 133.
 - Answer to inquiry of J. R. Muhleman; description and figure of the gall of *Cecidomyia salicis-siliqua*; figure of larva; food-plants, synonymy, and description of the pupa of the same.
- 1171. [RILEY, C. V.] Bee-nest. <Amer. Ent. and Bot., 1870, v. 2; May, p. 214, fig. 134; September, p. 307.
 - Answer to inquiry of J. R. Muhleman; description and figure of larva of Prosopis affinis in hot, low current-stem; probability that Coratina dupla breeds twice a year.

- 1172. [RILEY, C. V.] Beetles named. < Amer. Ent. and Bot., May, 1870, v. 2, p. 214.
 - Answer to inquiry of S. V. Summers; specimens identified; difference between Canthon lævis and C. chalcites.
- 1173. [RILEY, C. V.] Great discovery: Curculio extermination possible! <Amer. Ent. and Bot., June, 1870, v. 2, pp. 225-227.

 Notice: <Cultivator and Country Gentleman, 9 June, 1870, v. 35, p. 361.
 - Reprint of articles by J. E. Chamberlain and Mrs. H. Wier on the destruction of the Curculio; letter from W. B. Ransom; criticism of the same; means against Conotrachelus nenuphar; distinctions between C. nenuphar and Anthonomus quadrigibbus.
- 1174. [RILEY, C. V.] The death web of young trout. < Amer. Ent. and Bot., June, 1870, v. 2, pp. 227-228, figs. 143-144.
 - Value and progress of fish-culture; habits of Simulium piscicidium, with the observations of Seth Green and Sara J. McBride; figures larva and pupa of S. piscicidium and the image of S. molestum.
- 1175. [RILEY, C. V.] Insects injurious to the grape-vine. No. 9. <Amer. Ent. and Bot., June, 1870, v. 2, pp. 234-235, fig. 148. Reprint, with slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 65-68, fig. 27.
 - Treats of Pterophorus [= Oxyptilus] periscelidactylus; see No. 1301 for synopsis of contents
- 1176. [RILEY, C. V.] The apple Curculio. <Amer. Ent. and Bot., June, 1870, v. 2, p. 243, fig. 152.

 Figures of the image of Anthonomus quadrigibbus.
- 1177. [RILEY, C. V.] The new Curculio remedy. <Amer. Ent. and Bot., June, 1870, v. 2, p. 243.</p>
 Results of experiments in the use of Ransom chip-trap for Constructelus nenuphar.
- 1178. [RILEY, C. V.] Tarantula of Texas. <Amer. Ent. and Bot., June, 1870, v. 2, p. 244.

 Answer to inquiry of L. J. Stroop; the figure of Mygale hentzii given in volume one is somewhat incorrect.
- 1179. [RILEY, C. V.] Ailanthus silk-worm naturalized. <Amer. Ent. and Bot., June, 1870, v. 2, p. 244.

 Answer to inquiry of A. S. Fuller; introduction and naturalization of Atta-
 - Answer to inquiry of A. S. Fuller; introduction and naturalization of Attacus cynthia.
- 1180. [RILEY, C. V.] Cypress gall. <Amer. Ent. and Bot., June, 1870, v. 2, p. 244, fig. 153.
 - Answer to inquiry of J. P. Stelle; description and figures of gall and description of gall and image of *Cecidomyia cupressi-ananassa* n. sp. on cypress; figures breast-bone of the larva.
- 1181. [RILEY, C. V.] Tent-caterpillar of the forest. < Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to the inquiries of A. M. Brown and J. H. Evans; habits, food-plants, and ravages of Clisiocampa sylvatica [=disstria] and C. americana.

- 1182. [RILEY, C. V.] Worm-boring into peach. < Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to inquiries of W. C. Flagg, A. C. Hammond, and M. M. Hooton; food-plants and description of larva of Xylina cinerea [=Lithophane antennata].
- 1183. [RILEY, C. V.] Insects named. Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to inquiry of A. Engelman; descriptions of Smilia auriculata and Membracis ampelopsidis; both species common on grape-vines.
- 1184. [RILEY, C. V.] Lice on "snow-balls. <Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to inquiry of Mrs. C. L. Seymour; means against Aphidida.
- 1185. [RILEY, C. V.] Twig-borer. <Amer. Ent. and Bot., June, 1870, v. 2 p. 245, fig. 154.
 - Answer to inquiries of S. H. Kriedelbaugh and G. F. Merriam; figures of male and female Bostrichus [=Amphicorus] bicaudatus, which bore into the axils of grape-buds.
- 1186. [RILEY, C. V.] Bee enemy. < Amer. Ent. and Bot., June, 1870, v. 2, p. 245.
 - Answer to inquiry of F. Brewer; Staphylinus maculosus, found eating a bee, is rather a scavenger than an insect of prey.
- 1187. [RILEY, C. V.] Knots on apple-tree roots caused by root-lice. <Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of B. N. McKinstry; ravages of and means against Schizoneura lanigera on roots of young apple-trees.
- 1188. [RILEY, C. V.] Beetles named. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of W. W. Daniells; Ithycerus noveboracensis injures appleleaves.
- 1189. [RILEY, C. V.] Bag-worm. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of M. M. Hooten; occurrence of young larva of Thyridopteryz ephemeræformus on peach-trees; manner in which they carry their cases.
- 1190. [RILEY, C. V.] The larder-beetle. <Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of S. H. Kriedelbaugh; food-habits and description of the larva and image of Dermestes lardarius.
- 1191. [RILEY, C. V.] Water-bug. <Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of W. H. Harrington; habits and description of Ranatra fusca.
- 1192. [RILEY, C. V.] Gregarious worms on horse-chestnut. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of W. R. Howard; oviposition, food-plant, and parasite of *Tortrix* [= Cacoccia] rileyana.
- 1193. [RILEY, C. V.] Pupa of the disippus butterfly. <Amer. Ent. and Bot., June, 1870, v. 2, p. 246, fig. 155.
 - Answer to inquiry of T. Montgomery; figure and colors of pupa of Limenitis disippus; food-plants and hibernations of larva.

- 1194. [RILEY, G. V.] Prickly-rose gall. <Amer. Ent. and Bot., June, 1870, v. 2, p. 246.
 - Answer to inquiry of J. Cochrane and J. P. Stelle; brief description of the gall of *Bhodites bioolor* on wild rose.
- 1195. [BILEY, C. V.] Insects feeding on sap of black-walnut. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246. Answer to inquiry of M. Barrett; food-habits of Psocus venosus.
- 1196. [RILEY, C. V.] Locust-borer, <Amer. Ent. and Bot., June, 1870, v. 2, p. 246.

 Answer to inquiry of "Arbor;" means against Arhopalus [= Cyllene] robinia.
- 1197. [RILEY, C. V.] To exterminate cockroaches. < Amer. Ent. and Bot., June, 1870, v. 2, p. 246.

 Answer to inquiry of R. F. Weitbree; means against cockroaches.
- 1198. [RILEY, C. V.] The white-lined morning Sphinx. (Deilepkila lineata, Fabr.) < Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 257-258, figs. 162-164. Reprint, with additions and slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 140-142, figs. 60-62.
 - Treats of Deilephila lineata; see No. 1301 for synopsis of contents.
- 1199. [RILEY, C. V.] Descriptive entomology. <Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 258-261. Abstract: <Cultivator and Country Gentleman, 6 April, 1871, v. 36, p. 218.
 - Critical review of a remark by J A. Lintner; calculation of the cost, labor, and extent of a work containing the description and figure of every existing species of insect.
- 1200. [RILEY, O. V.] The tent-caterpillar of the forest. (Clisiocampa sylvatica, Harr.) < Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 261-266, figs. 165-168. Reprint, with additions and slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 121-129, figs. 52-54.
 - Treats of Clisiocampa sylvatica [== disstria]; see No. 1301 for synopsis of contents.
- 1201. [RILEY, C. V.] The Ransom Curculio remedy. <Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 268-271.

 Discussion of the efficacy of the Ransom chip-trap against Constructelus nenuphar; extracts from and criticism of articles by E. S. Hull, W. B. Ransom, and others on this subject.
- 1202. [RILEY, C. V.] Insects injurious to the grape-vine. No. 10. <Amer. Ent. and Bot., July-August, 1870, v. 2, pp. 272-273, fig. 170. Reprint, with additions and slight changes. <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 68-69, fig. 28. Treats of Spilosoma virginica; see No. 1301 for synopsis of contents.
- 1203. [RILEY, C. V.] The Walsh entomological cabinet. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.</p>
 Remarks on the purchase and disposal of the entomological collection of B. D. Walsh.

- 1204. [BILEY, C. V.] The current-worm! < Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.
 - Comments upon the confusing way in which some horticultural publications treat of the currant-worm, without discrimination of species.
- 1295. [RILEY, C. V.] Water larva. < Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.

 Answer to inquiry of F. Mather; habits of the larvæ of Ephemeridæ.
- 1206. [RILEY, C. V.] Large black potato-beetles. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.

Answer to inquiry of R. S. Elliott; Epicauta corvina? injuring potato-vines in Kansas.

- 1207. [RILEY, C. V.] Destroying cherry plant-lice. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.

 Answer to inquiry of G. C. Brackett; means against Myzus cerasi.
- 1208. [RILEY, C. V.] Caterpillars on grape-vines. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.

 Answer to inquiry of G. A. Watson; larvæ of Alypia octomaculata and Acronycta oblinita found on grape-vines; food-plants of the latter species.
- 1209. [RILEY, C. V.] Ash-gray blister-beetle. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 275.

 Answer to inquiry of P. H. Foster; Lytta cinerea [= Macrobasis unicolor] feeding on the three-thorned Acacia.
- 1210. [RILEY, C. V.] Specimens lost. <Amer. Ent. and Bot., July—August, 1870, v. 2, p. 276.

Answer to inquiry of C. H. Roberts; food-habits and description of larva of Gortyna nitela and of an undetermined moth on peach-trees.

1211. [RILEY, C. V.] White willow worm. < Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.

Answer to inquiry of S. H. K.; food-plants and description of larva of and

means against Nematus ventralis.

- 1212. [RILEY, C. V.] Bark-lice on grape-vine and raspberry saw-fly. Amer.Ent. and Bot., July-August, 1870, v. 2, p. 276.

 Answer to inquiry of S. Thompson; Lecanium [= Pulvinaria] vitis common on grape vines in Furgree and North America; its ovincettics and common on grape vines in Furgree and North America; its ovincettics and common on grape.
 - grape-vines in Europe and North America; its oviposition and occurrence in Illinois; description and means against the larva of Sclandria [= Monophadnus] rubi.
- 1213. [RILEY, C. V.] Apple-tree borer; variations in the two-striped Saperda. Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Answer to inquiry of D. B. Wier; colorational variations in Saperda bivittata [=candida]; abundance and ravages of Capsus oblineatus [=Lygus pratensis].
- 1214. [RILEY, C. V.] The plum Curculio breeds in apple. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.

 Answer inquiry of E. Leming; Constructedus nemephor breeds in the fruit of apple.

- 1215. [RILEY, C. V.] Cecropia worm. August, 1870, v. 2, p. 276.
 - Answer to inquiry of J. F. Thompson; occurrence and ravages of the larva of Attacus cocropia on apple-trees.
- 1216. [RILEY, C. V.] Gigantic rhinoceros beetle. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Answer to inquiry of L. G. Saffer; variations in coloration of Dynastes tityus.
- 1217. [RILEY, C. V.] Roman-nosed pupa. <Amer. Ent. and Bot., July-August, 1870, v. 2, p. 276.
 - Answer to inquiry of E. D. Van Winkle; food-habits of Limenitis ursula and L. disippus; the pupe of the two species are alike.
- 1218. [RILEY, C. V.] The onward march of the Colorado potato-beetle. A word to our Canadian neighbors. <Amer. Ent. and Bot., September, 1870, v. 2, pp. 289-291, fig. 181.
 - Remarks on the spread of *Doryphora decemlineata* into Ontario, and means of checking it; efficacy and proper use of Paris green; other remedies; *Lebia grandis* a natural enemy of the larvæ.
- 1219. [RILEY, C. V.] The tarnished plant-bug. (Capsus oblineatus, Say). <Amer. Ent. and Bot., September, 1870, v. 2, pp. 291-293, fig. 182. Reprinted, with additions and slight changes, from <2d Ann. Rept. State Ent. Mo., March, 1870, pp. 113-115, fig. 83.
 - Treats of Capsus oblineatus [=Lygus pratensis]; see No. 1127 for synopsis of contents.
- 1220. [RILEY, C. V.] Osage orange for the mulberry silk-worm.

 <Amer. Ent. and Bot., September, 1870, v. 2, p. 293.

 Reprint and critical review of S. Cornaby's article on the above; important
 - Reprint and critical review of S. Cornaby's article on the above; important articles should not be published anonymously.
- 1221. [RILEY, C. V.] Insects injurious to the grape-vine. No. 11. <Amer. Ent. and Bot., September, 1870, v. 2, p. 295, fig. 185. Reprint with slight changes. <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 77-79, fig. 34.
 - Treats of Pelidnota punctata; see No. 1301 for synopsis of contents.
- 1222. [RILEY, C. V.] The slug on pear and cherry trees. <Amer. Ent. and Bot., September, 1870, v. 2, p. 296.
 - Reprint and critical review of "Addi" on the above; ravages of and means against Sciandria [= Eriocampa] cerasi.
- 1223. [RILEY, C. V.] Appendix to joint-worm article published in vol. 1, No. 8 < Amer. Ent. and Bot., September, 1870, v. 2, pp. 296–297.
 - Introduction to and explanation of the preparation and publication of Walsh's Eurytomides. See No. 384.
- 1224. [RILEY, C. V.] Entomology indeed run mad! < Amer. Ent. and Bot., September, 1870, v. 2, p. 305.
 - Critical review of Mark Miller's article on the current-worm; Nomatus vontricosus [= ribesii] confounded with Euftchia ribearia.
- 1225. [RILEY, C. V.] Red spider. < Amer. Ent. and Bot., September, 1870, v. 2, p. 305.
 - Ravages and transformations of Tetranychus telarius,

- 1226. [RILEY, C. V.] Insects named. < Amer. Ent. and Bot., September, 1870, v. 2, p. 306.
 - Answer to inquiry of J. K. Kidd; food-habits of Silpha peltata [= americana] and allied forms, of Calosoma scrutator and of Laphria [= Dasyllis] thoracica; mimicry between Laphria and Bombus and its use.
- 1227. [RILEY, C. V.] Caterpillar of white-marked tussock moth. Amer. Ent. and Bot., September, 1870, v. 2, p. 306, fig. 186.
 Answer to inquiry of G. C. Brackett; figure of larva and description of image of Orgyia leucostigma; posture of male in repose; habits of female; Saperda bivittata [= candida] usuall; perishes if it has not changed to a pupa before the death of the tree; Uhrysobothris femorata lives for weeks on dead wood.
- 1228. [RILEY, C. V.] Does the apple Curculio go underground to transform? <Amer. Ent. and Bot., September, 1870, v. 2, p. 306.
 - Answer to inquiry of W. Muir; Anthonomus quadrigibbus transforms within the fruit; it does not attack stone fruit.
- 1229. [RILEY, C. V.] Walnut caterpillars. < Amer. Ent. and Bot., September, 1870, v. 2, p. 306.
 - Answer to inquiry of G. M. Levette; seasons, habits, and food-plant of and means against Datana ministra; description of larva.
- 1230. [RILEY, C. V.] Striped blister-beetle. <Amer. Ent. and Bot., September, 1870, v. 2, p. 306, fig. 187.</p>
 Answer to inquiry of A. Galt; means against Epicauta vittata and other Me

loide on potato vines; figure of Epicauta vittata.

- 1231. [RILEY, C. V.] Grape-vine Fidia. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307, fig. 188.

 Answer to inquiry of J. Hetzel; habits, ravages, and food-plants of and
- means against Fidia viticida; figure of this species.

 1232. [RILEY, C. V.] Some interesting insects. < Amer. Ent. and Bot., September, 1870, v. 2, p. 307.
 - Answer to inquiry of A. S. Fuller; ravages of Bruckus obsoletus in Lima beans; of Butalis cerealella in flint corn in stems and of Lobesia [=Eudemis] botrana in blossoms of blackberry; description of the larva case of Physita nebulo [=Acrobasis indiginalla] and of galls of Cecidomyia tubicola on Carya.
- 1233. [RILEY, C. V.] The green hag-moth. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307.
 - Answer to inquiry of S. B. Shaw; food-plants, and synonymy of Callochlora viridis [= Parasa chloris]; description of larva and imago; the larva described by Reakirt does not belong to this species.
- 1234. [RILEY, C. V.] The antiopa butterfly. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307.
 - Answer to inquiry of A. S. Moss; description of larva and image of Vanessa antiopa; the larva feeds on willow; vernacular name and comparative abundance.
 - 1235. [RILEY, C. V.] Rose-gall and pupa of archippus butterfly. <Amer. Ent. and Bot., September, 1870, v. 2, p. 307, fig. 189.
 - Answer to inquiry of L. B. Custar; figure of pupa of Danais archippus; descriptions of three undetermined galls on rose-leaf formed by Rhodites sp.?

 10 ENT

- 1236. [RILEY, C. V.] White grubs in strawberry beds. < Amer. Ent. and Bot., September, 1870, v. 2, p. 307.
 - Answer to inquiry of J. B. Miller; food-habits of larva and description of imago of Cyclocephala immaculata.
- 1237. [RILEY, C. V.] Larva of the thoas swallow-tail < Amer. Ent. and Bot., September, 1870, v. 2, p. 308.
 - Answer to inquiry of E. H. Sprague; food-plant and description of larva of Papilio thous [=cresphontes]; the larva rare in Missouri.
- 1238. [RILEY, C. V.] Larva of clubbed tortoise-beetle. '<Amer. Ent. and Bot., September, 1870, v. 2, p. 308.
 - Answer to inquiry of A. R. Bodley; Cassida, Coptocycla and Deloyala feed on Convolvulaceæ with the exception of Deloyala [=Coptocycla] clarata which feeds on Solanaceæ.
- 1239. [RILEY, C. V.] The banded Ips in calyx of pear. <Amer. Ent. and Bot., September, 1870, v. 2, p. 308.</p>
 Answer to inquiry of G. C. Brackett; food-habits and description of Ips fasciatus.
- 1240. [RILEY, C. V.] The larder-beetle. < Amer. Ent. and Bot., September, 1870, v. 2, p. 308, fig. 191.
 Answer to inquiry of F. S. Sleeper; figures of larva and image and of mag
 - nified hair of larva of Dermestes lardarius; ravages in collections of pre-
- 1241. [RILEY, C. V.] Moth named. Amer. Ent. and Bot., September, 1870", v. 2, p. 308.
 - Answer to inquiry of E. M. Hale; description of Ctenucha latreillana [=rir-ginica]; its abundance in 1870 in the vicinity of St. Louis, Mo.
- 1242. [RILEY, C. V.] The little Cicada. <Amer. Ent. and Bot., September, 1870, v. 2, p. 308.

 Answer to inquiry of G. O. Hardeman: occurrence of Cicada [=Melamagalta]
 - Answer to inquiry of G. O. Hardeman; occurrence of Cicada [=Melampsalta] parvula in Missouri.
- 1243. [RILEY, C. V.] The brown mantispian. <Amer. Ent. and Bot., September, 1870, v. 2, p. 308.</p>
 Answer to inquiry of G. C. Bracket; Mantispa brunnes common, predaceous,
- 1244. [RILEY, C. V.] Small reddish snout-beetle on apple. <Amer. Ent. and Bot., September, 1870, v. 2, p. 30.

and beneficial.

- Answer to inquiry of J. Weed; foed-habits and description of Anthonomus oratægi.
- 1245. [RILEY, C. V.] Prickly rose-gall. < Amer. Ent. and Bot., September, 1870, v. 2, p. 309, fig. 192.
 - Answer to inquiry of "Subscriber;" description and figure of galls and description of Rhodites bicolor.
- 1246. [RILEY, C. V.] Questions answered. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiries of K. Parsons; formation and use of portable cases by Tineina; habits and character of the young of Mytilaspis pomicorticis [=pomorum]; food-habits of ants.

- 1247. [RILEY, C. V.] Hog-caterpillar of the vine infested with parasites. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of J. M. Wilson; occurrence of larva of Darapsa [=Ampelophaga] myron parasitized by Apanteles congregatus.
- 1248. [RILEY, C. V.] Larva of Abbot Sphinx. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of S. E. Todd; occurrence of larva of Thyreus abbotii on grape-vine; means against the same.
- 1249. [RILEY, C. V.] Cecropia worm. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of E. G. Hofman; occurrence of Attacus cecropia on plum.
- 1250. [BILEY, C. V.] Flat-headed borer in soft maples. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of L. R. Elliott; ravages of and means against Chryso-bothris femorata on soft maple; ravages of Arhopalus [= Cyllene] robinia on black-locust.
- 1251. [RILEY, C. V.] Cherry plant-lice and their foes. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309, fig. 193.
 - Answer to inquiry of C. H. Roberts; occurrence on cherry-trees of Aphis [= Mysus] cerasi, and of its enemies, the larva of Hippodamia convergens and Syrphus sp.; figure of the larva, pupa, and image of the former; positions in which both species transform.
- 1252. [RILEY, C. V.] Grape vine flea beetle. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of C. H. Roberts; ravages of Haltion chalybea; means against this species and Sciandria vitis [= Blennocampa pygmæa] on grape-vines.
- 1253. [RILEY, C. V.] Blood sucker and pear-slug. <Amer. Ent. and Bot., September, 1870, v. 2, p. 309.
 - Answer to inquiry of G. A. Watson; food-habits of *Pirates* [= *Melanolestes*] picipes and of Hemiptera in general; means against *Selandria* [= *Erio-oampa*] cerasi.
- 1254. [RILEY, C. V.] The codling-moth. (Carpocapsa pomonella, Linnaeus.) < Amer. Ent. and Bot., October, 1870, v. 2, pp. 321-322. Reprint: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 101-104.
 - Treats of Carpocapsa pomonella; see No 1301 for synopsis of contents.
- 1255. [RILEY, C. V.] Insects injurious to the grape-vine. No. 12. <Amer. Ent. and Bot., October, 1870, v. 2, pp. 327-328, figs. 204-205. Reprint, with slight changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 79-81, figs. 35, 36; Cultivator and Country Gentleman, 1 June, 1871, v. 36, p. 343.
 - Treats of Graptodera [= Haltica] chalybea; see No. 1301 for synopsis of contents.
- 1256. [RILEY, C. V.] The fall army-worm. <Amer. Ent. and Bot., October, 1870, v. 2, pp. 328-329, figs. 206-207.
 - Seasons, food-plants, and ravages of Laphygma frugiperda; figures larvæ of Leucania unipuncta and Laphygma frugiperda; comparison between the latter and Heliothis armigera.

- 1257. [RILEY, C. V.] The rape butterfly; our new cabbage pest. <Amer. Ent. and Bot., October, 1870, v. 2, p. 338. Spread of and means against *Pieris rape*.
- 1258. [RILEY, C. V.] Paris-green for the Curculio. <Amer. Ent. and Bot., October, 1870, v. 2, p. 338.

 Inefficacy of the use of Paris-green against Conotrackelus nonuphar.
- 1259. [RILEY, C. V.] Beetles working in wheat, oats, and rye; the grain Silvanus. <Amer. Ent. and Bot., October, 1870, v. 2, p. 339, fig. 208.
 - Answer to inquiry of M. H. Boye; figure, description, and ravages of Silvanus surinamensis; means against the same and against Calandra graneria in grain; habitat and probable origin of the Silvanus.
- 1260. [RILEY, C. V.] Beetles in dried English currants. <Amer. Ent. and Bot., October, 1870, v. 2, p. 339.

Answer to inquiry of T. V. Munson; Silvanus surinamensis breeding abundantly in dried English currents.

- 1261. [RILEY, C. V.] The same in flouring mills. <Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of S. Blanchard; abundance of Silvanus surinamensis in flouring-mills.
- 1262. [RILEY, C. V.] Carolina Sphinx. < Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of W. R. Howard; the larva of Macrosila [= Protoparce] carolina feeds on tobacco.
- 1263. [RILEY, C. V.] Insects clustered on apple-trees. <Amer. Ent. and Bot., October, 1870, v. 2, p. 339.

 Answer to inquiry of R. L. Ham; habits and food of Psocus renesses.
- 1264. [RILEY, C. V.] Larvæ named. < Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of T. W. Gordon; stinging powers of the larvæ of Empretia stimulea and Hyperchiria varia [= io]; Microgaster sp. parasitic upon Macrosila quinquemaculata [= Protoparce celeus].
- 1265. [RILEY, C. V.] Mite-gall on sugar-maple. <Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of A. Furnas; description of gall of Acarus aceris-crumena n. sp. on leaves of sugar-maple; similar mite-galls occur on plum and cherry.
- 1266. [RILEY, C. V.] Cheese-fly and blow-fly. <Amer. Ent. and Bot., October, 1870, v. 2, p. 339.
 - Answer to inquiry of B.; skippers in cheese are larvæ of Piophila casei; those found in bacon of Calliphora vomitoria.
- 1267. [RILEY, C. V.] Fall army-worm. < Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiries of K. Kelsey and G. Pauls; seasons, ravages, and food-plants of Laphygma frugiperda; seasons and food-plants of Leucania uni-puncta.

- 1268. [RILEY, C. V.] Larva of imperial moth; Thoas swallow-tail. <Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of G. M. Dodge; larva of Ceratocampa [= Eacles] imperiate feeds on maple.
- 1269. [BILEY, C. V.] Large Asilus fly. < Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of L. G. Saffer; food-habits of Promachus vertebratus, P. [= Erax] bastardii, and Asilus missouriensis [= Proctacanthus milbertii]; occurrence of undetermined galls under white-oak trees.
- 1270. [RILEY, C. V.] Mole cricket. <Amer. Ent. and Bot., October, 1870, v. 2, p. 340.

 Answer to inquiry of V. K. Deyo; habits of Gryllotalpa borealis.
- 1271. [RILEY, C. V.] A rare capture in Illinois. <Amer. Eut. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of H. S. Bontell; geographical distribution of Callidryas philea; occurrence of Thysania zenobia in Iowa.
- 1272. [RILEY, C. V.] Hag-moth larva. < Amer. Ent. and Bot., October, 1870, v. 2, p. 340, fig. 209.
 - Answer to inquiries of D. M. Hunter and G. Pauls; figure of larva and description of imago of Limacodes [= Phobetrum] pithecium; larva found on apple-tree; formation of its cocoon.
- 1273. [RILEY, C. V.] Insects named. < Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of Mrs. E. U. B.; larva of *Papilio asterias* feeds on paranip; that of *Alaria* [= *Rhodophora*] *florida* on evening primrose; habits of the imago of the latter.
- 1274. [RILEY, C. V.] Some friends and foes. <Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of C. W. Spaulding; food-habits of Harpactor [= Milyas] cinctus, Mysia [=Anatis] 15-punctata, and Calosoma calidum; Prionus imbricornis bred from grape-vine roots.
- 1275. [RILEY, C. V.] The royal-horned caterpillar. <Amer. Ent. and Bot., October, 1870, v. 2, p. 340.
 - Answer to inquiry of J. T. Hodgen; larva of Ceratocampa [=Citheronia] regalis found on persimmon.
- 1276. [RILEY, C. V.] Caterpillars named. <Amer. Ent. and Bot., October, 1870, v. 2, p. 341, fig. 210.
 - Answer to inquiry of E. H. King; food plants and descriptions of larvæ of Papilio asterias and of Acronycta oblinita; function of the osmaterium of the former; vernacular name and figure of larva, cocoon, and image of the latter.
- 1277. [RILEY, C. V.] The Abbot Sphinx; parasites on its larva. <Amer. Ent. and Bot., October, 1870, v. 2, p. 341.
 - Answer to inquiry of T. W. Gordon; description of Thyreus abbotii; life-history of Microgaster sp. parasitic on larvæ of Sphingidæ.
- 1278. [RILEY, C. V.] Crane-flies; rose-bugs; ants. < Amer. Ent. and Bot., October, 1870, v. 2, p. 341.
 - Answer to inquiry of J. W. Potts; food-habits of Tipula sp. and of Macro-dactylus subspinosus; structure and habits of the several sexes of Formicida.

- 1279. [RILEY, C. V.] Cabbage-worms. < Amer..Ent. and Bot., Oetober, 1870, v. 2, p. 341.
 - Answer to inquiry of B. H. Foster; ravages of and means against Pieris rapa.
- 1280. [RILEY, C. V.] The unicorn prominent. <Amer. Ent. and Bot., October, 1870, v. 2, p. 341.
 - Answer to inquiry of E. Payne; description of larva and image of Notodonta [= Cœlodæys] unicornis; food-plants and mimicry of the larva.
- 1281. [RILEY, C. V.] Insects injurious to the grape-vine. No. 13. <Amer. Ent. and Bot., December, 1870, v. 2, pp. 353-359, figs. 218-219. Reprint with changes: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 84-96, figs. 37-40. See: <Bull. de la Soc. Cent. d'Agric., 1870.
 - Treats of Phylloxera vastatrix; see No. 1301 for synopsis of contents.
- 1282. [RILEY, C. V.] The fall army-worm. < Amer. Ent. and Bot., December, 1870, v. 2, pp. 363-365, figs. 221-223.

 Descriptions of egg, larva, pupa, and image of Prodenia autumnalis n. sp. [= Laphygma frugiperda]; variation of the images; seasons, habits, and ravages of and means against the same; figures of larva and image of Prodenia commelinæ and of image of Leucania unipuncta.
- 1283. [RILEY, C.V.] The so-called web-worm of young trout. < Amer. Ent. and Bot., December; 1870, v. 2, pp. 366-367. Editorial remarks appended to S. J. McBride's communication; description of Simulium piscicidium n. sp.
- 1284. [RILEY, C. V.] Hybrid between a grape-vine and a hickory. Amer. Ent. and Bot., December, 1870, v. 2, p. 373.

 Correction of error in mistaking a gall of Cecidomyia vitis-pomum for a hybrid fruit.
- 1285. [BILEY, C. V.] Death of noted entomologists. <Amer. Ent. and Bot., December, 1870, v. 2, p. 373.

 Notice of death of Julius Lederer and J. T. Lacordaire.
- 1286. [RILEY, C. V.] Osage orange for the mulberry silk-worm. <Amer. Ent. and Bot., December, 1870, v. 2, p. 373.

 Explanation of diff rences in experience in feeding silk-worms on osage orange.
- 1287. [RILEY, C. V.] Insects named. < Amer. Ent. and Bot., December, 1870, v. 2, p. 373.
 - Answer to inquiry of Mrs. M. Chappellsmith; Astoma [= Trombidium] locustarum paraeitic on grasshoppers; similar mites on other insects; food-habits of Nemobius vittatus and Orocharis saltator.
- 1288. [RILEY, C. V.] Locust-borer. < Amer. Ent. and Bot., December, 1870, v. 2, p. 373.
 - Answer to inquiry of W. R. Howard; seasons and method of oviposition of Arkopalus [== Cyllens] robiniæ in black-locust; description of eggs and imago of the same.
- 1289. [RILEY, C. V.] The northern lady-bird; its larvæ. < Amer. Ent. and Bot., December, 1870, v. 2, p. 373.
 - Answer to inquiry of C. E. Billings; food-habits and description of larva of Epilachna borealis.

- 1290. [RILEY, C. V.] Not eggs, but parasitic cocoons. <Amer. Eut. and Bot., December, 1870, v. 2, p. 373.
 - Answer to inquiry of R. Couch; occurrence and true nature of cocoons of Microgaster sp. on larva of Darapsa [= Ampelophaga] myron.
- 1291. [RILEY, C. V.] The cabbage Plutella. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of C. E. Bessey; ravages of *Plutella cruciferarum* on cabbages.
- 1292. [RILEY, C. V.] Gigantic rhinoceros beetle. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374, fig. 224.
 - Answer to inquiry of "Subscriber;" figure of male and description of the male and female of Dynastes tityus; distribution; food-habits of larva.
- 1293. [RILEY, C. V.] Bee-bread devoured by worms. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 - Answer to inquiry of L. C. Francis; food-habits of Ephestia zew [=interpunctella] and Galleria cereana; ravages of the former in old beehives.
- 1294. [RILEY, C. V.] Rape butterfly. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.

 Answer to inquiry of J. E. Cowden; ravages of *Pieris rapæ* on cabbages.
- 1295. [RILEY, C. V.] Grape-leaf gall. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374.</p>
 Answer to inquiry of H. C. Beardslee; occurrence of *Phylloxera vastatrix* at Painesville, Ohio.
- 1296. [RILEY, C. V.] Bean weevil. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.

 Answer to inquiry of H. Kleinhaus; ravages and distribution of Bruchus obsoletus.
- 1297. [RILEY, C. V.] Cabbage insects. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.

 Answer to inquiry of W. R. Howard; ravages, distribution, and synonyms of Plutella oruniferarum; first occurrence of Strackia [= Murgantia] histri-
- 1298. [RILEY, C. V.] Two-striped walking stick. < Amer. Ent. and Bot., December, 1870, v. 2, p. 374.

 Answer to inquiry of C. R. Edwards; occurrence of Spectrum bivittatum in

onica in Missouri.

- Answer to inquiry of C. R. Edwards; occurrence of Spectrum bivitiatum in Kentucky.

 1209 [Ruley C. V.] Ladder spider < Amer. Ent. and Rot. Decem.
- 1299. [RILEY, C. V.] Ladder spider. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.
 Answer to inquiry of L. G. Saffer; explanation of the vernacular name of Epeira [=Argiope] riparia.
- 1300. [RILEY, C. V.] Sugaring for moths; preserving larvæ. <Amer. Ent. and Bot., December, 1870, v. 2, p. 374.

 Answer to inquiry of H. S. Boutwell; occurrence of Papilio thous [=cres-
 - Answer to inquiry of H. S. Boutwell; occurrence of *Papilio thoas* [=cresphontes] in Illinois; directions for the preparation of a fluid for preserving caterpillars.

1301. RILEY, C. V. Third annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <6th Ann. Rept. State Board of Agric. for 1870, April, 1871, pp. 176+7, 73 figs. Separate: Jefferson City, Mo., April, 1871, pp. 176+7, 73 figs.

CONTENTS.

3

5

	PREFACE
NOXIOUS INSECTS.	
tophora	The whole vas food for ferent par Vast exte plicity of guished b family, 9-larve, 10-
Inculio, Conotrachelus nonephar	The common y It is single hibernate the beetle in other Habits of chip-trap successful ber of Cur character more expe specimens the Curcu catcher de catcher, 2 machine, 2 22—Hoote operation, Sigalphus donbt abo and obser 25; its de 25; it atts ural histo the imago, zon Curcu description parasites o tion of pa
nthonomus quadrigibbus	The apple Cur

How it differs from the plum Curculio, 30—Its natural history, 30— Food plants, 30-The punctures it drills into the fruit, 31-The egg, 31-The larva and its habits, 31-The larva transforms within the fruit, 32-Amount of damage it does, 33-How its

1301. RILEY, C. V.—Continued.

work in the fruit can be distinguished from that of other insects, 33—It is very injurious in Southern Illinois and parts of Missouri, 33—It is less injurious to apples than the plum Curculio, 33—Injury done by it to pears, 33—The rot in apples principally produced by it and by the plum Curculio, 34—Season of its appearance, 34—Remedies and preventive measures, 34—Description of larva and pupa, 35.

The plum-gouger, Coccotorus scutellaris

How it differs from the preceding species, 39—Injury caused by it,
39—Food-plants, 40—It is often mistaken for the plum Curculio,
40—Season of its appearance, 40—Holes bored by it in the fruit,
41—The larva lives within the kernel of the fruit, 41—Remedies, 41

The pea-weevil, Bruchus pisi

Insect enemies of the garden-pea, 44—Characters of the Bruchidæ,
45—Habits of other species of Bruchidæ, 45—Frequent occurrence
of its larva in green peas, 45—Characteristics of the beetle, 46—
It is in all probability an indigenous North American insect, 46—
The beetle does not sting the peas, 46—The eggs are fastened by
the female beetle on the outside of the pod, 47—The larva and
its habits, 47—Its transformations, 47—Remedies and preventives, 48—Examination of peas intended for seed, 48—Concerted
action necessary to exterminate the insect, 48—Mr. Saunders's account of the occurrence of the pea-weevil in Canada, 49—Other
preventive measures, 49—Birds destroying the insect, 50.

1301. RILEY, C. V.—Continued.	
The corn Sphenophorus, Sphenophorus sculptilis	59
The cocklebur Sphenophorus, Rhodobanus 13-punctatus	60
Insects injurious to the Grape-vine The grape leaf-folder, Desmia maculalis Its geographical distribution, 61—Generic characters, 61—Characters of the moth, 61—Sexual differences, 62—Habits of the larva, 62—Remedy and prevention, 62—Natural enemies, 62—Description of the larva, 62.	61 61
The grape-vine epimenis, Psychomorpha epimenis. Its larva formerly mistaken for that of the pearl wood nymph, 63— Characters of the moth and sexual differences, 63—Habits of the larva when about to transform, 64—Its grape-vine feeding habits formerly unknown, 64—Description of the larva, 64; of the chrysalis, 65.	63
The grape-vine plume, Oxyptilus periscellidativius	65
The common yellow bear, Spilosoma virginica	. 68
The smeared dagger, Acromycta oblinita. The larva is polyphagous, 70—Characters of the larva, pupa, and imago, 70—Remedy, 70—Parasites, 71—Description of imago and larva, 71; of the pupa, 72.	70
The pyramidal grape-vine worm, Pyrophila pyramidoides	72
The grape-root borer, Sciapteron polistiformic Its distribution, 75—Distinguishing characters of the larva, 76— Its transformation, 76—Characters of the moth, 76—Description of the image and its sexual differences, 76—Work of the larva on grape-roots, 77—Remedies, 77.	75
The spotted Pelidnota, Pelidnota punctata	77
The grape-vine flea-beetle, Haltica chalybea	79

101

	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	155
1301.	RILEY, C. V.—Continued.	
	The grape-vine Colaspis, Colaspis flavida Fitch's account of the insect, 82—Varieties of the beetle, 82—The larva in all probability attacks sometimes that of the leaf-folder,	81
	82-It lives on the roots of the strawberry, 82-Larva of the European Colaspis barbara, 82-Difference in habits of larvæ of allied species, 83-Singular characters of the larva of the grape-vine Colaspis, 83-Work of the larva on strawberry roots, 83-Rem-	•
	edy, 84—Description of the larva, 84. The grape-leaf gall-louse, Phylloxera vastatrix	. 84
	Its life-history not yet fully studied, 84—Previous accounts of the insect by Fitch, Shimer, and Walsh, 85—The root disease in France, 85—The Phylloxera vastatrix recognized as the cause of this disease, 85—Identity of the gall-louse with the root-inhabit-	, 0
	ing insect, 86—The American and European insects are identical, 86—Remedies tried in France, 86—The disease directly caused	
	by the Phylloxera, 87—Injury done by the Phylloxera in Missouri, 87—Forming of the gall, 87—Propagation of the lice and	
	multiplication of the galls, 88—The gall-lice descend in the lat- ter part of the season to the roots, 88—Change of the insect after	
	passing from the leaves to the root, 88—Questions still to be settled in the life-history of the Phylloxera, 88—Rare occurrence of the	
	winged form, 89—The insect can be transported from one place to	
	another on roots, 89—It hibernates on the roots, 89—Grape-vines	
	that should be planted, 89—Number of indigenous species of the	
,	grape-vine, 90—Grape-vines which are most seriously infested with the grape-leaf louse, 90—Danger in planting the Clinton	
	among other grapes, 91—Insects acquiring different food-habits	
	as illustrated in the apple-magget and the pine-leaf scale, 92—	
	The different forms of the grape-leaf louse, 93—Discussion on the	
	proper place of the insect in the classification, 93—On Dr. Shi-	
	mer's proposed new families Dactylosphæridæ and Lepidosaphidæ,	
	93—Objections to Fitch's specific name vitifolia, 95—Identity of	
	the European with the American insect, 95—The apple-root louse	
	is identical with the woolly Aphis, 95—The gall-inhabiting form of the Phylloxera identical with the root-inhabiting type, 96—	
	Characters of the genus Phylloxers and its place in the system, 96.	
	THE COLORADO POTATO-BEETLE AGAIN, Doryphora 10-lineata	79
	Its onward march, 97—It invades the Dominion of Canada, 97—	•
	How it crossed Lake Michigan, 97—It will probably spread	
	through Ontario unless preventive measures are taken, 98—Ex-	
	cellent chance to prevent its spread in Canada, 98—The Paris	
	green remedy, 99-It is efficient if judiciously applied, 99-It	
	does not affect the tuber, 99—Natural checks to the increase of	
	the potato-beetle, 100—The great Lebia destroying the larvæ, 100—Bogus experiments, 100—The true remedy, 101—How to	

early varieties of potatoes, 101. THE CODLING-MOTH AGAIN, Carpocapsa pomonella Hay-bands around the trunk of the tree more effectual than rags placed in the fork, 102-The codling-moth is single-brouded in the more northern countries, but double-brooded in the latitude of St. Louis, 102-Sexual differences of the moth, 103-Sexual characters in the genera Argynnis and Grapta, 103-The codlingmoth also infests peaches, 103.

prevent the insect from becoming too numerous, 101-Planting

1301.	RILEY, C. V.—Continued.	
1301.	RILEY, C. V.—Continued. THE CORN-WORM alias BOLL-WORM, Heliothis armigera Its geographical range, 104—Injury done by it to corn, 104—It attacks tomatoes and other plants, 105—Food-plants of the stalk-borer, 105—The egg of the corn-worm, 105—Mr. Glover's account of the habits of the boll-worm, 106—The larva is very variable in color, 107—Its transformations, 107—Number of annual broods, 107—Amount of damage done by it, 107—Remedies, 108—Attracting the moth by sweets, 108—Heard's moth-trap, 109. The fall army-worm, Laphygma frugiperda Reports of its appearance in 1870, and how it was generally mistaken for the true army-worm, 109—It was also mistaken for the boll-worm, 111—Injury caused by it, 111—How it differs from	104
	the true army-worm, 112—It is a very variable species in the image state, 113—The spiderwort owlet-moth, and how it differs from the fall army-worm moth, 113—Number of annual broads and time of appearance of the fall army-worm, 114—The eggs and how they are deposited, 114—Preventive measures, 115—It is never injurious during two consecutive years, 115—Parasitic checks, 116—Description of the image, 116; of the varieties and the earlier states, 117.	
	THE APPLE-TREE TENT-CATERPILLAR, OR AMERICAN LACKEY-MOTH, 'Clisiocampa americana	117
	Parasites and enemies, 120. The Tent-Caterpillar of the forest, Clisiocampa disstria The egg-mass and how the eggs are deposited by the female moth, 121—Development of the larva, 122—Fitch's description of the full-grown larva, 123—Confusion arising from want of uniform rule in describing larvæ, 123—The cocoon, 124—The chrysalis and the moth, 124—The web spun by the caterpillar, 124—Mr. Ferris's observations on differences in habits, appearance, and food-plants of the caterpillar, 125—Phytophagic varieties or species, 127—Food-plants of the caterpillar, 127—Its destructive powers, 127—Remedies, 128—Natural enemies and parasites, 128—Summary, 129.	121
	THE FALL WEB-WORM, Hyphantria cunea	130
	THE BLUE-SPANGLED PEACH-WORM, Callimorpha lecontei	132
	THE ASH GRAY PINION, Lithophane cinerea Food-plants of the larva, 135—Transformations of the insect, 135— Characters of the moth, 135—Description of larva and imago, 135—Remarks on allied species, 136.	134

BENEFICIAL INSECTS.

THE GLASSY-WINGED SOLDIER-BUG, Campyloneura vitripennis It preys upon different species of leaf-hoppers, 137—Its larva and pupa, 138—How it seizes its prey, 138—Coloration of the insect, 138—It was never observed before to attack the leaf-hoppers of the grape-vine, 139—Habits of the Phytocorida, 139.	137
INNOXIOUS INSECTS.	
THE WHITE-LINED MORNING SPHINX, Deilephila lineata	140
Two of our common butterflies	142
Synonymy of its specific name, 143—Its geographical distribution, 143—Characters of the Danaida, 143—Sexual differences, 143—Food-plants, 144—Hibernation, 144—Two annual broods, 144—Description of the egg, 144—The larva and its larval changes, 145—How the horns of the larva become longer at each molt, 145—The full-grown larva, 146—Interest attached to the metamorphoses of insects, 146—How the larva becomes a chrysalis, 147—The hardened chrysalis, 147—Duration of the chrysalis state, 147—The issuing of the butterfly, 148—Protective colors of insects, 146.	143
sects, 148—Nauseous odor of the archippus butterfly in all its stages, 149—The Tachina-parasite of the archippus larva, 149—Action of the parasitized larva, 149—How the Tachina larva and other insects prepare the lid of their puparia, 149—Characters of this Tachina-fly, 150—Difficulties of the study of the Tachinariæ, 150—Two forms occurring in Tachina archippivora, 150—The butterfly often congregating in immense swarms, 151—Probable reasons of this assembling in swarms, 152.	
The disippus butterfly, Limenitis disippus	153
Distinguishing characters of the Nymphalidæ, 153—Food-plants and geographical range, 153—The egg, 153—Description of the egg, 154—Development of the larva, 154—Description of the mature and young larva, 154—The insect hibernates as young larva, 155—Case prepared by the larva for its winter quarters, 155—Modifications of the case, 156—Peculiar habit of the autumnal larvæ, 156—Parasites, 157—Tachina-fly, 157—The Disippus eggparasite, 157—The disippus Microgaster, 158.	,
Mimicry as illustrated by these two butterflies, with some remarks	
on the theory of natural selection	159
the archippus and disippus butterflies, 163—The ursula butter-	
fly, 163—Its caterpillar and chrysalis undistinguishable from those of the Disippus, 163—Its image has no protective color, 164—Mr. Bennett's objections to the theory of natural selection producing mimetic forms, 165—Mr. Scudder's objections, 166—	(

Discussion of Mr. Scudder's arguments, 167-Mr. Murray's ob-

- 1301. RILEY, C. V.—Continued.
 - jections to the connection between mimetic resemblances and natural selection, 170—Natural selection not the only power producing mimicry, 171—Reasons for discussing in this report the theory of natural selection, 172—Natural selection involves belief in the doctrine of evolution, 173—Darwiniam is neither irreligious nor atheistic. 174.
- 1302. RILEY, C. V. Snout-beetles injurious to fruits. <Trans. Ill. Hortic. Soc. for 1870, 1871, n. s., v. 4, pp. 89-124, figs. 1-11. Reprint, with additions and omissions: <3d Ann. Rept. State Ent. Mo., April, 1871, pp. 5-44, figs. 1, 3, 4, 7-14. See No. 1301 for synopsis of contents.
- 1303. RILEY, C. V. Bark-lice on rose bushes. <Moore's Rural New Yorker, 24 June, 1871, v. 23, p. 393. S.-b. No. 4, p. 3.

 Answer to inquiry of W. A. French; occurrence of and means against Locanium rosæ [= oleæ] and Diaspis rosæ on rose bushes.
- 1304. RILEY, C. V. Canker-worms; not army-worms. < Moore's Rural New Yorker, 24 June, 1871, v. 23, p. 393.
- 1305. RILEY, C. V. How to distinguish between Limenitis disippus Godt., and L. ursula Fabr., in their preparatory states. <Ca. Ent., July, 1871, v. 3, pp. 52-53, fig. 24.
 - Description and figures of structural characters distinguishing the larva and pupa of *Limenitis disippus* from those of *L. ursula*; see No. 1306.
- 1306. RILEY, C. V. Friendly notes. <Ca. Ent., September, 1871, v. 3, pp. 117-119.
 - Distinctive structural characters of pupse of Limentis disippus and L. ursula; criticism of W. Couper's articles; description, food, and habits of Gelechia [=Holcocera] glandulella n. sp.; review of statements concerning the dimorphism of Grapta interrogationis.
- 1307. RILEY, C. V. Friendly criticism. <Gardener's Mo. and Hortic., November, 1871, v. 13, p. 341.
 Inefficacy of the roller remedy for potato-beetles and grasshoppers; identifies "Meloloniha philophaga" as Lachnosterna quercina [=fusca].
- 1308. RILEY, C. V. Parasites on fowls. The chicken mite. < Moore's Rural New Yorker, 16 December, 1871. S.-b. No. 5, p. 31.</p>
 Answer to inquiry of H. Hales; injuries to fowls caused by Dermanyssus galling? [= avium].
- 1309. RILEY, C. V. The American Entomologist. <Cultivator and Country Gentl., 21 December, 1871, v. 36, p. 809. Reprint: <Gardener's Mo. and Hortic., January, 1872, v. 14, p. 23. <Ca. Ent., January, 1872, v. 4, p. 19.
 - Announcement of the continued suspension of publication of the American Entomologist.
- 1309a. [RILEY, C. V.] (Lxodes bovis.) < Rept. Comm. of Agric. on the diseases of cattle. Washington, 1871, p. 118, fig. Description and figure of Lxodes bovis n. sp.</p>
- 1310. RILEY, C. V. The acorn moth. Holcocera glandulella, n. sp. <Ca. Ent., January, 1872, v. 4, pp. 18, 19.
 - Description of larva and image of Holcocera glandufella n. sp.; habits of larva; food-plant and description of larva of Balaninus rectus.

1311. RILEY, C. V. Fourth annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <7th Ann. Rept. State Board of Agric. for 1871, April, 1872, pp. 146+6, 66 figs. Separate: <Jefferson City, Mo., April, 1872, pp. 146+6, 66 figs. Notice: <Horticulturist, August, 1872, v. 27, p. 251. <Ent. Mo. Mag., July, 1872, v. 9, p. 47.

CONTENTS.

^{*}Extract in <Sci. Amer., 25 May, 1872, v. 25, p. 351.

1911	DIVEY C V Continued	
1011.	RILEY, C. V.—Continued. mies, 28—The Pennsylvania soldier-beetle and its larva, 28—The	
	two-lined soldier-beetle and its larva, 29—Description of the	
	larva of Telephorus bilineatus, 30—Summary, 30.	
	The periodical Cicada, Tibicen septendecim	30
	Reproduction, with corrections and additions, of the chronolog-	
	ical table of Broods I to VI from Report I. 31.	
	The grape-vine Colaspis again, Colaspis flavida	34
	Redescription of the larva from well-preserved specimens, 34.	-
	THE HABLEQUIN CABBAGE-BUG, Murgantia histrionica	35
	It was not known in Missouri prior to 1870, 35—Its geographical	
1	range and color variations, 35-Insect enemies of the cabbage-	
	plant, 35—Dr. Lyncecum's account of its habits and injury	
	caused by it in Texas, 36—Its appearance in Missouri in 1870,	
	36—The egg, 37—The larva and pupa, 37—Several annual broods,	
	37—The mature bug, 37—Injury caused by it, 37—Its congener	
	in Europe, 38.	
	THE RASCAL LEAF-CRUMPLER, Acrobasis indignella	36
	Its proper scientific name, 38—It is hardly noticed in summer time,	
	38—Injury caused by it, 39—It hibernates as larva, 39—Habits of the larva, 39—The larval case, 39—Characteristics of the moth,	
	39—Food-plants, 39—Remedies, 40—Natural enemies, 40—Le	
	Baron's description of Tachina physics, 40—Limneria fugitiva, 41—	
	Description of imago, larva, and chrysalis of the rascal leaf-	
	crumpler, 41—Description of the variety nebulella, 42.	
	THE WALNUT CASE-BEARER, Acrobasis juglandis	42
	Other case-bearers enumerated, 42—The case of the walnut case-	
	bearer, 42—Differences between the moth and that of the rascal	
	leaf-crumpler, 43—Natural enemies, 43—Description of Perilitus	
	indigator, 43.	
	THE APPLE-LEAF SKELETONIZER, Pempelia hammondi	44
	Its work on the leaves of apple-trees, 44—The worm and the chrys-	
•	alis, 44—Appearance of the moth, 45—Mr. Hammond's account	
	of the injury caused by the worm, 45—The European Acrobasis consociella, 45—Remedies and parasites, 45—Description of the	
	imago, larva, and pupa, 46.	
	THE GREEN APPLE LEAF-TYER, Teras cinderella	46
•	It occurs almost always in company with the foregoing, 46—	•0
	Characters and habits of the worm, 46—The chrysalis, 47—De-	
	scription of imago, larva, and chrysalis, 47.	
	THE LESSER APPLE LEAF-FOLDER, Toras malivorana	47
	Its larva and pupa closely resemble those of the foregoing species,	
	47—Mr. Wier's account of its habits, 48—Remedy, 49.	
	THE APPLE-LEAF BUCCULATRIX, Bucoulatrix pomifoliella	49
	It is not very injurious in Missouri, 49—Account of damage caused	
	by it in New York, 50—The worm and its habits, 50—Its trans-	
	formations, 50—Season of the appearance of the moth, 50—Rem-	
	edies, 50—Habits of Bucculatrix thuislla, 51—Description of larva	
	and pupa of the apple-leaf Bucculatrix, 51.	E 1
	THE APPLE-TWIG BORER, Amphicerus bicaudatus	51
	52—The holes made by it in the twigs, 52—The holes are made	
	only for food and protection, 52—The insect breeds probably in	
	the sap-wood of forest trees, 52—The larva of Sinoxylen basilare	
	mistaken for that of the apple-twig borer, 52—Remedy, 53.	

72 72

BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	161
1311. RILEY, C. V.—Continued.	
INSECTS INJURIOUS TO THE GRAPE-VINE	53
The red-shouldered Sinoxylon, Sinoxylon basilare	53
Characteristics of the insect as imago, larva, and pupa, 54-Dam-	
age done by it to grape-vines, fruit trees, and hickory, 54-De-	
scription of the larva and pupa, 54.	
GRAPE DISEASE	55
The grape-leaf gall-louse, Phylloxera vastatrix	55
Its scientific name, 55-The law of priority, 55-European grape-	
vines valueless in the eastern United States, 55-Deterioration	
of some of our native vines, 56—Climatic reasons for the failure	
of European vines, 56—The principal cause of this failure is the	
Phylloxera, 56—Further proof of the identity of the American	
with the European insect, 57—Reasons for the identity of the	
gall-louse with the root-louse, 57 - Further facts respecting	
the habits of the root-louse, 58—Underground forms, 58—The	
young lice and their habits, 59—Hibernation, 59—The pupa, 59—	
The winged female, 59—Susceptibility of different vines to the	
attacks of the louse, 60—Classification of the North American	
grape-vines according to their practical importance, 60—Synopsis	
of the true grape-vines of the United States, by Dr. George En-	
gelmann, 60—Difficulty of separating the cultivated varieties of	
grape-vine, 61-Importance of a proper classification of culti-	
vated grape-vines, 62-Importation into Europe of resisting	
American vines, 62—Enumeration of the cultivated varieties and	
their susceptibility to the disease, 63—No variety is entirely ex-	
empt from the attacks of the root-louse, 64—American vines	
which resisted the root-louse in Europe, 64—Means of contagion	
from one vine to another, 64—The spread of the root-lice from	
one vine to another, 64—Transportation of the louse upon the	
roots of seedlings and cuttings, 64—Spread of the disease in	
France, 64—The winged female of the root-louse and her func-	
tion, 65—The vernal leaf-gall, 65—Preference of the winged	
Phylloxera for the Clinton grape and its allies, 65—Power of	
flight of Phylloxera caryafolia, 66—Deposition of the eggs upon	
the leaf, 66—Probable reasons why the injuries of the Phylloxera	
are greater in Europe than in America, 66—Ontward and more	
visible effects of the root-disease, 67—Practical suggestions, 67—	
No need to destroy the Clinton vines, 67—Influence of soil on	
the intensity of the disease, 67—Remedies, 68—Destruction of	
the gall-lice, 68—Destruction of the root-lice, 68—Results of ex-	
periments in France with various substances, 68—Irrigation and	
submersion, 69—Résumé of the insect's history, 69—No need of	
unnecessary alarm, 70.	

BENEFICIAL INSECTS.

Introductory
The Morus multicaults fever and its reaction, 72-Increasing atten-
tion lately given to silk-culture in America, 72-North America
well adapted to the raising of silk, 73—General outlines of the
natural history of the eight species of silkworms treated of in
this report, 74.

The mulberry silkworm, Sericaria mori.....

75

Its past history, 75-Earliest silk-culture in China and India. 75-Its introduction into Europe, 76-Value of silk produced in France, 76-The "Silk Supply Association" in England, and its objects, 76-Countries exporting raw silk, 77-The name given to silk by different nations, 77-History of the mulberry silkworm in America, 77-Its introduction and failure of earlier efforts, 77-Renewal of silk-culture within the past decade, 77-Silk manufacture in the United States, 78-Favorable prospects for raising silk in this country, 78-Silk-growing in California, 79-Mr. Prevost's "California Silk-grower's Manual," 79-False statements and exaggerations in Prevost's book, 79-Sale of Californian silkworm eggs, 80-Disastrous effect of the Franco-Prussian war on the egg trade, 81-Success of silk-culture in Calitornia dependent on the ability to reel the silk, 81-Silk-culture in Kansas, 82-Mons. Boissière's silk establishment and its chances of success, 82-Silk-growing in Missouri, 83-The fall season in Missouri eminently propitious for rearing silkworms, 84-Natural history of the silkworm, 84-Races of the silkworm produced by domestication, 84-Effects produced on the insect by domestication, 85-The egg, 86-Larva and larval changes, 86-Cocoon, chrysalis, and moth, 87-No insect parasite of the silkworm in Europe, 87-The "Uji" disease in China and Japan, 87-Diseases of the silkworm, 88; the Muscardine, its effect and cause, 88; the Pébrine disease, its symptoms and cause, 89; nature and origin of these diseases, 90-Other diseases of the silkworm, 91-Best varieties or races, 90-Different forms of cocoons produced by different races, 92-How best to rear silkworms, 92-Rearing a very simple process, 92-Character of climate of the Japanese silk districts, 93-Keeping the eggs during the winter, 93—Hatching of the eggs, 93—Room and building for the rearing of the worms, 93-The feeding net or fillet, 94-Importance of carrying all the worms simultaneously through their molts, 94-Regularity of feeding, 95-Regulating the temperature, 95—Cocoonery, 96—Choking the chrysalis, 96—Egglaying, 97-Selecting and fastening the cocoons for breeding purposes, 97-Treatment of the female moths after copulation, 97-Preservation of the eggs, 97-Reeling, 98-Great skill required to accomplish the work properly, 98-Classification of raw silk, 98-Preparing the cocoons for reeling, 98-Objects of and manipulations in reeling, 99-Best food for the worms, 100-Varieties of the mulberry, 100-Cultivation of the mulberry, 100-Osage . orange as silkworm food, 100-Introduction of the osage orange into France, 100-Experiments in America with feeding silkworms on osage orange, 101-Advantages and disadvantages of the osage orange, 102.

The cecropia silkworm, Attacus cecropia.....

Changes made in its scientific generic name, 103—General appearance of the moth, 103—Fitch's explanation of the specific name, 104—Food-plants, 104—The cocoon, 104—Value of the cocoon as compared with that of the polyphemus moth, 105—How the moth issues from the cocoon, 105—The moth immediately after hatching, 106—The egg, 106—Description of the larval changes,

106—Strange habit of birds of using the empty cocoon as a store-house, 107—The cecropia worm can not be classed as an injurious insect, 107—Samia columbia ought to be considered a variety of cecropia, 107—Parasites, 107—The long-tailed Ophion, 107; its mode of oviposition, 108; habits of its larva, 108—The cecropia Tachina-fly, 108; how it affects its victim, 108; its larva and imago, 109—The Mary Chalcis-fly, 109; how it escapes from the cocoon of the moth, 109; description of the imago, 110—The cecropia Cryptus, 110; its habits, 110; description of the two sexes, 111; how it differs from the allied species, 111.

The yama-maī silkworm, Antherœa yama-mai

Its native home and food-plant, 130—Its introduction into Europe and Australia, 130—Attempts at raising it in America, 130—The egg and how it should be kept over winter, 131—Climate most favorable for the worm, 131—The worm and its resemblance to the leaf, 132—Food-plants, 132—Larval changes, 132—The cocoon and the value of its silk, 133—The moth and its habits, 133—Difficulties of rearing the worm, 133—Mr. Adams's account of rearing the yama-maī worm in Japan, 134—The Uji parasite, 136.

The perny silkworm, Antheræa pernyi

Its native home, 137—How it differs from the preceding, 137—Larval changes, 137—The cocoon and its silk, 137—The moth, 137—Its culture in China, 138—The tusseh silkworm, 138.

112

121

123

125

130

1311.	RILEY, C. V.—Continued.	
	Summary Successful silk culture possible in this country, 138—Comparative value of the different species of silkworms, 138.	13 8
	INNOXIOUS INSECTS.	
	THE HORNED PASSALUS, Passalus cornutus. Its frequent occurrence in old logs, 139—The noise produced by the beetle, 139—It occurs only in decaying wood, 140—The larva and its exceptional character, 140—Description of the larva, 140—Previous description of Passalus larvæ, 141—Description of the pupa, 141.	139
	THE GREAT LEOPARD MOTH, Eopantheria scribonia	141
	THE ISABELLA TIGER MOTH, Pyrrharotia isabella	143
·	THE ACORN MOTH, Holcocera glandulella The egg laid in acorns that have been ruined by Balaninus rectus, 144—Habits of the larva, 144—Characters of larva and moth, 145—Description of imago, larva, and pupa, 145.	144
1312.	RILEY, C. V. Thomas Wier's apple-worm trap. Amer. Agr April, 1872, v. 31, pp. 142–143, 1 fig. Description and figure of Wier's apple-worm trap.	ric.,
1313.	[RILEY, C. V.] Remarkable parasitic fungus. <sci. 1872="" 26,="" 347.="" 40],="" [v.="" amer.="" and="" description="" figure="" fungus="" fusca.<="" grub,="" infesting="" lachnosterna="" larva="" may,="" n.="" of="" p.="" s.,="" td="" the="" unidentified="" v.="" w=""><td></td></sci.>	
1314.	RILEY, C. V. Cut-worm lion. <colman's 15="" ju<br="" rural="" world,="">1872, fig. Sb. No. 5, p. 47. History of the larva and image of Calesoma calidum, an enemy to cut-work</colman's>	
1315.	RILEY, C. V. Cut-worms. < Cultivator and Country Gentlem 20 June, 1872, v. 37, p. 392. Dandelions and other weeds furnish food for young cut-worms which he in the fall; land should be kept clear of weeds at that season.	·
1316.	RILEY, C. V. Flat-headed apple tree-borer in horse chestrest Colman's Rural World, 22 June, 1872, fig. Sb. No. 5, p. Answer to inquiry of S. S. R.; natural history of larva and image of Chebothris femorata.	47.
1317.	RILEY, C. V. A new insect. < Western Planter, 29 June, 18 Sb. No. 5, p. 49. Answer to inquiry of F. Halsinger; Nysius n. sp., very destructive to tatoes.	
1318. 1	RILEY, C. V. Codling-moth; jarring down infested fr <cultivator 1872,="" 37,="" 4="" 4<br="" and="" country="" gentleman,="" july,="" p.="" v.="">Excrement outside of the fruit indicates the exit of the worm, with a</cultivator>	122. ome

are visible.

- 1319. RILEY, C. V. Food for silkworms. <Colman's Rural World, 13
 July, 1872. S.-b. No. 5, p. 40.
 Osage orange superior to red-mulberry.
- 1320. RILEY, C. V. The apple maggot-fly, Trypeta pomonella Walsh. Amer. Agric., July, 1872, v. 31, pp. 263-264, 2 figs.

 Answer to letter of J. H. Spatter; comparative descriptions and figures of larvæ, pupæ, and imagos of Trypeta pomonella and Carpocapsa pomonella; figures showing injuries of both species; distribution and means against Trypeta pomonella.
- 1321. RILEY, C. V. Worms on Dutchman's pipe. <Colman's Rural World, 3 August, 1872, fig. S.-b. No. 5, p. 136.

 Answer to inquiry of J. T. C.; life-history of Papilio philenor.
- 1322. RILEY, C. V. Apple-leaf worm. The apple-leaf skeletonizer. <Colman's Rural World, 10 August, 1872, fig. S.-b. No. 5, p. 138.
 - Answer to inquiry of L. R. Bryant; habits and times of appearance of Pempelia hammondi.
- 1323. RILEY, C. V. Eggs in grape-canes and apple-twigs. <Amer. Agric., August, 1872, v. 31, p. 302, figs. 1-7.
 - Figures of eggs and of twigs punctured by Orocharis saltator, Œcanthus niveus, and Ceresa bubalus; figures images of the same; habits, descriptions of eggs, and images of the three'species.
- 1324. RILEY, C. V. Borers in evergreens. <Gardener's Mo. and Hortic., December, 1872, v. 14, p. 373.
 - Larvæ of Monohammus titillator the worst enemy of the white-pine; another species of Corambycidæ injurious to red-cedars; a host of bark-borers affect all kinds of evergreens.
- 1325. RILEY, C. V. Einige unserer schädlicherer Insekten. <St. Louis, Mo., 1872, p. 35, il. S.-b. No. 19, pp. 174-190.
 - Causes of the decadence of viticulture in the United States; habits, ravages, specific identity, and figures of several stages of the leaf-inhabiting and root-inhabiting forms of Phylloxera vastatrix; identity of the European and American grape Phylloxera; susceptibility of the different species of grape to the attacks of the same; principal manifestations of attack; preventive and remedial measures; description and figures of Conotrachelus nemsphar; habits, ravages, parasites, enemies, and means against it; description of Ransom's trap, Hull's, Ward's, and Hooten's machines for capturing the images; figures of the machines. Natural history, food-plants, distribution, enemies, parasites, and means against Doryphora decembinata; figures of all stages of the same; use of Paris green and other powders; mechanical devices for collecting the insects.
- 1326. RILEY, C. V. Harvest mites. < Amer. Nat., January, 1873, v. 7, pp. 16-19, fig. 5. Extract: < Colman's Rural World, 21 June, 1876, 2 figs. S.-b. No. 10, pp. 170, 171.
 - List of insect parasites on man; description, figures, habits of and means against Leptus [= Tetranychus] americanus n. sp. and L. [= T.] irritans n. sp.
- 1327. RILEY, C. V. Vanessa antiopa. <Ent. Mo. Mag., January, 1873, v. 9, p. 195.
 - Habits and food-plants of Vanessa antiopa.

1328. RILEY, C. V. Entomological correction. <Cultivator and Country Gentleman, 6 March, 1873, v. 38, p. 149.

Lema trilineata does not occur in southwestern Missouri, as stated by W. R. Howard.

1329. RILEY, C. V. Fifth annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <8th Ann. Rept. State Board of Agric. for 1872, 18 April, 1873, pp. 160+8, 75 figs. Separate: <Jefferson City, Mo., 18 April, 1873, pp. 160+8, 75 figs. Review by A. S. Packard, jr.: <Amer. Nat., August, 1873, v. 7, pp. 471-477, figs. 115-130. Reply to Packard and rejoinder by Packard: <Amer. Nat., March, 1874, v. 8, pp. 181-188.

CONTENTS.

3

46

46

Entomology, its relations to agriculture and its advancement* Definition of entomology, 5-The term "insect," 5-The four subkingdoms of the animal kingdom, 6-Characteristics of insects, 7—Classification of insects, 8—Hymenoptera, 9—Coleoptera, 10-Lepidoptera, 11-Hemiptera, 12-Diptera, 13-Orthoptera, 14—Neuroptera, 14—Osculant and aberrant groups, 15—Importance of entomology as a study, 17-Economic importance of entomology, 18-Relation of insects to agriculture, 18-Damage inflicted by insects to agriculture and horticulture, 19-Progress of economic entomology, 19-Economic entomology in Europe, 19—Insects in America much more injurious than in Europe, 21— State entomologists and entomologists in public positions in the United States, 21-How to counterwork noxious insects, 23-Prevention, 24—Checking the spread of injurious insects, 24— Cure, 25-Modification of Dr. Hull's Curculio-catcher, 25-Encouragement of the natural enemies of injurious insects, 26-Duties of a State entomologist, 27-Annual report, 27-Correspondence and other duties, 28-How to collect, preserve, and study insects, 29-Appliances for collecting, 29-Use of the umbrella, 29-The knapsack, 30-The hand net, 30-The frame of the net, 30—The bag of the net, 31—Use of the sieve, 31—Sugaring, 32-Attracting moths by light, 32-How to kill insects, 32-The cyanide bottle, 32-Use of chloroform, 33-Entomotaxy, 34—Insect-pins, 34—How to pin insects, 34—How to mount small insects, 34—Spreading-board, 35—Drying-box, 36—Insect boxes and cabinet, 37—Mr. Lintner's boxes for Lepidoptera, 38—Substances for lining insect-boxes, 40-Preserving insects in the collection, 41-Relaxing specimens, 41-Breeding insects, 41-Breeding-cage, 42-Note-book of the collector, 43-How to transmit insects, 44-Text-books, 44.

NOXIOUS INSECTS.

NOTES OF THE YEAR

^{*}Pages 17-27 were reprinted under title Eutomology in <Rept. State Board Agric. Kansas for 1873, 1874, pp. 266-274.

•	
1329. RILEY, C. V.—Continued.	
worm in California, 49-Enemies and parasites, 49-The ring-	
legged Pimpla, 49-The delicate long-sting, 50-Other enemies,	
51—Efficiency of the spined soldier-bug, 51—False doctrines for	
exterminating the codling-moth, 51.	
The Colorado potato-beetle, Doryphora 10-lineata	52
Its comparative harmlessness in 1872, 52—New food-plants, 52—	
Its progress eastward up to 1872, 52—Experiments with Paris	
green, 53—New enemies, 53—The rust-red social wasp, 54—The	
rose-breasted grosbeck, 54.	
The apple-twig borer, Amphicerus bicaudatus	54
It attacks also pear twigs, 54—It has been bred from grape canes	
by Dr. Shimer, 54.	
Egg of the horned Passalus, Passalus cornutus	55
The egg, 55—The newly hatched larva, 55—Rapid development of	
the insect, 55.	==
Egg of the common May beetle, Lacknosterna fusca	55
Description of the eggs and how they are deposited, 55.	E 0
Egg of the broad-necked Prionus, Prionus laticollis	56
Characteristics of the eggs and where they are deposited, 56.	56
Eggs of American tent-caterpillar, Clisiocampa americana	90
Counterworking the tobacco-worm, Protoparce celeus	56
Mr. White's method of planting the Jamestown weed among the	30
potatoes and poisoning the blossoms thereof, 56.	
THE GRAPE PHYLLOXERA, Phylloxera vastatrix	57
Its popular name, 57—Accounts of the unusual mortality among	0.
grape-vines in the spring of 1872, 57—Causes of this mortality	
given in the accounts, 59—Excessive drought and overbearing,	
60—The Phylloxera is the true cause, 60—Actual proof of the	
Phylloxera having caused the mortality among grape-vines, 61—	
Influences that favored the increase of the lice in 1872, 61—Im-	
portance of a full understanding and management of the Phyl-	
loxera, 62—Its range in North America, 62—Its spread in Europe,	
63-Inconstancy in the habits of the gall-lice, 63-The leaves of	
the Clinton vine no longer affected by the gall-lice since 1871,	
63-Method of formation of the leaf-gall, 64-Relative immunity	
of American vines in Europe, 64—Propagating American varie-	
ties from cuttings, 65-Appreciation of American vines in Eu-	
rope, 66-Careless statement published as to the immunity of	
Labrusca vine, 66-Grafting the grape-vine, 67-New theories,	
67-The Phylloxera is the true cause and not the effect of the	
disease, 67—Mr. Laliman's theory that the insect has always ex-	
isted in Europe, 68-Oidium tuckeri of Europe and America iden-	
tical, 69-Mr. Saunders's account of the presence of Oidium tuckeri	
in America, 70—Means of contagion of the disease from one vine to	
another, 69—Flying capacity of the winged Phylloxera, 70—The	
male louse, 71—Remedies, 71—Efficacy of carbolic acid and soot,	
71—Value of submersion or irrigation, 72—Sprinkling with quick-	
lime, ashes, etc., 72-Mr. Lichtenstein's experiments to allure	
the lice, 72—Experiments with carbolic acid, 73.	
THE OYSTER-SHELL BARK-LOUSE OF THE APPLE, Mylilaspis pomorum	73
Its occurrence in Missouri, 74-Its appearance in Luray County,	
Missouri, 74-Mr. Hauan's account of its spread, 75-Its occur-	

rence in Southern Missouri, 76—Its occurrence in Mississippi and Georgia, 78—Its appearance in Kansas, 79—It is double-brooded

in the South, 79-The waxy secretion of Homoptera, 80-The

1329. RILEY, C. V .- Continued.

newly hatched louse, 80-The larval scale, 81-Development of the female scale, 81—Growth of the male scale, 82—The male . louse, 83-Rare occurrence of the winged male, 84-Agamic multiplication, 85-Mode of spreading, 85-Food-plants, 86-Varieties of the apple-tree preferred by it, 86-Enemies and parasites, 87; mites, 87; Apheliaus mytilaspidie and Dr. Le Baron's account of its habits, 88-Easy transportation and introduction of the Aphelinus, 90-Remedies, 90-Application of oily substances, 90—Bibliographical and descriptive, 91—The generic name, 91— Signoret's classification of the Coccide, 92—Specific name, 92— Characteristics and habits of three allied species, 93-A new name necessary for our apple-tree species, 94—Description of the eggs and the winged male, 95; of the male and female scale and of the female louse, 96. **97** THE PINE-LEAF SCALE-INSECT, Chionaspis pinifolii Nature of the malady caused by it, 97-Natural history of the insect, 98—The male, 99—There are two annual broods, 99—It is confined to the pines proper, 100-Natural enemies, 100; the twice-stabbed lady-bird, 100; the painted lady-bird and description of its larva, 101-Remedies, 101-Stripping the old leaves, 102-Application of powdered and liquid substances, 102. THE HICKORY BARK-BORER, Scolytus 4-spinosus 103 Accounts of the damage caused by it in Missouri, 103-Habits of the European Scolytus destructor, 104-Various kinds of hickory attacked by the hickory bark-borer, 105-Its natural history, 105-Natural enemies, 106-Description of the three-banded Spathius, 106-Mr. Cresson's description of Bracon scolytivorus, 106—Remedies, 107—Description of the image, 107—Is it different from Scolytus 4-spinosus ? 107. THE ROSE CHAFER, Macrodactylus subspinosus..... 108 Great damage caused by the beetle in 1872, 108—Remedy, 109— Harris's account of its natural history, 109. THE FALSE CHINCH-BUG, Nysius angustatus 111 lt was not known as injurious before 1872, 111-Accounts of injury

^{*}Pages 119-125 were reprinted under title Insect punctures, in < Rept. State Board Agric. Kansas for 1873, 1874, pp. 274-279, figs. 47-59.

L 329.	RILEY, C. V.—Continued.	
	Probable eggs of the jumping tree-cricket. Orocharis saltator, 119-	
	The egg-punctures on grape-cane, 119—Characters of the egg, 119—General appearance of the cricket, 120.	
	Eggs of the snowy tree-cricket, Counthus niveus, 120-Trees and	
	shrubs attacked by this cricket, 120—The egg, 120—Habits and	
	natural history, 120—Injury done by it, 121—The eggs mistaken	
	by Fitch for those of the Buffalo tree-hopper, 121. Egg-punotures of the Buffalo tree-hopper, Caresa bubalus, 121—De-	
	velopment of the larva, 121—Characters and habits of the per-	
	fect insect, 122—Egg-punctures of some unknown tree-hopper,	
	192.	
	Egg-punctures of the frosted lightening-hopper, Paciloptera prui-	
•	ness, 122—Development and habits of the insect, 122.	
	Egg-punctures probably of Orchelimum glaberimum, 123-The egg,	
	123—Characters and habits of the imago, 123.	
	Eggs of the oblong-winged katydid, Amblycorypha oblongifolia,	
	123—They have been mistaken by Harris for those of the broad-	
	winged catydid, 123—Ovipositors and modes of egg-laying in	
	the three katydids occurring in Missouri, 124—Increase in thick-	
	ness of katydid eggs before hatching, 124. STINGING LARVÆ	125
	The potato-worm falsely considered as a stinging caterpillar, 125—	
	General harmlessness to man of insect larvæ, 125—Stinging cat-	
	erpillars in Europe and North America, 126-List of such larvæ	
	in United States, 126-Description of larva and chrysalis of	
	Acronycta xylinoides, 126.	
	The buck moth or mais moth, Hemileuca maia	127
	Flight and characteristics of the moth, 127—The egg and mode of	
	egg-laying, 128—Appearance of the full-grown larva, 129—	
	Larval changes, 129—Color variations of the larva, 130—Habits of the young larva, 130—The sting of the larva, 131—The pupa,	
	131—Issuing of the moth, 132—Irregularity and retardation in	
	development, 132; the species benefits thereby, 132—Food-plants,	
	132—Natural enemies, 132.	
	The io moth, Hyperckiria io	133
	The male and female moth, 133—The egg, 134—The larva and its	
	urticating power, 134-Larval changes, 135-Pupation, 135-	
	Food-plants, 136—Parasites, 136.	
	THE GREEN-STRIPED MAPLE-WORM, Dryocampa rubicunda	137
	Account of its occurrence in great numbers in Kansas, 137—Injury	
	done by it to soft maples, 138—The egg, 138—Larval changes, 138—The chrysalis, 139—The image, 139—Natural enemies, 139—	
	Description of Belvoisia bifasciata, 140—Remedies, 141.	
	INNOXIOUS INSECTS	
•		
	THE HELLGRAMMITE FLY, Corydalus cornuius	143
	The eggs, 143—Respiratory apparatus of the larva, 143—Habits of	
	the larva when about to transform, 143—The pupa, 144—Sexual	
	difference in the image, 145. THE GOAT-WEED BUTTERFLY, Paphia glycerium	145
	The egg, 146—Habits of the larva, 146—Larval changes, 146—	140
	Transformation, 147—New food plant, 147—Simulation of the	
	color of the food-plant by the larva, 147—Two annual broods,	
	149. Without tion of the image 149. Natural anamies 140	

On a new genus in the lepidopterous family Tineidæ with re-MARKS ON THE FERTILIZATION OF YUCCA, Pronuba yuccasella...... Generic characters of Pronuba, 150-Description of Pronuba yuccasella, 151-Plants requiring the aid of insects for fertilization. 152—Fertilization of orchids, 152—Fructification of Yucca, 153— Yuccas must rely on insects for fertilization, 153-Insects frequenting Yuccas, 154-Diurnal and nocturnal habits of Pronuba, 154-How the female moth fertilizes the plant, 154-Oviposition, 155-The larva within the young fruit, 155-Description of the larva, 155-Only a small percentage of fruit not infested by the larva, 156-The larva leaves the capsule and enters the ground for hibernation, 156-Mutual adaptation of plant and insect, 156-The moth doubtless occurs wherever Yuccas grow wild, 157--Easy transportation of the cocoon, 158--Further facts regarding the fructification of Yucca filamentosa and gloriosa, 158-Yuccas seeding in Europe, 159-Range of the insect, 159-Mr. Stainton's opinion on the characters on the Yucca moth, 160.

150

- 1330. RILEY, C. V. New York without a State entomologist. < Moore's Rural New Yorker, 5 May, 1873. S.-b. No. 8, p. 103.

Needs of a State entomologist for New York; Salix humilis infested with larva of Plectrodera scalator.

- 1331. RILEY, C. V. Tent-caterpillar of the forest. <N. Y. Tribune, 23 May, 1873. S.-b. No. 8, pp. 64-65.
 - Answer to inquiry of S. T. Gilbert; means against eggs of Clisiocampa sylvatica [= disstria].
- 1332. RILEY, C. V. Apple-tree borer. <N. Y. Tribune, 23 May, 1873, S. b. No. 8, p. 64.
 - Answer to inquiry of J. Durbin; means against Saperda bivittata [= candida].
- 1333. RILEY, C. V. Punctured grape-canes. <N. Y. Tribune, 23 May, 1873. S.-b. No. 8, p. 64.
 - Answer to inquiry of E. Snyder; food-habits and means against *Ecanthus niveus*.
- 1334. RILEY, C. V. The codling moth. Weir's trap. <Amer. Agric., May, 1873, v. 32, p. 184, figs.
 - Unsatisfactory results of experiments with the Weir trap; relative value of materials used; criticism of J. S. Parker's theory for the extermination of Carpocapsa pomonella; this species breeds in apples, wild crabs, pears, peaches, and plums; figures females and the abdomen of the males of Pimpla annulipes and Macrocentrus delicatus parasitic upon the apple-worm.
- 1335. RILEY, C. V. Influence of extreme cold on the Curculio. <Gardener's Mo. and Hortic., May, 1873, v. 15, pp. 137-139.
 - Critical review of T. T. Southwick's article of same title; extract from 3d Ann. Rept. State Ent. Mo.

- 1336. RILEY, C. V. On a new genus in the lepidopterous family *Tineidæ*, with remarks on the fertilization of *Yucca*. <Trans. Acad. Sci. St. Louis, June, 1873, v. 3, pp. 55-64, 2 figs. Reprinted, with omissions and slight changes, from <5th Ann. Rept. State Ent. Mo., 18 April, 1873, pp. 150-160, figs. 74-75. Review by P. C. Zeller: <Verh. k.·k. zool.·bot. Ges. Wien, 1876, Jahrg. 1875, Bd. 25, pp. 340-342. Reply to Zeller: <Trans. Acad. Sci. St. Louis, January-March, 1876, v. 3, pp. 325-326.
 - Description and figures of *Pronuba* n. g. and of larva and image of *P. yuccasella* n. sp. See No. 1329 for synopsis of contents.
- 1337. RILEY, C. V. Supplementary notes on *Pronuba yuccasella*. <Trans. Acad. Sci. St. Louis, June, 1873, v. 3, pp. 178-180, fig. 3. Extract: <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 131-135, fig. 3.
 - Descriptions and figures of male and female pupa of *Pronuba yuccasella*; hibernation, imagination, seasons, and distribution of the same; species of *Yucca* pollinated by it.
- 1338. RILEY, C. V. [To destroy the cotton-worm.] <Ill. Jour. of Agric., June, 1873. Reprint: <Colman's Rural World, 1873; <Rural Alabamian, July, 1873, v. 2, pp. 289-293; <Mobile Register, 1873; <Farmer's Advocate, 1873. Extract: <6th Ann. Rept. State Ent. Mo., 1874, pp. 17-18.
 - Ravages of Aletia argillacea [=xylina] in the cotton fields of Southern United States; inefficiency of measures hitherto employed against them; recommends the use of Paris green; directions for its use; natural history, description of egg, larva, and image of the Aletia; hibernation of the image.
- 1339. RILEY, C. V. [Imported plants and insects.] <Trans. Acad. Sci. St. Louis, July, 1873, v. 3, pp. 42-43 Proc.

 Verbal communication; inequality of the exchange of plants and insects be-

tween Europe and North America; extent and causes of the same.

- 1340. RILEY C. V. [Mimicry and protective resemblances.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, pp. 44-45 Proc.
 - Verbal communication; mimicry of Danais archippus by Limenitis disippus and consequent greater abundance of the latter than of L. ursula.
- 1341. RILEY, C. V. [Silk-worms fed with osage orange.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 47 Proc.

 Verbal communication; successful rearing of silk-worms on Maclura aurantiaca; exhibition of cocoons made by these worms.
- 1342. RILEY, C. V. [On the cause of deterioration in some of our native grape-vines.] <Trans. Acad. Sci. St. Louis, July, 1873, v. 3, pp. 51-52 Proc.
 - Verbal communication; abstract from 4th Ann. Rept. State Ent. Mo.; failure of grape-vines attributed largely to the ravages of *Phylloxera vastatrix*.
- 1343. RILEY, C. V. [Insects affecting the ailanthus.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, pp. 53-54 Proc. Mentions Œta compta [= punctella] and Attacus cynthia.

- 1344. RILEY, C. V. [Posthumous papers by B. D. Walsh.] < Trans.

 Acad. Sci. St. Louis, July, 1873, v. 3, p. 77 Proc.
 - Verbal communication; disposition of sundry entomological manuscripts left by B. D. Walsh.
- 1345. RILEY, C. V. [Remarks on Simulium piscicidium.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 79 Proc.
 - Verbal communication; nature and supposed ravages of Simulium piscicidium.
- 1346. RILEY, C. V. [On Antherwa yama-mai as a silk-producer.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 84 Proc.
 - Verbal communication; superiority of Antheresa yama-mai to other silk-worms tried as substitutes for Sericaria mori.
- 1347. RILEY, C. V. [On galls growing on wild sage.] < Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 84 Proc.

 Verbal communication; occurrence of three distinct undescribed galls on
- 1348. RILEY, C. V. [On a larva of Scenopinus sp. from the human lungs.] <Trans. Acad. Sci. St. Louis, July, 1873, v. 3, p. 90

Artemisia tridentata in Utah.

- Verbal communication; a larva of Scenopinus sp. found in human expectoration; usual food-habits of the larvæ of this genus.
- 1349. RILEY, C.V. Agricultural editorial excursion. < Prairie Farmer, 1873, v. 44: 2 August, p. 241; 9 August, p. 248; 16 August, p. 256; 23 August, p. 265; 30 August, p. 273; 6 September, p. 281. Reprint: < Colman's Rural World, 1873, 2, 9; 16, 23, and 30 August, 6 and 13 September. See: < Colorado Mountaineer, 1 August, 1877. S. b. No. 14, p. 102.

 Brief notes on locusts.
- 1350. RILEY, C. V. Enemies of the elm. <N. Y. Tribune, 7 August, 1873. S.-b. No. 8, pp. 56-57. Reprint: <Gardener's Mo. and Hortic., August, 1876, v. 18, p. 246.
 - Answer to inquiry of T. S. Watson; natural history of and means against Galeruca calmariensis [= xanthomelæna].
- 1351. RILEY, C. V. Entomological information. <N. Y. Tribune, 16
 August, 1873. S. b. No. 8, p. 58.
 - Answer to inquiry of J. W.; description of a number of traps for alluring and destroying insects.
- 1352. RILEY, C. V. "Controlling sex in butterflies." <Amer. Nat., September, 1873, v. 7, pp. 513-521. Separate: <Salem, Mass., August, 1873, pp. 9.
 - Critical review of Mary Treat's article of same title; females require more nourishment than males; sex determined in the egg; oviposition of Papilio and Anisota [= Dryocampa]; larvæ can not be forced to eat more than is natural to them; experiments on Thyridopteryx ephemeræformis, Orgyia leucostigma, Clisiocampa americana, Hyperchiria io, Hemileuca maia, and Anisota [= Dryocampa] rubicunda to determine the effect of the stinting of food upon the determination of sex; effect of this upon the number of molts and the longevity of Orgyia leucostigma and Megatoma serra; note on alternation of generations in Cynips [= Andricus] querous-operator and C. [= A.] querous-operatola; nature of parthenogenesis.

- 1353. RILEY, C. V. Cotton caterpillar.—Boll-worm. <Rural Alabamian, October, 1873. S. b. No. 16, p. 121.</p>
 Criticism of recent articles on Anomis [= Aletia] xylina, Heliothis armigera, and Phylloxera vastatrix.
- 1354. RILEY, C. V. On the oviposition of the Yucca moth. <Amer. Nat., October, 1873, v. 7, pp. 619-623. Abstract: <Trans. Acad. Sci. St. Louis, 10 December, 1873—25 April, 1874, v. 3, pp. 208-210. Reprint, with slight changes and omissions. <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 131-135, fig. 38.
 - See No. 1363 for synopsis of contents.
- 1355. RILEY, C. V. Phylloxera: correction. <Gardener's Mo. and Hortic., November, 1873, v. 15, p. 342.

 Critical review of report of remarks before Academy of Natural Sciences of Philadelphia.
- 1356. RILEY, C. V. Hackberry butterflies. Descriptions of the early stages of Apatura lycaon, Fabr., and Apatura herse, Fabr.; with remarks on their synonymy. <Trans. Acad. Sci. St. Louis, 10 December, 1873, v. 3, pp. 193-208, figs. 3-6. Reprint, with slight changes. <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 136-148, figs. 39-42.
 - Treats of A. lycaon [=celtis] and A. herse [=clyton]. See No. 1363 for synopsis of contents.
- 1357. RILEY, V. C. Economic entomology. <Trans. Kansas State Board Agric. for 1872, 1873, pp. 292–325, 18 figs. Extract: <Sci. Amer.
 - Characterization of entomology; importance of and means against noxious insects; list of imported noxious insects and plants; chapters on Anisota [=Dryocampa rubiounda], Eriosoma pyri [= Schizoneura lanigera], Paleacrita vernata, Galleria cercana, Estrus ovis, Bruchus pisi, Macrodaciylus subspinosus, Conotrachelus nenuphar, and Nysius destructor [= angustatus]. In general, descriptions and figures of the several stages and accounts of the habits and food-plants of and means against these insects, with some accounts and figures of their enemies and parasites, are given. Describes especially larva of Anisota [= D.] rubiounda and of Paleacrita vernata and of all active stages of Nysius destructor [= angustatus]. Figures larva, puparium, and imago of Pipiza radicum, imagos of Nothrus ovivorus, Microgaster [= Apanteles] militaris, Calosoma scrutator, C. calidum, and Blissus leucopterus and imago and nests of Eumenes fraternus.
 - Some of the chapters are based upon and some are reprinted from the 1st, 2d, and 3d Ann. Repts. State Ent. Mo.
- 1358. BILEY, C. V. Curculios on pears. <Ill. Journ. Agric., 1873.
 S.-b. No. 16, pp. 109-110, figs. 1-2.
 Habits, history, and means against Conotrachelus nenuphar, C. oratægi, and
- Anthonomus quadrigibbus; figures the two last-named species.

 1359. RILEY, C. V. Length of thread of the silk-worm. < Popular Sci.
- Monthly, February, 1874, v. 4, p. 508.

 Correction of statements in A. de Quatrefage's "Silk-worms and sericulture"

as to the length and weight of the fiber in a single cocoon of Sericaria mori.

1360.	RILEY,-C. V. A new (?) ægerian maple-borer. <amer. 123-124.<="" 1874,="" 8,="" nebruary,="" pp.="" th="" v.=""><th>ſat.,</th></amer.>	ſat.,
	Critical review of P. Gennadius's "A new ægerian maple-borer." Trochi acericolum is the well-known Ægeria acerni, injurious to maple-trees.	lium
1361.	RILEY, C. V. Entomology in Missouri. <amer. 181-188.<="" 1874,="" 8,="" mai="" nat.,="" pp.="" td="" v.=""><td>rcb,</td></amer.>	rcb,
	Reply to criticisms of A. S. Packard; discussion on the number of segment the head of an insect, on classification, and on Mytilaspis pomicon [=pomorum].	
1362.	RILEY, C. V. Economic entomology. < Amer. Nat., March, 18 v. 8, pp. 189-190.	B74,
	Demand for more attention to and better representation of the agricult interests of the country. Inadequacy of measures propounded for wholesale destruction of noxious insects.	
1363.	RILEY, C. V. Sixth annual report on the noxious, beneficial, other insects of the State of Missouri, made to the State Bo of Agriculture, pursuant to an appropriation for this purp from the legislature of the State. <9th Ann. Rept. State Bo of Agric. for 1873, March, 1874, pp. 169+12, 55 figs. Separately September 1874, March, 1874, pp. 169+12, 55 figs.	ard pose ard
	Table of contents	3
	NOXIOUS INSECTS.	
	Notes of the Year	9 9
	The Colorado potato-beetle, Doryphora 10-lineata How it has affected the price of potatoes, 11—New food-plants, 11—Its progress eastward during 1873, 12—Improved methods of using Paris green, 13—Device for jarring off the bugs, 14—Euro- pean publications on the insect, 15—Danger of its introduction into Europe, 15—Precautionary measures to be taken in Europe, 16.	11
	The cotton worm, Aletia xylina Paris green suggested as remedy, 17—Address before the National Agricultural Congress, 17—Mr. Glover's summary on experience with Paris green, 19—Experiments with the poison, 19—Johnson's sprinkling machine, 20—Patents on Paris green, 20—The	17
	Royal mixture, 21—Hibernation of the insect, 22—Natural enemies, 23—Geographical range, 23—Position of the moth when alighting, 24.	
	The canker-worm, Paleacrita vernata; Anisopteryx pometaria Dr. Le Baron's summary of remedies, 24—Mr. Milliken's experience with the rope and tin trap, 25—The Paris green remedy, 26—A new trough, 26—Birds which destroy the worm, 27—Mr. Mann's observations on the insect, 28—Two species have hitherto been confounded, 28—The English sparrow and the increase of the white-marked tussock-moth, 29.	24

1363. RILEY, C. V.—Continued.

INSECTS INJURIOUS TO THE GRAPE-VINE The grape Phylloxera, Phylloxera vastatrix..... The term "Phylloxera," 30-Bibliographical history, 30-Characters of the genus and its position in the system, 33-Biological history, 33-Different forms which the insect assumes, 33-The gall-inhabiting form, 34-The root-inhabiting form, 38-Balbiani's discovery of the true sexual individuals and the winter egg of Phylloxera querous, 41-The winter egg not essential to the winter life of the grape Phylloxera, 42-Polymorphism in Phylloxera, 43—Conclusive proof of the identity of the root- and galllice, 44-Practical considerations, 44-The more manifest and external effects of the disease, 44-Mode of spreading, 45-Swarming of winged specimens of Phylloxera caryæfoliæ, 45-Power of flight in the grape Phylloxera, 45—Spread of the disease through the winged females, 46-Susceptibility of different vines to the disease, 46-Practical importance of knowing the more resisting and more susceptible vines, 46-List of the cultivated species and varieties of vines indicating their relative susceptibility to Phylloxera, 47—Prophylactic means of coping with the disease, 48—Grafting the more susceptible on to the roots of resisting vines, 48-Other preventive measures, 50-Natural enemies, 50-Insect enemies of the gall-louse, 50—Enemies of the root-louse, 52-The Phylloxera mite, 52-Different forms assumed by mites in their development, 52-Hoplophora arctata, 53-Peculiarities of the Oribatida, 54-Direct remedies, 55-Submersion a perfect remedy, 55-Application of pure insecticides without satisfactory results, 56-Range of the insect in America, 57-Injury caused by

history, 64—Conclusion, 65.

Appendix and notes to the article on grape Phylloxera.....

Diagnosis of Phylloxera vastatrix, 66—Influence of the insect's

it in America, 58—Reasons why the insect is more injurious in Europe, 59—False theories, 60—The Phylloxera is the cause and not the effect of the disease, 60—It is a native American insect and has been imported into Europe, 62—It is peculiar to the grape-vine, 63—The American oak Phylloxera and its natural

puncture, 67—The supposed male of the gall-louse, 67—The non-cultivation of the Clinton grape, 67—Transient nature of the leaf-galls, 68—Constancy of the differences between the forms assumed by the insect, 68—Supposed sexual individuals, 68—Number of generations annually produced, 69—Number of molts, 69—Transplanting root-lice on to the leaves, 69—Nature of the swelling on the roots produced by Phylloxera, 70—The true grape-vines of the United States, by Dr. G. Engelmann, 70—Exceptional instances where the European vine has succeeded in America, 76—Grafting the more susceptible on to the roots of the resisting varieties, 78—Descriptions of Tyroglyphus phylloxera and Hoplophora arctata, 81—Efficacy of inundating the vineyards, 82—Facts showing that the disease of grape-vines in America is principally caused by Phylloxera, 82—Description of Phylloxera

- The grape-vine Epimenis, Psychomorpha epimenis, 87—Habits and characters of the larva, 87—Description of larva and chrysalis, 88.
- 2. The beautiful wood nymph, Eudryas grata, 83—Characters and food-plants of the larva, 88—The moth and the egg, 89—Description of the egg and larva, 89—Of the chrysalis, 90.

66

1363. RILEY, C. V.—Continued.	
INSECTS INJURIOUS TO THE GRAPE-VINE-Continued.	
3. The pearl wood nymph, Eudryge unio, 99—Its larva almost un-	
distinguishable rom that of the foregoing species, 96-Food-	
plant, 91—Distinguishing characters of the moth, 91—Mr. Lint-	
ner's description of the larva, 92—Mr. Lintner on the difference	
between the larva of Eudryae grata and E. unio, 93.	
 The eight-spotted ferester, Alypia octomaculata, 94—Description of the larva, 94; chrysalis and image, 95. 	.'
Summary, 95—Comparison of the larve of these four species, 95— Remedies 96.	
THE RED-LEGGED HAM-BEETLE, Neorodia ruspee	96
Its popular name, 96—It has been the cause of an interesting law-	
suit, 96; entomological information necessary to a just verdict,	
98-Another case before a jury requiring botanical knowledge,	
98-Injury caused by the beetle in St. Louis, 99-The eggs, and	
how they are deposited, 99—The larva and its habits, 99—The	
pupa, 99—Prevention, 100—Other species associated with it, 100—	
Habits of other species of Clerida, 101—How Necrobia ruficollis	
saved the life of Latreille, 101—Description of the larva of Ne-	
crobia rufipes, 101—Description of the pups, 102.	•
THE CLOVER-HAY WORM, Asopia costalis	102
Its geographical distribution, 102—It has probably been imported	
from Europe, 101—Its past history and accounts of damage	
caused by it, 102—Its natural history, 105—Remedies, 105—De-	•
scription of larva and chrysalis, 106; of the image, 107—Its al-	
lied congener, Asopia olinalis, 107. THE LEGGED MAPLE-BORER, Ægeria acerni	107
The burrows of the larva, 108—Appearance of the moth, 108—Re-	107
marks on the nomenclature of the species, 108—Injury caused	
by it to maples, 109—It attacks only trees already injured by	
some cause or another, 109—Remedies, 110—Description of imago,	
larva, and pupa, 110.	
THE RASPBERRY ROOT-BORER, Bembecia marginala	111
Work of Oberea perspicillata in cases of raspberry and blackberry,	
111-Injury caused by the root-bever, 111-General appearance	
and habits of the worm, 112—Description of image and larva,	
113.	
THE NORTHERN BRENTHIAN, Bupealls windle	113
Its occurrence and distribution, 113-Characteristics of the beetle,	
114—Fighting habits of the males, 114—How the male assists the	
female in ovipositing, 115—How much time is required for the	
transformations of the insect, 115—Habite of the larva, 115—De-	
scription of the larva, 115; of the pupa, 116—Real position of	
the brenthians in the system, 116—The specific and generic names, 116—How another larva has been mistuken for that of the	
Northern brenthian, 117—Description of this larva, which evi-	
dently belongs to the Tonobrionida, 118.	
THE JUMPING SUMACH-BRETLE, Blepharida rhois	118
Sumach industry in Europe and America, 118-Species of sumach	
possessing economic value, 119-Appearance and habits of the	
beetle, 119-The egg-masses, 120-Development and habits of	
the larva, 120—Transformation, 120—The two annual broods,	
120—Injury caused by it to the sumachs, 121—Geographical	
range, 121—Remedies, 121—Description of larva and pupa, 121—	
Variations of the beetle, 122.	

12 ENT

BENEFICIAL INSECTS.

THE UNADORNED TIPHIA OR WHITE GRUB PARASITE, Tiphia inornata ... 123 Other enemies of the white-grab, 123-Cocoon and larva of Tiphia, 123-It is undoubtedly a parasite of the white-grub, 124-Characters of the genus Tiphia, 124-Habits of Tiphia femorata, 124-Habits of the Tiphia larva, 124—Characters of the Unadorned Tiphia, 125-Beetle parasitic upon it, 125-Description of larva and imago, 126-The species has been described under three different names, 126. INNOXIOUS INSECTS. THE DOMINICAN CASE-BEARER, Coscinoptera dominicana..... 127 Characters of the beetle, 127-Mode of oviposition, 128-Oviposition of Chlamys plicata, 128-The young larva and its foodhabits, 129-The case of the larva, 129-Characters of the larva, 130-Time required for its development, 130. THE YUCCA MOTH, Pronuba yuccasella 131 Its natural history completed, 131-Description of the chrysalis, 131—Hatching of the chrysalis, 132—Method of oviposition, 133— Oviposition is followed by pollination, 134—The egg in the young fruit and the embryo larva, 134-It is the only insect that can well fertilize the Yuccas, 135. HACKBERRY BUTTERFLIES..... 136 The European purple Emperor, 136-Insufficient former account of their earlier states, 136-Species of Celtis in Missouri, 137. The eyed Emperor, Apatura celtis..... 137 The full-grown larva, 137—Habits of the larva, 138—Pupation, 138-The image, 139-The egg and the young larva, 139-Two annual broods, 139—Hibernation of the young larva, 140. The tawny Emperor, Apatura clyton..... 140 How it differs from the eyed emperor, 140-The egg, 141-Larval changes and habits, 141—Egg parasite, 142. Bibliographical 142 The generic name, 142-The specific names of the two species, 143-Fabricius's original description of lycaon and herse, 144-Other species of the genus in the United States, 145-On the validity of alicia, 145-Descriptions of the earlier states of both species, 146—Their popular names, 148—The scientific specific names, 149. KATYDID8 150 General considerations, 150—Spring in Europe and America, 151-Stridulating noise produced by crickets and grasshoppers, 152-Sounds inaudible to man, 152-Grasshoppers, katydids, locusts, 153—Habits and general appearance of our katydids, 154—They oviposit above ground, 154-Their ovipositors not rudimental, The angular-winged katydid, Microcentrum retinerve 155 It is the most common species in Missouri, 155—General character of the insect, 155—The eggs, 155—Erroneous statement concerning the egg, 156-How the female deposits the eggs, 156-Number of eggs laid by the female, 158—Hatching of the larva, 158— Food of the larva, 158-Changing from the pupa to the perfect

•	•	1
1363.	RILEY, C. V.—Continued.	
	state, 159—Its song, 159—It is capable of domestication to a cer-	
	tain degree, 160—Description of the immature states, 161—Nat-	
	ural enemics, 162—The back-rolling Wonder, an egg-parasite,	
	162—Curious habit of the female Antigaster, 162—Description of	
	Antigaster mirabilis, 163—Striking sexual differences, 163.	104
	The narrow-winged Katydid, Scudderis curvicanda	164
	egg, 165—Its song, 166—Description of larva and pape, 166.	
	The broad-winged Katydid, Cyrtophyllus concavus	167
	It is the true katydid, 167—Distinguishing characters, 167—Mode	
	of oviposition as observed in confinement, 167-Mr. Jaeger's	
	erroneous statement regarding the oviposition, 168-Its song,	
	168.	
	The oblong-winged katydid, Amblycorypha oblongifolia	169
	It has not yet been found in Misseuri, 169—Distinguishing charac- ters, 169—The eggs not yet known, 169—Abnormal female speci-	
	men, 169.	
1384	RILEY, C. V. The Colorado potato-beetle abroad. <n. td="" y.<=""><td>Tri-</td></n.>	Tri-
TOOT.	bune, 1 April, 1874. Sb. No. 10, p. 37.	T11-
	· · · · · · · · · · · · · · · · · · ·	
	Criticism of some articles on <i>Doryphora decemlineata</i> published in Europe, and suggestion of the insect into Europe, and suggestion	
	genger or the importantion or the intect into transfer and suggested	T OF

RILEY, C. V. Cabbage-lice. <N. Y. Tribune, 8 April, 1874.
 S.-b. No. 10, pp. 33-34.

means to prevent such introduction.

- Answer to letter from Mrs. M. Walker; means against Aphie brassion.
- 1366. RILEY, C. V. Peach-borers. <N. Y. Tribune, 8 April, 1874. S.-b. No. 10, p. 34.
 - Answer to letter from W. J. Clary; a peach-tree whose base was surrounded by ice in winter was free from borers; this observation of slight significance.
- 1367. RILEY, C. V. Apply soap. <N. Y. Tribune, 8 April, 1874. S.-b. No. 10, p. 34.
 - Answer to letter from T. H. Wakeley; means against Saperda bivittata [=can-dida] and Chrysobothris femorata.
- 1368. RILEY, O. V. Meadow enemy. <N. Y. Tribune, 8 April, 1874. S.-b. No. 10, p. 34.
 - Answer to letter from D. Freeman; ravages of a species of Tipula in grass lands in California; characters of the genus; habits of and means against the same.
- 1369. RILEY, C. V. A remedy for the cotton-worm. <N. Y. Tribune, 22 April, 1874. S.-b. No. 10, pp. 25-26. Reprint: <Vicksburg Herald, 1 May, 1874.
 - Experiments with Paris green; advocacy of its use against Aletta argillaces [=xylina]; directions for and successful results of this use.
- 1370. RILEY, C. V. Descriptions of two new subterranean mites. <Trans. Acad. Sci. St. Louis, 25 April, 1874, v. 3, pp. 215-216, figs. 8-9.
 - Description and figures of Tyroglyphus phylloxera n. sp. [p. 215] and of Hoplophora arotata n. sp. [p. 216]; food-habits of the same,

- 1371. RELEY, C. V. The habits of Polistes and Pelopœus. < Amer. Nat., April, 1874, v. 8, pp. 229-231.
 - Critical review of P. R. Uhler's article "On a remarkable wasp's nest found in a stump in Maryland;" probably the nests and specimens of Pelopous lunatus [= comentarius] were mistaken for those of Polistes fuscatus [= metricus], in which case the observations criticised present no remarkable features.
- 1372. RILEY, O. V. The plum Curculio; natural history and how to catch him. <N. Y. Semi-weekly Tribune, 1 May, 1874. S.-b. No. 10, pp. 18-19; 38-40. Extract: <Cultivator and Country Gentleman, 14 May, 1874, v. 39, p. 310. <New England Farmer, 18 July, 1874, [v. 53,] n. s., v. 29, p. 1. See: <Indust. Record, 5 February, 1875. S.-b. No. 16, pp. 46-48.
 - Natural history of and means against Constructelus nonupher; figures of the several stages of the insect and of machines for catching the same.
- 1373. RILEY, C. V. The apple-worm; natural history; remedies. <N. Y. Tribune, 20 May, 1874. S.-b. No. 10, pp. 8-9; 37-38. Description and figures of the several stages of Carpocapes pomenella; habits of and means against the same.
- 1374. RILEY, C. V. The grape Phylloxera. <Popular Sci. Monthly, May, 1874, v. 5, pp. 1-16, 7 figs. Reprinted, with slight changes, from <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 30-46.

 See No. 1363 for synopsis of contents.
- 1375. RILEY, C. V. Rose chafers on grape-vines. <Colman's Rural World, 20 June, 1874, fig. S.-b. No. 17, pp. 103, 104.

 Natural history and means against Macrodactylus subspinosus; figure of the same.
- 1376. RILEY, C. V. More about the grape-vine pest. <Popular Sci. Monthly, June, 1874, v. 5, pp. 158-170, 10 figs. Reprinted, with changes and omissions, from <6th Ann. Rept. State Ent. Mo., April, 1874, pp. 47-60, 64-65.

 See No. 1363 for synopsis of contents.
- 1377. RILEY, C. V. Scale insects on magnolia. <N. Y. Tribune, 15
 July, 1874. S.-b. No. 10, p. 24.
 Answer to inquiry of R. H. Day; occurrence of an undescribed Lecanium sp.?

on the leaves of Magnolia fuscata.

- 1378. BILEY, C. V. Confounding friend with foe. <N. Y. Tribune, 15 July, 1874, 2 figs. S.-b. No. 10, p. 24.</p>
 Answer to inquiry of E. J. Day; usefulness of Arms [= Podieus] spinosus
 - Answer to inquiry of E. J. Day; usefulness of Arma [= Podious] spinosus figure of the same. Injury to plums by Anthonomus prunicida [= Coccotorus scutellaris]; figure of and means against the same.
- 1379. RILEY, C. V. The Colorado potato-beetle in New York. <N. Y. Tribune, 15 July, 1874. S.-b. No. 10, p. 24.

 Answer to letter from J. H. P.; eastern limit of the range of Deryphora dependineats in 1873 and 1874.

- 1380. RILEY, C. V. Large willow-worm. <N. Y. Tribune, 15 July, `1874. S.-b. No. 10, p. 24.
 - Answer to letter of J. H. P.; food-plants, habits, and means against Cimber laportei [= americana].
- 1281. RILEY, C. V. Black blister beetles on potatoes. <N.Y. Tribune, 22 July, 1874. S.-b. No. 10, p. 17.
 - Answer to inquiry of A. Barr; ravages of and means against Epicauta puncti-
- 1382. RILEY, C. V. Pear-tree slug. <N. Y. Tribune, 22 July, 1874. S.-b. No. 10, p. 17.
 - Answer to letter of C. S. B.; means against Sclandria [= Ericcampa] cerasi.
- 1383. RILEY, C. V. The plug-ugly theory. <N. Y. Tribune, 22 July, 1874. S.-b. No. 10, p. 17.
 - Answer to letter of O. J. B.; insertion of sulphur, calomel, and similar powders in the trunks of trees has no effect upon the sap or upon the insects which injure the trees.
- 1384. RILEY, C. V. Cockscomb elm-gall. <N. Y. Tribune, 22 July, 1874. S.-b. No. 10, p. 17.
 - Answer to inquiry of J. S. Ruby; Thelazes [= Colopha] ulmicola forms excrescences on the American elm.
- 1385. RILEY, C. V. Pitcher-plant insects. <Hartford [Conn.] Daily Courant, 15 August, 1874, v. 38, No. 195, p. 1. See: <N. Y. Tribune Extra: Lecture and Letter series No. 21, August, 1874, pp. 56-58, fig. Reprint: <Nature, 8 October, 1874, v. 10, pp. 463-465, figs. 1-2. Abstract: <Sci. Amer., 12 September, 1874 [v. 45], n. s., v. 31, p. 168. <Amer. Nat., November, 1874, v. 8, pp. 684-687. Reprint of abstract: <Hardwicke's Science Gossip, December, 1874, v. 10, pp. 272-275, figs. 179-182. Extract: <Ca. Ent., November, 1874, v. 6, pp. 207-214, figs. 25-26. Reprint of extract: <Proc. Amer. Assoc. Adv. Sci. for 1874, June, 1875, v. 23, pp. 18-25, 2 figs. Separate of reprint: <Salem, Mass., December, 1874, pp. 18-25, figs.
 - Insect-catching habits of Sarracenia variolaris; list of its victims; habits and figures of all stages of Xanthoptera [= Exyra] semicrocea and Sarcophaga sarracenia; capture of insects by other plants.
- 1386. RILEY, C. V. On the habits and transformations of Canthon hudsonias, Forst.; the common "tumbledung." <Hartford [Conn.] Daily Courant, 18 August, 1874, No. 11162, v. 38, No. 197, p. 2. Reprint: <N. Y. Tribune Extra: Lecture and Letter series, No. 21, August, 1874, pp. 75–76.
 - Breeding habits and oviposition of Canthon hudsonias [= lavis].
- 1387. RILEY, C. V. On the larval habits of the cantharid genera Epicauta and Henous. <Hartford [Conn.] Daily Courant, 18 August, 1874, No. 11162, v. 38, No. 197, p. 2. See: <N. Y. Tribune Extra: Lecture and Letter series, No. 21, August, 1874, p. 76.
 - Habits and hypermetamorphosis of *Melos angusticollis*; probably parasitic food-habits of young larve of *Epioguta* and *Henous*; the images phytophagons,

1388. RILEY, C. V. Humming-bird moths caught by the tongue.

<Moore's Rural New Yorker, 29 August, 1874, v. 30, p. 140.

Method of capture of insects, particularly moths by the flowers of Physican-

thus albens; capture of insects by other flowers.

1389. RILEY, C.V. The io moth (Saturnia io). < Illust. Journ. Agric., August, 1874, figs.

Answer to inquiry of G. Barter; natural history of Saturnia [= Hyperchiria]

io; figures larva and & and Q imagos.

- 1390. RILEY, C. V. Descriptions and natural history of two insects which brave the dangers of *Sarracenia variolaris*. <Trans. Acad. Sci. St. Louis, 1 September, 1874, v. 3, pp. 235-240, figs. 10-11.
 - Description of leaf of Sarracenia variolaris and its function of entrapping and digesting insects; description and figures of eggs, larva, chrysalis, and image of Santhoptera [=: Exyra] semi-crocea and larva, puparium and image of Sarcophaga sarracenia n. sp.; habits of the two species; comparison of S. sarracenia with S. carnaria and of the genera Musca, Calliphora, and Sarcophaga; notice of A. S. Packard's "Transformation of the common housefly."
- 1391. [RILEY, C. V.] [Discussion on entomology.] <Trans. Ill. State
 Hortic. Soc. for 1873, 1874, n. s., v. 7, pp. 100-104.
 Description, habits, and means against Anisopteryx; effect of late plowing;
 - enemies of Doryphora 10-lineata; habits of Gastrophilus equi.
- 1392. [RILEY, C. V.] Note on leaf-hopper. <Trans. Ill. State Hortic. Soc. for 1873, 1874, n. s., v. 7, p. 138.

 Means against Erythroneura [= Typhlocyba] vitis.
- 1393. [RILEY, C. V.] [Notes on the strawberry crown borer.] < Trans.

 Ill. State Hortic. Soc. for 1873, 1874, n. s., v. 7, p. 147.

 Description, habits, and means against Tyloderma fragariæ.
- 1394. RILEY, C. V. Lecture on entomology. <Trans. Ill. Hortic. Soc. for 1873, 1874, n. s., v. 7, pp. 172–178, figs. 1–3. Reprint: <3d Ann. Rept. Sec. State Pomol. Soc. Mich. for 1873, 1874, pp. 443–448. Abstract: <Rept. U. S. Commis. Agric. for 1873, 1874, [30 March, 1875], pp. 389–390.
 - Habits of and means against Carpocapsa pomonella; habits, transformations, prolificacy of and means against Phylloxera vastatrix; description and figures of leaf-galls; figures of root- and gall-forms; history and meaning of the word "Phylloxera;" statement by D. B. Wier concerning the invention and use of Wier's apple-worm trap.
- 1395. RILEY, C. V. "Walking-sticks or specters" becoming injurious. <N. Y. Weekly Tribune, 11 November, 1874, fig. S.-b. No. 23, p. 103.
 - Ravages in Yates County, N. Y., vernacular names, habits of, and means against Diapheromera femorata; description of its eggs; figure of image.
- 1396. RILEY, C. V. Entomological notes. <Sci. Amer., 5 December, 1874 [v. 45], n. s., v. 31, p. 356.
 - Pemphigus imbricator found on beech; vesicatory potato-beetles, Meloidæ; cow manure and cow urine one of the earliest supposed remedies for Phylloxera.

- 1397. RILEY, C. V. The bark louse. <Beach, A. E. The Science Record for 1874, N. Y., 1874, p. 356.

 Discovery of the male of Mytilaspis conchifermis [= pomorum].
- 1398. RILEY, C. V. Codling-moth heresies. <N. Y. Tribune, 2 January, 1875. S.-b. No. 13, p. 163.</p>
 Criticism of paper of S. B. Peck; Carpocapsa pomonella confines its attacks to one apple and does not enter the ground.
- 1399. RILEY, C. V. Shall we scrape our trees? <N. Y. Tribune, 6
 February, 1875. S.-b. No. 10, p. 10.
 Careful scraping of trees early in spring is beneficial as a safeguard against
 the attacks of many injurious insects.
- 1400. RILEY, C. V. What are army-worms? <N. Y. semi-weekly Tribune, 6 February, 1875.

 Differences between Laphygma frugiperda and Loucania unipuncta.
- 1401. RILEY, C. V. The hickory bark-borer, Scolytus carye. <Colman's Rural World, 6 February, 1875. S.-b. No. 17, p. 105.

 Natural history of Scolytus caryæ [= 4-epinosus].
- 1402. RILEY, C. V. Genuine vs. bogus chinch-bugs. <N. Y. Tribune, 10 February, 1875, figs. S.-b. No. 10, p. 10. Answer to inquiry of C. H. Cushing; food-babits and figures of Nysius destructor [= angustatus] and Blissus leucopterus; larva of Deilephila lineata feeds on purslane.
- 1403. RILEY, C. V. Remedies for Phylloxera. <N. Y. Tribune, 10
 February, 1875. S.-b. No. 10, p. 22.
 - At present no insecticide is effective against Phylloxera vastatrix; grafting of more susceptible varieties on the roots of the least susceptible advised.
- 1404. RILEY, C. V. Newest facts of grape Phylloxera. <N. Y. Tribune, 10 February, 1875. S. b. No. 10, pp. 21-22. Succession of different forms of individuals in the cycle of development of Phylloxera described; seasons at which the different forms appear; places

in which eggs are laid by the winged females.

- 1405. RILEY, C. V. Is the Colorado beetle poisonous? <N.Y. Weekly Tribune, 17 February, 1875. S. b. No. 10, p. 9.</p>
 Insists upon the poisonous nature of the fumes from scalded or burning masses of Doryphora decomlineata, in opposition to statements by Prof. T. J. Burrill.
- 1406. RILEY, C. V. Notes of Phylloxera. <N. Y. Tribune, 4 March, 1875. S.-b. No. 10, pp. 20-21.</p>
 Critical review of A. S. Fuller's "Distribution of the grape-losse;" maintains the specific identity of the so-called gall-inhabiting and root-inhab-

iting forms of Phylloxera, and the North American origin of this insect.

1407. [RILEY, C. V.] The Colorado potato-beetle abroad. <N. Y. Weekly Tribune, 17 March, 1875. S.-b. No. 10, pp. 14-15. Statement and criticism of measures adopted by several European Governments to prevent the introduction of Doryphora decembranes into their countries.

- 1408. RILEY, C. V. Bud-eating insects. <Cultivator and Country Gentleman, 25 March, 1875, v. 40, p. 183.

 Habita food-plants and manne against Agrotic condens: means against field.
 - Habits, food-plants, and means against Agrotis sounders; means against field cut-worms.
- 1409. RILEY, C. V. Description of a new species of Agrotis. < Proc. Bost. Soc. Nat. Hist., March, 1875, v. 17, pp. 286–288. Description of Agrotis morrisoniana n. sp. [p. 286].
- 1410. RILEY, C. V. On the summer dormancy of the larva of *Phyciodes***myctois*, Doubleday, with remarks on the natural history of the species. <Proc. Amer. Assoc. Adv. Sci. for 1874, 4 June, 1875, v. 23, pp. 108-112. Separate: <Salem, Mass., March, 1875, pp. 108-112. Abstract: <Hartford [Conn.] Daily Courant, 18 August, 1874, No. 11162, v. 38, No. 197, p. 2. See: <N. Y. Tribune Extra: Lecture and Letter series, No. 21, August, 1874, p. 75.
 - Observations on *Phyciodes nycteis* and *Argynnis bellona*; significance of the same; detailed description of the larva and pupa of *P. nycteis*.
- 1411. RILEY, C. V. Descriptions of two new moths. <Trans. Acad. Sci. St. Louis, 1 February [March], 1875, v. 3, pp. 240-242, figs. 12-13.
 - Descriptions and figures of Xanthoptera [== Exyra] ridingsii n. sp. and Cerura multiscripta n. sp.
- 1412. RILEY, C. V. The climate for Doryphora. <N. Y. Tribune, 2
 April, 1875. S.-b. No. 10, p. 16.
 - Comments on letter from D. L. Garver; *Doryphora decemlineata* would be likely to thrive as well in the climates of Europe as in North America; its transportation across the ocean is not improbable.
- 1413. RILEY, C. V. [On an Acridium eaten out by ants.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 102 Proc. Verbal communication; inner soft parts of an Acridium americanum eaten out by Myrmica minuta.
- 1414. RILEY, C. V. [On the chrysalis of *Pronuba yuccasella*.] <Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 106 Proc. Verbal communication; adaptation of pupa of *Pronuba yuccasella* to its needs of prying its ways through the soil.
- 1415. RILEY, C. V. [On regulating sex in insects.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 108 Proc.

 Verbal communication; critical review of Mrs. Treat's "Controlling sex in butterflies"; female insects need more neurishment than the males, but sex is determined in all animals at conception.
- 1416. RILEY, C.V. [On the peculiarities of Nephila plumipes.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 109 Proc.

 Verbal communication; comparison of the structure of male and female Nephila plumipes; habits and silk-production of the female.
- 1417. RILEY, C. V. [On the peculiarities of the Mexican honey-ant.]

 Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 109 Proc.
 Verbal communication; structural and functional division of the neuters of Myrmecocystus mexicanus into two distinct kinds.

- 1418. RILEY, C. V. [On the peculiarities of *Physianthus albens.*] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 109 Proc.
 - Verbal communication; the flowers of *Physianthus albens* so constructed as to hold large *Sphingida* fast by the tongue.
- - Verbal communication; method of capture of insects, particularly moths, by the flowers of *Physianthus albens*; records the capture of a number of *Noctuidæ* and of *Sphingidæ*, especially *Deilephila lineata*; *Norium oleander* and *Enothera grandiflora* are said to capture Sphinx-moths in Europe.
- 1420. RILEY, C. V. [On the Yucca borer.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 139 Proc.
 - Verbal communication; entomological interest attaching to the Yucca; abode and synonymy of Megathymus yucca.
- 1421. RILEY, C. V. [New biological facts regarding the grape Phylloxera.] < Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, pp. 147–148 Proc.
 - Sequence of development of certain forms of *Phylloxera*; sexual forms of three species of *Phylloxera* obtained; alleged discovery by J. Lichtenstein of the winged form of *P. vastatrix* on *Quercus coccifera* in Europe, discredited, and this form considered by E. G. Balbiani as a new species, named *P. lichtensteinii*.
- 1422. RILEY, C. V. [On the connection of locust invasions with the occurrence of drought.] <Trans. Acad. Sci. St. Louis, 6 April, 1875, v. 3, p. 163 Proc.
 - Verbal communication; no connection between the occurrence of drought and of locust invasions.
- 1423. RILEY, C. V. Seventh annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture pursuant to an appropriation for this purpose from the legislature of the State. <10th Ann. Rept. State Board of Agric. for 1874, April, 1875, pp. 7 + 196 + 4, 40 figs. Separate: <Jefferson City, Mo., April, 1875, pp. 7 + 196 + 4, 40 figs.

PREFACE III
TABLE OF CONTENTS VII

1

NOXIOUS INSECTS.

51

71

80

90

1423. RILEY, C. V.—Continued.

THE COLORADO POTATO-BEETLE-Continued.

man indirectly through the soil or through the plant, 13-The beetle eats as well as the larva, 14-It passes the winter as imago, 14-New food-plants, 14-New means of destruction, 15-The Gray sprinkler, 15-The proper scientific name of the beetle, 16-Mr. Carrière's ridiculous statements, 17.

THE CHINCH-BUG, Blissus leucopterus.....

Its disastrous work in 1874, 19-Circular distributed among farmers, 19-Appearance and transformations of the chinch-bug, 20-The short-winged form, 20-Description of the chinch-bag and its earlier states, 21-Its past history in America 22-Its past history in Missouri, 22-Destructive powers of the chinch-bug, 24—Its injuries in 1874, 24—Its injuries in Missouri in 1874, 25— Food-plants, 26-Time required for the commete development of various insects, 27-Number of annual broods, 27-Its rapid increase, 28-Where the eggs are laid, 28-Flight of the chinchbug, 29—Its migrations on foot, 30—Heavy rains destructive to it, 30-Direct remedies, 31-Irrigation, 31-Preventive measures, 32-Burning, 32-Rolling, 33-Manuring and early sowing, 34-Mixing seed, 34—Preventing the migrati n of the bugs from one fie'd to another, 35-Importance of winter work and combined action, 36-Other possible remedies, 37-Abstaining from the cultivation of grains, 38-Natural enemies, 38; lady-birds, 39; lace-wing fly and habits of its larva, 40; the insidious flower-bug and the many-banded robber, 41-Birds destructive to the chinchbug, 41-Discussion of other proposed remedies and preventive measures, 41-The chinch-bug injurious to stock, 43-Prognosticating, 44-Unnecessary lears, 44-Bogus chinch-bugs, 45-The false chinch-bug, 46-The insidious flower-bug and the ash-gray leaf-bug, 47-The flea-like negro-bug, 48-Recapitulation, 49.

Appendix to the article on the chinch-bug List of correspondents who replied to the circular, 51-Questions answered by correspondents, 52-Answers given by correspondents, 53.

THE FLAT-HEADED APPLE-TREE BORER, Chrysobothris femorata Extent of its ravages in fruit and shade trees, 71-Its natural history, 72-Natural enemies, 73-Chalcid larva, 74-The cherished Bracon, 75—The useful Labena, 75—Ants, 76—Remedies, 76— Keeping young trees vigorous and healthy, 77-Coating the trunks and larger branches with soap and other greasy substances, 77-Scraping, 78-Wrapping wire gauze around the trunk, 79-Cutting out the newly-hatched larva, 79.

CANKER-WORMS, Paleacrita vernata; Anisopteryx pometaria..... Confusion regarding the two species of canker-worms, 80-Distinguishing characters of the spring canker-worm, 80-Description of egg, larva, and chrysalis of Anisopteryx vernata, 82-How the fall canker-worm differs from the preceding, 83-Description of egg, larva, and chrysalis of Anisopteryx pometaria, 84-Practical importance of distinguishing the two species, 85-Comparative description of the spring and fall canker-worms, 86-Conclusion, 88-Extracts from the original essay on the canker-worm by W. D. Peck, 89.

THE GRAPE PHYLLOXERA, Phylloxera vastatrix..... Completion of its natural history, 90-The true sexual individuals discovered, 91-Epitome of the life-history of the grape Phyllox-

. THE GRAPE PHYLLOXERA-Continued.

era, 91—Different forms presented by the species, 93—Its power to change its habit, 93-Specific identity of the gall- and rootlouse, 94—Untrustworthy experiments made by the Department of Agriculture, 95-Proof of the identity of the two forms, 95-The gall-louse is but a transient form, 96-Where do the winged females lay their eggs 196-The winged female lays the egg wherever she is carried by the wind, 97-Particular part of the vine chosen by the winged female for laying her eggs, 98-The true sexual individuals, 98—Injury done by Phylloxera in America. during the year 1874, 99-Range of the insect in America, 101-Does it occur in South Carolina and Georgia ! 102—The Phylloxera in California, 103-Injury done during the year 1874 in France, 103-Its spread in Europe, 104-Direct remedies, 105-Natural enemies, 106-Susceptibility of different varieties of grape-vine, 106-Grafting as a means of counteracting the work of Phylloxera, 108-Underground grafting, 109-Methods of grafting above ground, 112-Roots to use as stock, 115-Varieties to graft, 116-American grape-vines abroad, 116.

THE ROCKY MOUNTAIN LOCUST, Caloptenus spretus.....

Its natural history, 121-Method of egg-laying, 121-The egg, 122-The newly-hatched locust and its development, 122-Where the eggs are laid by preference, 123-The invading swarms are formed by a single species, 124-Difference between the Rocky Mountain and the red-legged locusts, 125-Mr. Thomas's description of the red-legged locust, 126-Variations, 126-Measurements of Caloptenue femur-rubrum, 127-Description of Caloptenus spreius and of its larva and pupa, 129-Measurements of Caloptonus spretus, 130-Summing up the difference between the two species, 132-Chronological history, 132-Locust invasions in the Old World, 132-Accounts of earlier locust invasions in America, 133—Chronological history of the Rocky Mountain locust, 134—Earlier invasions, 135—Data regarding the invasion of 1867, 137—The invasion of 1873, 141—The invasion of 1874, 143—The invasion of 1874 in Missouri, 144—Questions addressed to correspondents in each county in Missouri regarding the locust, 144-Summary of the answers given by correspondents, 145-The invasion of 1874 in Kansas, 148—In Nebraska, 151—In Iowa and Minnesota, 153— In Colorado, 154-In Dakota and Manitoba, 155-Flight and ravages of a locust swarm, 156-Food-plants, 158-Time of appearance of invading swarms, 160—Eastern limits of locust invasions, 161-Native home of the species, 162-Explanation of the migratory instinct, 164—This locust can not thrive in the Mississippi Valley, 164-It is an subalpine insect, 165-What injury may be expected in Missouri in 1875, 166—Ravages of migratory locusts in the Atlantic States, 167—Description of the Atlantic migratory locust, 169-Differences between Caloptenus spretus, differentialis, and atlanis, 170-Injury from other, non-migratory, locusts, 171—The differential and two-striped locusts, 173—Enemies and parasites, 174—Birds destroying locusts or their eggs, 174—The 117

1423. BILEY, C. V.—Continued.

THE ROCKY MOUNTAIN LOCUST-Continued.

silky mite, 175—The locust mite and Dr. Le Baron's description thereof, 176—The anonymous Tachina-fly, 178—The common flesh-fly, 180—Remedies, 181—Classification of protective measures, 181—Natural agencies, 181—Destruction of the eggs, 181—Destruction of the young wingless locusts, 182—Driving off the winged locusts, 184—Further suggestions, 185—Locusts as food for man, 186—The popular and scientific names of the insect, 187—Prairie fires vs. the Rocky Mountain locust, 189.

Appendix to the article on the Rocky Mountain locust......

Letters of three correspondents from Texas and Kansas regarding locust invasions, 191.

1424. RILEY, C. V. The ways of bag-worms. <N. Y. Semi-weekly Tribune, 14 April, 1875, 3 figs. S.-b. No. 10, pp. 11, 36.

Figure of the larva-case of an undetermined species of Psychide and of the several stages of Thyridopteryx sphemoresformis; habits of and means against the latter.

- 1425. [RILEY, C. V.] [Poisonous qualities of the Colorado potato-beetle.] <N. Y. Tribune, 14 April, 1875. S.-b. No. 10, p. 17, 34.

 Mentions instances of poisoning from the fumes given out by scalded or crushed masses of Doryphora decembineata.
- 1426. [RILEY, C. V.] Cure for eanker-worm. <N. Y. Tribune, 21 April, 1875. S.-b. No. 10, p. 25.</p>
 Paris green recommended for destruction of canker-worms when other rem-
- edies are lacking. Paris green not to be used against all insects.

 1427. [RILEY, C. V.] Paris green: Its effects on plants and soils, and through them on man. <N. Y. Tribune, 12 May, 1875. S.-b. No. 10, pp. 15–16; 34–35.

Paris green is not injurious to plants or the soil, nor dangerous to man when properly applied.

- 1428. [RILEY, C. V.] Swellings on roots of Ampelopsis. <Colman's Rural World, 5 June, 1875. S.-b. No. 10, p. 1.

 Answer to letter of H. Hilker; swellings on roots of Ampelopsis possibly occasioned by root-lice, but not by Phyllozera.
- 1429. RILEY, C. V. Apple-tree plast-lice. <Colman's Rural World, 5 June, 1875.

Answer to letter of A. Moyer; occurrence of and means against Aphie mali.

1430. RILEY, C. V. [Torrubia elongata, the white-grub fungus.] <Colman's Rural World, 12 June, 1875, v—, p—.

Torrubia elongata proposed as the name for the white-grub fungus; figures of the same.

1431. RILEY, C. V. The Colorado potato-beetle, *Doryphora decemlineata*. <Garden [London], 24 July, 1875, v. 8, pp. 71-72, 5 figs. S.-b. No. 23, p. —.

Description and figures of all stages of Doryphora decembineats and D. juneta; migrations, habits, prolificacy, enemies, and parasites of D. decembineata; means against it; the use of Paris green; probability and probable method of its introduction into Europe and means of preventing such introduction; figures Lydella [= Exorista] doryphora, Harpactor [= Milyas] cinctus, Arma [= Podisus] spinosus, and Hippodamia convergens.

- 1432. RILEY, C. V. Locusts vs. chinch-bugs. <N. Y. Tribune, 4 August, 1875. S.-b. No. 10, p. 40.
 - Locust ravages are likely to prevent serious injury by *Blissus leucopterus*; the latter more noticeable than usual; excessive rains in 1875, also unfavorable to the development of *Blissus leucopterus*.
- 1433. RILEY, C. V. No locust injury in Kansas and Missouri this fall.
 <N. Y. Tribune, 1 September, 1875. S.-b. No. 10, p. 166.</p>
 Predicts immunity from attacks of Caloptenus spretus in the fall of 1875 in Kansas and Missouri.
- 1434. RILEY, C. V. Prof. Riley and the locusts. <St. Louis Daily Globe Democrat, 4 September, 1875, v. 1, No. 108, p. 3. S.-b. No. 10, pp. 158-159.
 - Reply to C. A. Davis's "Prof. Riley and the locusts;" food-plants, habits, parasites, and period of development of Caloptenus spretus; need of more extended investigation of this insect in its native haunts and breeding places.
- 1435. RILEY, C. V. The grape-leaf gall. <Cultivator and Country Gentleman, 9 September, 1875, v. 40, p. 567.

 Identity of the root and leaf-forms of Phylloxera rastatrix.
- 1436. RILEY, C. V. White grub fungus. <N. Y. Tribune, 6 October, 1875. S.-b. No. 10, p. 26.
 - Answer to inquiry of A. C. G.; brief history of Torrubia elongata, a parasite of Lachnosterna fusca.
- 1437. RILEY, C. V. Flying locusts in Illinois. <Cultivator and Country Gentleman, 28 October, 1875, v. 40, p. 679.

 Caloptenus spretus confounded with C. femur-rubrum by B. F. J.; C. spretus does not occur in Illinois.
- 1438. RILEY, C. V. Remarks on canker worms and description of a new genus of *Phalænidæ*. <Trans. Acad. Sci. St. Louis, 5 November, 1875, v. 3, pp. 273–280, figs. 14–21. Separate: <St. Louis, Mo., 1875, pp. 8, figs. Reprint: <8th Ann. Rept. State Ent. Mo., May, 1876, pp. 12–18, figs. 3–10. Notice: <Ca. Ent., November, 1875, v. 7, p. 219. <Proc. Bost. Soc. Nat. Hist., February; 1876, v. 18, p. 201.
 - Treats of Paleacrita [n. g.] vernata and Anisopteryx pometaria; see No. 1482 for synopsis of contents.
- 1439. RILEY, C. V. Notes on the natural history of the grape Phylloxera, *Phylloxera vastatrix*, Planchon. <Trans. Acad. Sci. St. Louis, 5 November, 1815, v. 3, pp. 281-287, fig. 22. Extract: <Amer. Nat., June, 1881, v. 15, pp. 483-484.
 - Discovery of the nidus in which eggs are deposited by the winged female of *Phylloxera vastatrix*; development, habits, description, and figures of the same; description of the male and of the impregnated egg; figure of the male of *P. carywcaulis*; summary of the natural history of *P. vastatrix*.
- 1440. [RILEY, C. V.] Grubs and guess-work. <N. Y. Semi-weekly Tribune, 12 November, 1875. S.-b. No. 10, pp. 17-18.
 - Characterizes some subdivisions of the Lamellicornia by their habits; contrasts the larvæ, imagos, and habits of Lachnosterna quercina [=fusca] and Lygyrua relictus; means against the former.

- 1441. R[ILEY], C. V. Entomological. Apple-tree borers; timber encourages them; new bag-worm. <Colman's Rural World, 13 November, 1875. S. b. No. 10, p. 188.
 - Timber-trees near apple orchards increase the likelihood of the attacks of Saperda bivittata [= candida] upon the apple-trees; habits and food-plants of Psyche confederata.
- 1442. RILEY, C. V. The army-worm; an important point yet to ascertain in its history. How it comes and goes; its natural enemies; preventive measures. <N. Y. Tribune, 16 November, 1875, 8 figs. S.-b. No. 10, pp. 29-31; 31-33.
 - Various applications of the name "army-worm;" seasons, habits, and natural enemies of and means against Leucania unipuncta; supposed method of oviposition; figures of it in its several stages, of Exorista flavicauda, Microgaster [= Apanteles] militaris, Ophion purgatum, and Pezomachus minimus, and of cocoons of the last; habits of these parasites.
- 1443. RILEY, C. V. The flying locusts in Illinois. <Cultivator and Country Gentleman, 25 November, 1875, v. 40, p. 744.

 Extent of swarms flying south over Kansas and Nebraska on September (4th?).
- 1444. R[ILEY], C. V. Scabby potatoes. <N. Y. Tribune, 15 December, 1875. S.-b. No. 10, pp. 20, 29.</p>
 The scab in potatoes is caused by imperfectly studied Acarina.
- 1445. [RILEY, C. V.] Not the Hessian-fly. <N. Y. Tribune, 15 December, 1875. S.-b. No. 10, pp. 20, 29.

 Answer to inquiry of J. H. K.; significance of the name and habits of Aphodius inquinatus.
- 1446. R[ILEY], C. V. How to destroy locusts. <Colman's Rural World, 23 December, 1875. S.-b. No. 10, p. 185.

 Answer to letter of C. Herschel; means against Acrididæ; recommends flooding and ditching.
- 1447. BILEY, C. V. Paris green as an insect destroyer. <N. Y. Tribune, 28 December, 1875. S.-b. No. 10, pp. 13-14; 28-29.</p>
 Paris green as a means against Doryphora decembineata; R. C. Kedzie's investigations show that Paris green is not deleterious to the soil or the crops.
- 1448. BILEY, C. V. Oak apple. <Amer. Cyclopædia, 1875, v. 12, pp. 558-559, 3 figs. S.-b. No. 14, pp. 47-48.</p>
 Description of galls of Cynips terminalis, C. [= Amphibolips] querous-sponsifica, and C. [=A.] q.-inanis; manner of their formation; figures the latter two galls and a parent fly; nature of galls and problems involved in their study.
- 1449. [RILEY, C. V.] Phylloxera. <Amer. Cyclopædia, 1875, v. 13, pp. 477-480, figs. 1-8. S.-b. No. 10, pp. 84-87.

 Habita, food-plants, and characters of the genus Phylloxera; natural history
 - of P. vastatrix; figures its several forms and the galls formed by it; history of the ravages in the vineyards of France.

- 1450. [RILEY, C. V.] Potato-bug. <Amer. Oyclopædia, 1875, v. 13, pp. 768-771, fig. 1-7. S.-b. No. 10, pp. 89-92.
 - Food-plants, original home, geographical distribution, natural history, enemies of, and means against Doryphora decominests; figures of it and of Lydella [= Exorista] doryphora, Arma [= Podisus] spinesus, Harpactor [= Milyas] cinclus, Mysia [= Anatis] 15-punctets, and Doryphora juncta.
- 1451. RILEY, C. V. Rocky Mountain locust. < Amer. Cyclopædia, 1875, v. 14, pp. 371-374, figs. 1-10.
 - Ravages, description, natural history, migrations, and enemies of and means against Caloptenus spretus; figures of it with details of its structure and habits; figures C. femur-rubrum, Astoma gryllarium [= Trombidium locustarum], Trombidium seriosum, and Sarcophage carnaria.
- 1452. RILEY, O. V. The hateful or Rocky Mountain locust, Caloptenus spretus. <N. Y. Tribune, 1875. Reprint: <Trans. Kans. State Hortic. Soc. for 1874, 1875, v. 4, pp. 172–176.
 - Answer to inquiry of Z. F. Hopkins; probable limit of the ravages and range of Caloptenus spretus in Kansas, Missouri, and neighboring States in 1875; means against and vernacular name of this species; vernacular names of Acrididæ and Locustidæ in general.
- 1453. RILEY, C. V. Prairie fires and hateful locusts: is there any connection between them? <N. Y. Tribune, 1875. Reprint: <Trans. Kans. State Hortic. Soc. for 1874-75, v. 4, pp. 176-180. Criticism of the effects of the drought, hot winds, locusts, and short crope in Kansas caused by the burning of the prairie grasses, as stated in Kansas Farmer, 23 September, 1874; benefits resulting from prairie fires by the destruction of hibernating Blissus lowopterus, Nysius destructor [== angustatus], etc.; relations of prairie fires to the origin and maintenance of prairies.
- 1454. RILEY, C. V. [Address on entomology.] <Trans. Ill. State Hortic. Soc. for 1874, 1875, n. s., v. 8, pp. 103-111. Partial reprint: <Trans. Kans. State Hortic. Soc. for 1874, 1875, v. 4, pp. 103-104.
 - Ageria rubi [= Bembecia marginata] injurious to blackberry and raspberry; the cause and cure of scab in apples; ravages, food-plants, seasons, habits, and means against Chrysobothris femorata; evidence for the identity of the leaf-and root-forms of Phylloxera vastatrix; means against the same.
- 1455. RILEY, C. V. Discussion of the honey-bee. <Trans. Ill. State
 Hortic. Soc. for 1874, 1875, n. s., v. 8, pp. 131-132.

 Apis mellifica as an enemy to horticulture; importance of insects in the fertilization of flowers.
- 1456. RILEY, C. V. Notes on locusts. <Trans. Ill. State Hortic. Soc. for 1874, 1875, n. s., v. 8, pp. 136-137.</p>
 Native habitat of Caloptenus spretus.
- 1457. RILEY, C. V. Nonsense about the Phylloxera. <Colman's Rural World, 12 January, 1876. S.-b. No. 10, p. 3. See: <N. Y. Tribune, 1876. S.-b. No. 10, p. 27.
 - There is no such species as the American corn-grape, which is reported capable of resisting the attacks of *Phylloxera vastatria*.

- 1458. RILEY, C. V. Small borer in apple-twig. <Colman's Rural World, 26 January, 1876. S.-b. No. 10, p. 1.
 - Answer to letter of F. Holsinger; description and habits of Pseuccrus supernotatus.
- 1459. RILEY, C. V. Worms on cottonwood. <Colman's Rural World, 26 January, 1876. S.-b. No. 10, p. 1.
 - Answer to letter of J. H. Davidson; occurrence of larvæ of Drasteria ereckthea on Populus monilifera and on Trifolium; description of the imago.
- 1460. RILEY, C. V. Ailanthus silk-worm in Missouri. <Colman's Rural World, 27 January, 1876. S.-b. No. 10, p. 2.
 - Answer to letter of "Subscriber;" extent and unprofitableness of the culture of Samia [= Atlacus] cynthia; naturalization of the worm in the United States.
- 1461. R[ILEY], C. V. Cause of smut in wheat. <Colman's Rural World, 26 January, 1876. S.-b. No. 10, p. 2.
 - Critical review of a report by Pulaski Grange, Tenn., on the cause of smut in wheat; absurdity of the report; occurrence of Brachytarsus variegatus in the smut, and of larvæ of Cocidomyia, Meromyza, and Chlorops in the lower joints of wheat; smut caused by Ustilago segetum.
- 1462. BILEY, C. V. Colorado potato-beetle's native home. <N. Y. Tribune, 9 February, 1876. S.-b. No. 10, pp. 12-13.

 Geographical distribution of *Doryphora decombineats* prior to 1869.
- 1463. RILEY, C. V. An entomological question. <Prairie Farmer, 26
 February, 1876, v. 47, p. 68. S. b. No. 10, pp. 4, 5; No. 42, pp.
 76, 77. See: <Prairie Farmer, 4 March, 1876, v. 47, p. 76.
 S. b. No. 10, p. 4.
 - Reply to Proximo's "An entomological question;" commendation of legislative efforts to effect the appointment of a national entomological commission; text of the two bills introduced into Congress; criticism of the same.
- 1464. RILEY, C. V. Insect ravages. An interesting letter from Prof. C. V. Riley. How to protect our agricultural interests; legislation, wise and otherwise; the duty of Congress. <St. Louis Daily Globe-Democrat, 4 March, 1876, v. 1, p. 3. S.-b. No. 10, pp. 5-8.
 - Importance and extent of injuries inflicted by insects in North America; notice of existing legislation upon means against injurious insects; statement of legislation needed for protection against locusts; criticism of bills introduced into Congress for the appointment of a national entomological commission.
- 1465. RILEY, O. V. Notes on the Yucca borer, *Megathymus yucca*, Walk. <Trans. Acad. Sci. St. Louis, 10 January-23 March, 1876, v. 3, pp. 323-344, figs. 25-31. Separate: <St. Louis, Mo., January, 1876, pp. 23, figs. 25-31. Reprint: <8th Ann. Rept. State Ent. Mo., May, 1876, pp. 169-182, figs. 40-55.
 - See No. 1482 for synopsis of contents; see No. 1602.

- 1466. RILEY, C. V. Entomology. An interesting lecture on the insect world. The subject considered both practically and scientifically. <St. Louis Daily Globe-Democrat, 25 March, 1876, v. 1, p. 3. S.-b. No. 10, pp. 161-164. Reprint, with omissions: <Ware's Valley Monthly, August, 1876, v. —, pp. 281-289. S.-b. No. 14, pp. 163-167.
 - Definition of entomology; claims of the science as a liberalizing study; its economic importance; metamorphoses, abundance, and almost omnipresence of insects.
- 1467. [RILEY, C. V.] The insect world. Lecture by Professor Riley at Washington University. A practical subject for fruit-growers. <St. Louis Republican, 26 March, 1876. S.-b. No. 10, pp. 178, 179.
 - Advantages and interest of the study of insects as compared with that of other animals; economic importance of the study; ravages of Glossins morsitans.
- 1468. RILEY, C. V. Legislation in regard to insects injurious to agriculture. <Nation, 30 March, 1876, v. 22, p. 208.</p>
 Amount of insect injuries in the United States; demand for an independent commission; duties and limits of such a commission.
- 1469. RILEY, C. V. Entomology. Another lecture by Professor Riley. How to counteract the ravages of insects; direct remedies; practical hints to farmers, etc. Some of the duties of the State entomologist. <St. Louis Daily Globe-Democrat, 1 April, 1876, v. 1, p. 3. S.-b. No. 10, pp. 135-138; 179-184. Reprint, with omissions: <Ware's Valley Monthly, September, 1876, v. —, pp. 369-380. S.-b. No. 14, pp. 168-173.
 - Classification of means against insects; cause of the destructive occurrence of insects; importation of foreign pests; history of the introduction and spread of *Pieris rapæ*, *Phylloxera vastatrix*, and *Doryphora decemlineata*; natural history of these insects; spread of insects by small degrees; advantageous means against certain insects; encouragement of enemies of and parasites on noxious insects; need of distinguishing friends from foes; need of co-operation and of legislation for the destruction of insects; duties of a State entomologist.
- 1470. [RILEY, C. V.] Scarlet mite. <Colman's Rural World, 12 April, 1876, fig. S.-b. No. 10, p. 160.
 - Answer to letter of G. W. Barnes; occurrence, habits, and figure of Trombidium sericeum; habits of T. holosericeum; use of T. tinctorium as a dye.
- 1471. RILEY, C. V. Hibernation of Amphipyra [=Pyrophila] pyramidoides. <Psyche, March [13 April], 1876, v. 1, p. 152.</p>
 Extract from 3d Ann. Rept. State Ent. Mo., pp. 72-73, with additional note; this species sometimes hibernates as a pupa, and doubtless frequently as a moth.
- 1472. RILEY, C. V. Bag. worms and borers. How to protect our shade-trees and insure their growth. How to render shade-trees healthy. Letter from the State entomologist,

 St. Louis Re-

- 1472. RILEY, C. V.—Continued.
 - publican, 14 April, 1876, No. 16843, p. 3, 3 figs. S.-b. No. 10, pp. 173-175; 175-178.
 - Causes of the death of shade-trees in the city of St. Louis; description, figures, and natural history of and means against Thyridopteryx ephemerosformis and Chrysobothris femorata.
- 1473. RILEY, C. V. The locust plague; how to avert it. < Proc. Amer. Assoc. Adv. Sci. for 1875, 1876, v. 24, pp. 215-222. Separate: < Salem, April, 1876, pp. 215-222.
 - Extent of the ravages of Caloptenus spretus in 1873, 1874, and 1875; classification of and special remarks on the several means to be employed against the same; need of more extensive investigation of the insect in its native haunts and breeding places.
- 1474. R[ILEY], C. V. Honey locust weevil. <Colman's Rural World, 26 April, 1876.
 - Answer to inquiry of E. H. B.; larva of Spermophagus robinice has legs and spins, a cocoon.
- 1475. RILEY, C. V. Apple and peach borers. <Colman's Rural World, 9 May, 1876. S.-b. No. 16, p. 11.
 - Answer to inquiry of F. H.; habits, natural history, and means against Chrysobothris femorata and Ageria [= Sannina] exitiosa.
- 1476. RILEY, C. V. Notes on the codling-moth. <Colman's Rural World, 17 May, 1876. S.-b. No. 13, p. 153.

 Natural history of Carpocapsa pomonella.
- 1477. RILEY, C. V. Plums and cotton. <N. Y. Weekly Tribune, 17 May, 1876. S.-b. No. 10, p. 167.
 - Answer to letter of J. C.; a belt of cotton-batting around the trunk of the tree is no protection against the injury to plums by Conotrachelus nenuphar.
- 1478. RILEY, C. V. Rose bug remedy. <N. Y. Weekly Tribune, 17May, 1876, fig. S.-b. No. 10, p. 167.
 - Answer to letter of Mrs. S. P. Smith; description, figure, habits, and preferred food-plants of and means against *Macrodactylus subspinosus*.
- 1479. RILEY, C. V. Smut in wheat. <N. Y. Weekly Tribune, 17 May,
 1876. S.-b. No. 10, pp. 167-168. Reprint: <Colman's Rural World, 14 June, 1876. S.-b. No. 10, p. 169.
 - Critical review of two quoted communications by A. S. and by L. Heskett, on the cause of smut in wheat; smut caused by the growth of *Ustilago segetum*; Brackytarsus variogatus breeds in various smuts; description of the beetle.
- 1480. [RILEY, C. V.] Potato-beetle; progress. <N. Y. Weekly Tribune, 17 May, 1876. S.-b. No. 10, p. 168.
 - Arrival of *Doryphora decemlineata* at the Atlantic coast; its ravages there and means against them.
- 1481. RILEY, C. V. Locusts as food for man. <Proc. Amer. Assoc. Adv. Sci. for 1875, 1876, v. 24, pp. 208-214. Separate: <Salem, May, 1876, pp. 208-214.
 - Reference to previous writings on the use of locusts as food for man; historical evidence of the extensive use of locusts as food; methods of preparing locusts for food; species used hitherto; use of Caloptenus spretus; peculiarities of individual taste or national custom.

1482. BILEY, C. V. Eighth annual report on the noxious, benefit	rial.
and other insects of the State of Missouri, made to the S	
Board of Agriculture, pursuant to an appropriation for	
purpose from the legislature of the State. <11th Ann. R	
State Board of Agric. for 1875, May, 1876, pp. 185+4, 55	
Separate: <jefferson 185+4,="" 1876,="" 55<="" city,="" may,="" mo.,="" pp.="" td=""><td>igs.</td></jefferson>	igs.
Preface	III
TABLE OF CONTENTS	V
NOXIOUS INSECTS.	
Mary Classics and market Developed 10 12 and a	•
THE COLORADO POTATO-BEETLE, Doryphora 10-lineata	1
Damage during the year, 1—Abundant in Atlantic States, 1—	
Swarming on Coney Island, 2—Injuring egg-plant, 2—Its scien-	
tific name, 2—Additional enemies, 3—Eaten by the crow, 3—Rem-	
edies, 3—Cost of applying Paris green, 3—Preparing the poison,	
3—Use of straw as a protection, 4—Machine for sprinkling, 4—	
Machine for brushing off the insects, 4—Experience with Paris	
green, 5—Experiments of Profs. R. C. Kedzie and Wm. McMurtrie	
show that it may be used with safety, 6—Trial of other remedies,	
6—The insect's native home, 8—The theory that it came from the	
Rocky Mountain region essentially correct, 10—Poisonous quali-	
ties of the insect discussed, 10.	••
CANKER-WORMS, Paleacrita vernata; Anisopteryx pometaria	12
Two species long confounded, 12—They differ generically; new	
genus (Paleacrita) proposed for one, 13—The two compared in	
all stages, 13, 17—Characters of the genus Paleacrita, 17—Distin-	
guished as spring and fall canker-worms, 17—Practical consid-	
erations from their differences of habit, 18—Stunting the larve	
does not produce male moths, 19—Traps recommended, 20, 21.	00
THE ARMY WORM, Leucania unipuncta	22
Its generic name, 22—The term "army-worm" applied to various	
insects, 23—Past history of the army-worm, 24—Known since	
1854 in Missouri, 27—It followed the 1871 conflagration around Peshtigo, Wis., 28—Its history in 1875; very general all over the	•
country, 28, 29—Its history in Missouri in 1875, 30—Sexual dif-	
ferences, 30—Sexual organs illustrated, 30, 32—Natural history	
of the species, 32—Illustrated in all states, 32, 33—It occurs in	
Europe, Asia, New Zealand, and Australia, 34—Description of	
the egg, 34—Where the eggs are laid, 34—Conclusions drawn	
from structure, 36, 37,—When the eggs are laid, 40—In what state	
does the insect hibernate?, 43—Habits of the worm, 45—Why	
it escapes detection when young, 45—Why it travels in armies,	
46—Time of its appearance, 46—Are there one or two broods?,	
47—The fall army-worm, 48—How distinguished from the real	
army-worm, 48—Plants preferred by the army-worm, 49—Its sud-	
den appearance and disappearance, 50—It swarms during wet pre-	
ceded by very dry seasons, 51—Its natural enemies, illustrated,	
52—Remedies, 54—Philosophy of winter burning, 54, 55—Preven-	
tion, 55—Summary of the leading facts concerning it, 56.	
THE ROCKY MOUNTAIN LOCUST, Caloptenus spretus	57
Previous experience in spring of 1867, 57—Predictions verified, 58—	
General outlook in spring of 1875, 60—Extent of country ravaged,	
60—The outlook in Missouri, 61—Country ravaged often as bare	
as in mild winds of Assumet by counting 69 Atchien County	

as in midwinter, 61—Account by counties, 62—Atchison County,

THE ROCKY MOUNTAIN LOCUST-Continued.

62-Andrew County, 62-Benton County, 63-Barton County, 63—Bates County, 63—Buchanan County, 64—Caldwell County, 64-Cass County, 64-Clay County, 67-Clinton County, 68-Dade County, 68-De Kalb County, 69-Gentry County, 69-Hickory County, 69-Holt County, 69-Henry County, 69-Jackson County, 69-Johnson County, 72-Lafayette County, 73-Nodaway County, 73-Newton County, 73-Pettis County, 73-Platte County, 73-Ray County, 74-St. Clair County, 75-Vernon County, 76-Condition of things in other States, 76-Kansas, 76-Nebraska, 79-Iuwa, 81-Minnesota, 81-Colorado, 84-Dakota, 85-Montana, 87-Wyoming, 88-Texas, 88-Indian Territory, 88-Manitoba, 89-Damage done in Missouri, 80-Destitution in Missouri, 91—Address of relief committee from Saint Louis Merchants' Exchange, 93-Cases of starvation, 94-The Governor's proclamation, 95-The locusts not a divine visitation, 97-Natural history; mode of molting illustrated, 98-Habits of the unfledged young, 100-Directions in which the young travel, 101-Rate at which they travel, 102-They reached but a few miles east of where they hatched, 102-Not led by "kings" or "queens," 103-The species taken for such, illustrated, 103, 104-The exodus in 1875, 104—Time of leaving of the winged insects, 104—Direction taken by the winged insects, 105-Destination of the departing swarms, 106-Native home of the species, 109-Views previously expressed confirmed, 110-Conditions of migration, 112-Conditions which prevent the permanent settlement of the species in Missouri, 113-Modification of the species by climatic conditions, 114, 155-Definition of the species, 114-How distinguished in all stages from species most nearly allied, 117—Experience in spring of 1875, 118—Contrast in summer and fall, 119— No evil without some compensating good, 120—Injury to fruit and fruit trees, 121-Food-plants, 121-Only one kind of plant not touched under all circumstances, 121—Changes that followed the locusts, 121-The widespread appearance of a new grass, ordinarily unnoticed, 122-Appearance of large worms, 123-The locusts did not return in the fall, 124-Natural enemies, 124-Remedies against the unfledged insects, 125—Artificial means of destroying the eggs, 125-Various means of destroying the unfledged young, 126-They are within man's control, 126-The proper ditch to make, 128-Machines used in Colorado, 129-Best means of protecting fruit trees, 130—How to avert locust injuries, 131—Prevention, 131-Legislation, both national and local, 132-Bills before the Forty-fourth Congress, 133—Need of a national entomological commission, 133—The bounties offered in Minnesota, 138-The requisites of a good bounty law, 138, 139-How a bounty law would work, 140-Suggestions, 140-Lessons of year, 142-Locusts as food for man, 143-They have been used from time immemorial, and are used extensively at the present day, 145-The Rocky Mountain species quite palatable, 146-Mode of preparation, 147-False opinions and predictions, 148-Unnecessary alarm caused by other species, 148-Injuries of native species in 1875, 150-Locust flights in Illinois in 1875, 151-They were composed of local species, 152, 153—Explanation of these flights, 154— Locust prospects in 1876, 155-No danger from them in Missouri, 156.

THE GRAPE PHYLLOXERA, Phylloxera vastatrix..... The injuries not great in Missouri in 1875, 157—Completion of its natural history, 157-Where the winged female lays her eggs, 157, 161—The sexed individuals illustrated, 158—Description of the true female, 159—Description of the impregnated egg, 159, 162-Practical considerations growing out of these latest discoveries, 163-Decortication of the bark to destroy the impregnated egg, 163-The insect may be imported from one country to another on cuttings as well as rooted plants, 163-Best time to attack the root-lice, 163-Phylloxera ravages in California, 163-Great destruction around Sonoma, 164-Need of action by the State authorities, 164—Occurrence of Phylloxera in the Southern States, 164-Report of committee appointed by the American Pomological Society, 165-Its occurrence in Georgia, 166, 167-American grape-vines in Europe, 167-Large demand for our vines, 167-The American vines flourishing in Southern France where the European varieties perish, 167—The orders for some varieties exceeded the supply, 168—Probable future demand, 168.

INNOXIOUS INSECTS.

NOXIOUS INSECTS-Continued.

SUPPLEMENTARY NOTES ON THE ARMY-WORM, Leucania unipuncta

Completion of its natural history, 182—Oviposition of the moth described, 183—Eggs described, 183—Conclusions previously arrived at verified, 184—Description of the different larval stages, 184.

1483. RILEY, C. V. Is the Colorado potato-beetle poisonous? < Colman's Rural World, 7 June, 1876. S.-b. No. 10, pp. 164-165.

Extract from the 6th Ann. Rept. State Ent. Mo., pp. 10-12; abstract and
critical review of Grote and Kayser's "Are potato-bugs poisonous?"; considers the experiments inconclusive.

1484. [RILEY, C. V.] The eggs of the army-worm. <Colman's Rural World, 7 June, 1876. S.-b. No. 10, p. 170. Reprint: <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 211 Proc. <Amer. Nat., August, 1876, v. 10, pp. 508-509.

Reasons why eggs of Leucania unipuncta have not been observed heretofore; verification of author's conclusions as to the probable method of eviposition of this species; description of eggs and young larvæ.

157

- 1485. R[ILEY], C. V. Persian insect powder. <N. Y. Tribune, 7 June, 1876. S.-b. No. 10, p. 168.
 - Answer to letter of G. W. Holmes; impossibility of prescribing means against unknown insects; Paris green unsuitable for protection of bearing fruit trees; pyrethrum recommended.
- 1486. R[ILEY], O. V. Cocoons of silkworms. <N. Y. Tribune, 7 June, 1876. S.-b. No. 10, p. 168.
 - Answer to letter of A. R. Sprout; description of larva and cocoon of Callosamia [= Atlacus] promethea; colors of imagos; food-plants of larvæ; deposition of eggs; silk of little value.
- 1487. [RILEY, C. V.] Is Paris green absorbed? <N. Y. Tribune, 7 June, 1876. S.-b. No. 10, pp. 168-169.
 - Answer to letter of M. F.; plants colored by certain tinctures placed at their roots; Paris green not absorbed into the tissues of plants in perceptible quantities, but neutralized in the soil.
- 1488. RILEY, C. V. Ditching for young locusts. <Colman's Rural World, 14 June, 1876. S.-b. No. 10, pp. 171-172; 203-204; 204-205.
 - Critical review of J. Stayman's article on same subject; proper width and depth of ditches to check the march of unfledged Caloptenus spretus and other locusts.
- 1489. RILEY, C. V. Periodical Cicada, "17-year locust." < N. Y. Semi-weekly Tribune, 23 June, 1876, 3 figs. S.-b. No. 10, pp. 166-167.
 - Occurrence of Cicada [= Tibicen] septendecim at Lexington, Va., in 1876; list of localities at which these insects will appear this year; chronological history of a brood; figures of larva, pupse, and imago.
- 1490. RILEY, C. V. Specific for Colorado potato beetle. <Colman's Rural World, 28 June, 1876. S.-b. No. 10, p. 160. Reprint, with omissions: <N. Y. Tribune. S.-b. No. 14, p. 7.
 - Answer to letter of R. Barbour; directions for the use of Paris green as a means against Doryphora decemlineata.
- 1491. RILEY, C. V. [Inquiries concerning the Colorado potato-beetle, and Mayor Brown's answer thereto.] < Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 170-172 Proc.
 - Verbal communication; critical review of answer of Mayor Brown to inquiries of A. C. Hardy de Boislieu as to the means of preventing the introduction of Doryphora decemiineata into Belgium; quotes the 6th Ann. Rept. State Ent. Mo., p. 16, showing that the fear of such introduction is not unfounded; inaccuracy of Mayor Brown's opinions; manner in which Doryphora decemiineata will probably reach Europe if at all.
- 1492. RILEY, C. V. [On the ravages of young locusts in western Missouri.] <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 179–180 Proc.
 - Verbal communication; alarming nature of the ravages of unfiedged Caloptonus spretus in western counties of Missouri at the present time; probable future departure of the winged locusts: means to be adopted against the locusts now.

- 1498. RILEY, C. V. [Lecture on the Rocky Mountain locust.] < Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 180 Proc.
 - Brief abstract of lecture; prediction of the speedily approaching end of the ravages of Caloptenus spretus in Missouri for 1875; plentiful crops to be expected subsequently.
- 1494. RILEY, C. V. [Predictions verified.] < Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 185 Proc.
 - Verbal communication; accuracy of author's conclusions as to the probable doings of Caloptenus spretus in Missouri later in the season.
- 1495. RILEY, C. V. [On changes in vegetation caused by locusts.] <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 188–189 Proc. See: <Amer. Nat., February, 1876, v. 10, p. 125.
 - Verbal communication; remarks on the extraordinary development of Vilfa vaginatora for a season after the ravages of Caloptonus spretus have occurred an illustration of "the struggle for existence;" the interesting character of such instances of abnormal multiplication of a species; criticism and defense of the expressions "struggle for existence" and "natural selection."
- 1496. RILEY, C. V. Jumping seeds and galls. <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 190-192 Proc. Reprint: <Gardener's Mo. and Hortic., July, 1878, v. 20, pp. 213-214. See: <Amer. Nat., February, 1876, v. 10, p. 125. Pacific Rural Press, 6 April, 1878.
 - Description of the seeds (of an unknown plant) which are inhabited by the larvæ of Carpocapsa salitians; manner in which these larvæ cause the seeds to roll and jump; habits of larva within the seed; description of the plant bearing the seeds; the seed of Tamariscus moved by the larva of Nanedes tamarisci, which feeds within it; description of the gall of Cymips [= Non-roterus] querous-salitatorius and of the jumping of these galls.
- 1497. RILEY, C. V. [On the use of Paris green as an insecticide.]

 Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 193. See:
 Amer. Nat., February, 1876, v. 10, p. 126.
 - Verbal communication; confirmation, by experiments of R. C. Kedzie, of author's conclusions in regard to the safety of Paris green as an insecticide; this substance metamorphosed into a less soluble form in the ground and held in the ground if not used to excess; when applied in small but sufficient quantities not injurious to plants; objections to the use of the "Potato-pest poison" made at the Lodi [N. J.] Chemical Works.
- 1498. RILEY, C. V. New use for the American Agave. < Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 195-196 Proc. See: < Amer. Nat., February, 1876, v. 10, p. 126.
 - Verbal communication; use of the dried flower-stalk of Agave americanus for lining insect-boxes.
- 1499. RILEY, C. V. [Food of insectivorous plants.] < Trans. Acad. Sci. St. Louis, June, 1876, v. 3, pp. 201-202 Proc.
 - Drosera, Dionæa and other plants digest, absorb, and appropriate nitrogenous matters; glands for the appropriation of animal food not yet found in the Sarracenias.
- 1500. RILEY, C. V. Parasites on bees. <Trans. Acad. Sci. St. Louis, June, 1876, v. 3, p. 212 Proc.
 - Bees in California infested with triungulins of a Meloë sp. which sometimes kill the bees; usual habits of such triungulins.

- 1501. RILEY, C. V. The locust pest. <Sci. Amer., 1 July, 1876 [v. 49], n. s., v. 35, p. 9.
 - Effects of water and cold on the eggs of Caloptenus spretus.
- 1502. RILEY, C. V. Swallows; bed bugs. < Colman's Rural World, 5
 July, 1876. S. b. No. 10, p. 166.
 - Answer to letter of R. R. Pierce; nests of Hirundo fulva generally infested by Acanthia lectularia; those of Hirundo americana rarely so; while this habit might cause annoyance about a dwelling, it is of little consequence when the bird nests under the eaves of a church.
- 1503. [RILEY, C. V.] Berry and cherry twigs. <N. Y. Tribune, 5 July, 1876. S. b. No. 10, p. 169.
 - Answer to letter of W. Keyser; eggs of *Œcanthus niveus* laid in twigs of *Rubus* and *Prunus*; description, habits of, and means against *Oberea tripunctata*.
- 1504. R[ILEY], C. V. Three worms and their work. <N. Y. Weekly Tribune, 12 July, 1876, 2 figs. S.-b. No. 14, p. 132.
 - Answer to letter of "Subscriber;" means against larve of Agrotidides, of Elaterides, and earth worms, Lumbrious sp.; eggs of Agrotis ypsilon laid in the spring; breeding-habits and economic importance of Lumbricus; figures of an elatrid larva and imago.
- 1505. [RILEY, C. V.] Mite parasites of the Colorado potato-beetle. <Mirror and Farmer, 15 July, 1876, v. 28, No. 29, p. 2. S.-b. No. 14, p. 221. Reprint: <Gardener's Mo. and Hortic., September, 1876, v. 18, p. 279. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 219 Proc. Note: <Amer. Nat., October, 1876, v. 10, p. 636.
 - Ectoparasitism of a gamasid mite [Uropoda americana] on Doryphora decemlineata; list of vertibrate enemies of D. decemlineata.
 - The reprints note the occurrence of *D. decemlineata* in New Hampshire and its ravages along the Atlantic coast. See No. 1610.
- 1506. [RILEY, C. V.] [Wheat insects.] < Blair [Nebr.] Times, 20 July, 1876. S.-b. No. 10, pp. 200-201.
 - Quotes from 1st Ann. Rept. State Ent. Mo., pp. 159-161; wheat injured in Nebraska in 1876 by Cecidomyia destructor and Meromyza americana; ravages of larva, description of imago and means against the latter; occurrence of Disonycha flaviventris in Nebraska.
- 1507. RILEY, C. V. A new enemy of wheat. <N. Y. Tribune, 21 July, 1876. S. b. No. 14, pp. 129-130; 130-131.
 - Answer to letter of W. Robson; occurrence, habits, and ravages of Leucania albilinea in Maryland, Pennsylvania, and Kansas; history and description of larva, pupa, and image of the insect; evolution of new habits and forms among insects.
- 1508. [BILEY, C. V.] Chinch-bug; bee-moth. <Colman's Rural World, 26 July, 1876. S.-b. No. 10, pp. 201-205.
 - Answer to letter of G. R. Christian; means against Blissus leucopterus and Galleria cercana; food, seasons, habits, and description of the latter.
- 1509. RILEY, C. V. The grape-root borer, *Ægeria polistiformis*. <Colman's Rural World, 26 July, 1876, 2 figs. S.-b. No. 10, pp. 205-206.
 - Answer to letter of F. J. Kron; experiments on means against *Ægeria* [= Soiapteron] polistiformis; habits of larva; figures of imago; geographical distribution.

- 1510. RILEY, C. V. Sweet-potato beetles; "beautiful bugs." <N. Y. Weekly Tribune, 26 July, 1876, 3 figs. S.-b. No. 10, p. 213. Answer to letter of W. Snowden: characters and habits of Cassidida: descrip-</p>
 - Answer to letter of W. Snowden; characters and habits of Cassididæ; description and figures of larvæ and imagos of Coptocycla [= Cassida] bivittata and C. [= C.] nigripes; descriptions of C. aurichalcea and C. guttata; these species feed on leaves of Ipomæa batatas and C. aurichalcea, also on leaves of Convolvulus and of Solanum dulcamara; means against them.
- 1511. [RILEY, C. V.] Those centennial insects. <N. Y. Weekly Tribune, 26 July, 1876. S.-b. No. 10, p. 219.</p>
 Criticism of communication of L. A. M.; list of principal insects preying upon stored corn.
- 1512. R[ILEY], C. V. Wheat midge; "rue-worms." <N. Y. Tribune, 2 August, 1876. S.-b. No. 10, p. 202.
 Larvæ of Papilio asterias feeding on Ruta graveolens; description, habits, seasons, ravages of, and means against Cecidomyia [= Diplosis] tritici.
- 1513. RILEY, C. V. Apple and peach borers. <Colman's Rural World,</p>
 9 August, 1876. S.-b. No. 14, pp. 3-4; 55-56. Extract: <Cultivator and Country Gentleman, 7 September, 1876, v. 41, p. 566.</p>
 Answer to inquiries of F. H.; Chrysobothris femorata distinguished from Saperda bivittata [= candida] and Egeria [= Sannina] exitiosa; habits and description of the first; habits of the last and means against both.
- 1514. RILEY, O. V. Large saw-fly. <Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 4.
 - Answer to inquiry of J. B. J.; description of larva and image of Cimber laportei [= americana]; its food-plants and transformations.
- 1515. RILEY, C. V. Cottony scale-insect on maples. <Colman's Rural World, 9 August, 1876. S.-b. No. 14, pp. 4-5.

 Answer to inquiry of C. F. Mills; Locanium accricola [= Pulvinaria innumerabilis] injurious to Accr Casycarpum at Springfield, Ill.; habits, description of the scale, and manner of oviposition; no great injury ever done by insects of this genus.
- 1516. RILEY, C. V. Hickory vs. locust borer. < Colman's Rural World,
 9 August, 1876. S.-b. No. 14, p. 5.
 Answer to inquiry of F. M. D.; description of larva and image and natural history of Arhopalus [= Cyllene] pictus; characters distinguishing this from
- A. [= C.] robinia; food-plants and seasons of the latter.

 1517. RILEY, C. V. Stag-beetle. <Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 5.

 Answer to inquiry of S. Lee; occurrence and characters of Lucanus elaphus;
- food of larva.

 1518. RILEY, C. V. Eggs of the angular-winged katydid. <Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 5.
- Rural World, 9 August, 1876. S.-b. No. 14, p. 5.

 Answer to inquiry of V. Kriegel; descriptions of eggs, images, and song of

 Microcentrum retinerve; habits and ravages and means against the same.
- 1519. RILEY, C. V. Experience with the Colorado potato-beetle. <Colman's Rural World, 9 August, 1876. S.-b. No. 14, p. 6.
 - Letter from N. Coleman; larvæ of Doryphora decembineata in confinement pupated on the surface of the ground; images eat the tubers underground.

- 1520. [RILEY, C. V.] An elm enemy. <N. Y. Semi-weekly Tribune, 11 August, 1876. S.-b. No. 14, pp. 54-55.
 - Habits and ravages of and means against Galeruca calmariensis [= xanthomelona].
- 1521. RILEY, C. V. The locust in 1876. <N. Y. Weekly Tribune, 16 August, 1876. S.-b. No. 14, p. 73.
 - The occurrence and ravages of Caloptenus spretus in Minnesota and Colorado in 1876; their absence from other portions of the country occasionally inhabited by them; their destruction by Astoma gryllaria [= Trombidium locustarum] and inability to maintain themselves in parts of the region they infest.
- 1522. [RILEY, C. V.] The war on "corn-worms." <N. Y. Semi-weekly Tribune, 18 August, 1876. S.-b. No. 14, p. 73.
 - Review of article by A. Brewster; ravages of and means against larve of Elateridæ, Agrotididæ and Lachnosterna quercina [=fusca] and the "grubworm," injurious to growing maize.
- 1523. RILEY, C. V. Spined soldier-bug. <Ohio Farmer, 19 August, 1876, v. 50, p. 118.
- 1524. RILEY, C. V. "Potato-pest poison." <Sci. Amer., 19 August, 1876 [v. 49], n. s., v. 35, p. 116.

 Condemning some of the patent poisons for Doryphora decemlineata.
- 1525. RILEY, C. F. Cottonwood borers. <N. Y. Weekly Tribune, 23
 <p>August, 1876. S.-b. No. 14, p. 55.

 Answer to inquiry of J. R.; food-plants and ravages of and means against Pleatrodera scalator.
- 1526. RILEY, C. V. Harmless insects. <N. Y. Weekly Tribune, 23
 August, 1876. S.-b. No. 14, p. 55.

 Answer to letter of a "Subscriber;" occurrence of Procus venous on appletrees; habits and molting.
- 1527. RILEY, C. V. Locust prospects. <N. Y. Tribune, 6 September,
 1876. S.-b. No. 10, pp. 213-215; No. 14, pp. 74-75. Reprint:
 <Prairie Farmer, 16 September, 1876, v. 47, p. 298. S.-b. No. 10, pp. 207-209.
 - Answer to letter of G. H. H.; movements and ravages of Caloptenus spretus in Aug., 1876, in Dakota, Minnesota, Iowa, Nebraska, and Colorado.
- 1528. RILEY, C. V. Some notes on potato-beetles. <Sci. Amer., 9
 September, 1876 [v. 49], n. s., v. 35, p. 164. Reprint: <Ohio
 Farmer, 16 September, 1876, v. 50, p. 179. S.-b. No. 14, pp.
 6, 7.
 - Critical review of S. R. M.'s "Facts about potato-beetles;" Doryphora decemlineata flies in the day-time, but not at night; feeds upon several species of plants; hibernates as an imago, which should be destroyed in early spring.
- 1529. RILEY, C. V. Entomological works wanted. <Colman's Rural World, 20 September, 1876. S.-b. No. 14, p. 2.
 - Answer to letter of J. W. Newman; notice of some works on entomology for the use of students.

- 1530. [RILEY, C. V.] Grape-leaf Philloxera enemy. <Colman's Bural World, 20 September, 1876. S.-b. No. 14, p. 3.
 - Answer to letter of A. Engleman; Leucopis sp. ? parasitic in galls of Phylloxera vastatrix and P. rilegi.
- 1531. [RILEY, C. V.] Grape-leaf gall. <Colman's Rural World, 20 September, 1876. S.-b. No. 14, p. 3.
 - Answer to letter of A. A. Briggs; Phylloxera rastatrix does comparatively little damage on the leaves of grape-vines; Clinton and Taylor vines peculiarly subject to the attacks of the leaf-inhabiting form; Concord vines seldom attacked by it and not seriously injured by the root-inhabiting form; means against the leaf-inhabiting form.
- 1532. RILEY, C. V. New locust theory wanted. <Colman's Rural World, 27 September, 1876. S.-b. No. 14, p. 62.
 - Answer to letter of W. T. D.; facts not theories wanted in regard to the flights of Caloptonus spretus.
- 1533. RILEY, C. V. Cecropia worm on elder. <Colman's Rural World, 27 September, 1876, 1 fig. S.-b. No. 14, p. 63.
 - Answer to letter of A. S. Van Winkle; description of larva, cocoon, and imago of Samia [= Attacus] cecropia; figure of larva; habits and seasons; availability of the cocoon for silk.
- 1534. RILEY, C. V. The harlequin cabbage-bug. <Colman's Rural World, 4 October, 1876. S. b. No. 14, p. 56.
 - Answer to letter of R. J. Waters; habits, ravages, distribution, and means against Strachia [= Murgantia] histrionica; colors of early stages and of images.
- 1535. [RILEY, C. V.] Butterfly chrysalis. <N. Y. Tribune, 13 October, 1876. S.-b. No. 19, p. 224. Reprint: <N. Y. Tribune, 21 October, 1876. S.-b. No. 10, p. 223.
 - Answer to inquiry of E. B. S.; occurrence of *Danais archippus* in swarms in central United States in autumn of 1876; colors of larva and pupa; larva feeds on *Asclepias*.
- 1536. [RILEY, O. V.] Domesticated katydid. <N. Y. Tribune, 18 October, 1876.
 S.-b. No. 19, p. 214. Reprint: <N. Y. Tribune, 21 October, 1876.
 S.-b. No. 10, p. 219.
 - Answer to letter of C. A. P.; longevity and food of Microcentrum retinerve.
- 1537. [RILEY, C. V.] Unjust accusation? <N. Y. Tribune, 21 October, 1876. S.-b. No. 10, p. 209; No. 19, p. 242.
 - Answer to inquiry of a "Correspondent;" Harpalus erraticus accused of destroying shrubbery; predaceous habits of Carabide.
- 1538. RILEY, C. V. The Rocky Mountain locust. <Colman's Rural World, 1876, v. —, 30 October, 6 November, 13 November.
 S.-b. No. 10, pp. 185–188. See: <Kansas Farmer, November, 1876.
 S.-b. No. 10, pp. 225–235. <N. Y. Tribune, October, 1876.
 S.-b. No. 10, pp. 215.
 - Verification of predictions in regard to the limitation of the ravages of Caloptenus spretus in western Missouri in 1876; gathering eggs, ditching, rolling, and burning recommended as means against them; recommends the employment of soldiers, of hogs, and of poultry, late planting, and the destruction of the locusts in their native country; Blissus leucopterus more

injurious in Western Missouri in 1876 than Caloptonus spretus; prospect of freedom from injury by the former for the next two years; Vilfa ragino-flora introduced into the eastern prairies by the locusts; limit to the eastern range of the locusts; they are not led by kings or queens; occurrence of larves of Deilephila lineata after the disappearance of the locusts; locust flights in Illinois.

- 1539. [RILEY, C. V.] Canker-worms at the West. <N. Y. Tribune, 31
 October, 1876. S. b. No. 17, p. 59.
 - Occurrence of *Paleacrita vernata* in Michigan in 1872 and in Ohio in 1874, 1875, and 1876; brief account of *P. vernata* and *Anisopteryz pometaria*; their habits, ravages, and means against them.
- 1540. RILEY, C. V. [The venation of Anisopteryx and variation in images.] < Ca. Ent., September [October], 1876, v. 8, pp. 178-179.
 - Variation in the venation of the wings of Anisopteryx pometaria and in the images of so-called species.
- 1541. RILEY, C. V. A new enemy of the grasshopper. <Lawrence [Kans.] Journal. Reprint: <Industrialist [Manhattan, Kans.], 2 November, 1876, v. 2, No. 30, p. 2. S.-b. No. 10, pp. 209-210. Letter of F. H. Snow with comments; eggs of Caloptenus spretus destroyed by larvæ of Anthomyia calopteni [=angustifrons]; characters of the larvæ; eggs destroyed by larvæ of some Ichneumon? [=Systachus oreas]; need of destroying eggs by artificial means.
- 1542. RILEY, C. V. Silk culture in Kansas. Nationalist, 10 November, 1876. S.-b. No. 14, p. 1.

 Importance of silk culture; practicability of the culture and prospects of its increase in the United States; measures for its promotion in Kansas; sac-
- 1543. RILEY, C. V. Bee killers: Asilus flies. <N. Y. Weekly Sun, 15 November, 1876. S.-b. No. 10, p. 201.
 Habits of Asilidæ, especially of Trapanea [= Promachus] apirora and Asilus missouriensis [= Proctacanthus milberti]; description of T. [= P.] apivora; larva of Asilus sericeus feeds on roots, of Rheum rhaponticum.

cessful raising of silk worms on osage orange [Maclura aurantiaca].

- 1544. RILEY, C. V. How to use Paris green for the cotton-worm.

 Colman's Rural World, 15 November, 1876. S.-b. No. 10,
 pp. 210-211. See: <N. Y. Tribune, 15 December, 1876. S.-b.
 No. 14, p. 7.
 - Answer to letter of C. W. Niver; directions for the use of Paris green by the dry and wet method; description of sprinkling machines.
- 1545. [RILRY, C. V.] Notodonta concinna. <Colman's Rural World, 15 November, 1876. S.-b. No. 10, p. 219.
 - Answer to letter of J. Barritt; description of larva of Notodonta [= Œdema-sia] concinna; habits, food-plants, and means against the same.
- 1546. [RILEY, O. V.] The dog-day harvest fly. <Colman's Rural World, 15 November, 1876. S.-b. No. 10, p. 220.
 - Answer to inquiry of C. A. U.; description of Cicada pruinces [= tibicen]; method of sonifaction; habits; notice of some manuals of entemology.

- 1547. [RILEY, C. V.] Snake-worms. < Colman's Bural World, 15 November, 1876. S. b. No. p. 10, p. 220.
 - Answer to letter of J. Armstrong; description of larvæ of Sciara sp.; their habit of moving in congregations; their abode and enemies.
- 1548. RILEY, C. V. Locust eggs. < Colman's Rural World, 15 November, 1876. S.-b. No. 10, p. 223.
 - Answer to letter of C. T.; female Acridide lay more than one litter of eggs; the occurrence of egg-like parasites in males has led to the supposition that the males bore eggs.
- 1549. RILEY, C. V. Locust flights east of the Mississippi. < Colman's Rural World, 22 November, 1876. S. b. No. 10, pp. 211-212; 221-222; 223-225. Reprint: Sci. Amer., 16 December, 1876 [v. 49], n. s., v. 35, p. 392, 2 figs. S.-b. No. 10, pp. 216, 217. <Trans. Kans. Acad. Sci., 1877, v. 5, pp. 62-64.</p>
 - Limitation of plants and animals to certain geographical regions; regions in which alone Caloptenus spretus survives; species confounded with C. spretus: occurrence of swarms of Acridium americanum in Ohio; description, geographical distribution, and ravages of the same; swarms of Caloptenus differentialis, C. atlanis, and C. femur-rubrum in Illinois; ravages of the same. The reprint in the Sci. Amer. contains figures of Caloptenus spretus and Acridium americanum.
- 1550. RILEY, C. V. Locusts again. < N. Y. Tribune, 22 November, 1876. S.-b. No. 10, p. 222. Effects of winter on the vitality of the eggs of Acridida.
- 1551. RILEY, C. V. The army-worm; its natural history complete-<Sci. Amer., 9 December, 1876 [v. 49], n. s., v. 35, p. 372, 4 figs. S.-b. No. 10, pp. 217-219. Reprint, with slight changes: <Proc. Amer. Assoc. Adv. Sci. for 1876, 1877, v. 25, pp. 279-</p>
 - Geographical distribution of Loucania unipuncta; figures of all stages and of ovipositor of female; place and manner of oviposition; description of eggs and young larvæ; number of annual broods; summary of the natural history of this species.
- 1552. RILEY, C. V. The apple-bark louse. <Ca. Farmer, 15 December, 1876. S.-b. No. 14, p. 50.
 - Seasons, habits, and ravages of Mytilaspis pomicorticis [= pomorum]; description of young larvæ; of males and females and formation of scales.
- 1553. RILEY, C. V. The apple maggot; a formidable enemy. < N. Y. Semi-weekly Tribune, 15 December, 1876. S.-b. No. 14, pp. 7-8. Answer to letter of P. M. Augur; description of larva and image of Trypeta pomonella; ravages, food-plants, habits, and means against the same; literature of the subject.
- 1554. RILEY, C. V. Entomological notes; confounding friend with foe. < Colman's Rural World, 20 December, 1876. S.-b. No. 14, p. 4. Description of scales and eggs of Diaspis harrisii [= Chionaspis furfurus] and

of colors of larva, pupa, and image of Chilocorus bivulnerus; ravages of the

Diaspis; habits and usefulness of the Chilocorus.

- 1555. RILEY, C. V. Locust injury next spring. The territory in Missouri that will probably suffer therefrom. <Colman's Rural World, 20 December, 1876. S.-b. No. 14, pp. 5-6. Reprint: <Industrialist [Manhattan, Kans.], 17 February, 1877, v. 2, No. 44, p. 4. S.-b. No. 14, p. 49.
 - Prediction of ravages of Caloptenus spretus to occur in spring of 1877 in Missouri.

similarity in habits of Oncideres amputator and E. putator to those of O.

- 1556. [RILEY, C. V.] Amputating insects. <N. Y. Semi-weekly Tribune, 29 December, 1876, 2 figs. S.-b. No. 14, pp. 8-9.

 Description of the work of Elaphidion putator [= villosum] and Oncideres cingulata; habits of both; figures of larva, pupa, and image of the latter;
- oingulate.

 1557. [RILEY, C. V., et al.] The Rocky Mountain locust or grasshopper, being the report of proceedings of a conference of the Governors of several Western States and Territories, together with several other gentlemen, held at Omaha, Nebr., on the 25th and 26th days of October, 1876, to consider the locust problem; also a summary of the best means now known for counteracting the evil. <St. Louis, 1876, 8°, pp. 3+58, 8 figs.

See: <N. Y. Tribune, 1876. S.-b. No. 10, p. 223.

- Preface, proceedings, pp. 1-36. Practical considerations and suggestions for the suppression of Caloptenus spretus; description, oviposition, transformations, and habits of the same; its enemies and parasites; means against it; description of Anthonyia calopteni n. sp. [= angustifrons]. Figures. Prepared by J. S. Pillsbury, P. Pusey, and C. V. Riley.
- 1558. BILEY, C. V. Potato pests. Being an illustrated account of the Colorado potato-beetle and the other insect foes of the potato in North America, with suggestions for their repression and methods for their destruction. <New York: Orange Judd Company [1876], pp. 108, 49 figs., map. Review: <Cultivator and Country Gentl., 11 January, 1877, v. 42, p. 25. Reply to review: <1bid., 1 February, 1877, v. 42, p. 78. S.-b. No. 14, pp. 56-57.

TABLE OF CONTENTS.

TREFACE	7
INTRODUCTION	g
THE COLORADO POTATO-BRETLE, Doryphora 10-lineata	11
Its past history	11
Prediction that it would reach the Atlantic, 12—Its march across the country, 13—It reaches the Atlantic, 14—Its swarming in large cities, 16—Its occurrence out at sea, 17.	
The insect's native home	12
When it first attacked the potato, 18.	17
Rate at which it traveled	21
How it traveled	21
Mostly in the beetle state, 21—Assisted by man, 21—Tendency to migrate in swarms, 23.	•
It spreads, but does not travel in the sense of leaving one district for	
another	23

V	
RILEY, C. V.—Continued.	
THE COLORADO POTATO-BEETLE-Continued.	
Area invaded by it	9
Causes which limit its spread	2
Intense heat in the South, 25—Excessive dryness in the mountains, 26.	
How it affected the price of potatoes	9
The modification it has undergone	2
Its natural history	2
First made known in 1863, 28—The female capable of laying 1,000 eggs, 29—Three broods a year, 28.	
Ita poisonous qualities	20
Exhalations from the crushed bodies injurious, 29.	
Its food-plants	30
The number increases with each year, 32—Varieties of potatoes	
preferred, 33.	
The beetle cats as well as the larva	33
Its natural enemies	34
Birds which feed upon it, 35–36—Domestic fowls, 36—Reptiles, 36—	
Spiders and ites, 36-38—True insects, 39—Rust-red social wasp,	
40—Lady-birds, 40—43—Ground-beetles, 44—45—Rove-beetles, 46—	
Blister-beetles, 46—Soldier-bugs, 47-51—Tachina-fly, 52—Asilus-	
flies, 53.	
Remedies	54
· · · · · · · · · · · · · · · · · · ·	
Mechanical means of destruction, 55—Pincers for, 56—Sun-sealding, 56—Horse-machine, 57—Machines for collecting, 58, 59—Poi-	
sonous applications to the plant, 60—Paris green, 61—Different modes of using Paris green, 62–65—Other poisonous applications	
tested, 66—Patent poisons, 68.	
The use of Paris green	69
Its influence on the plant, 70—Its influence on the soil, 71—Its in-	•
fluence on man indirectly through the soil or through the plant, 74.	
Bogus experiments	75
Alarm about the insect abroad	76
Unnecessary prohibition of traffic in American potatoes, 77—How	
the insect will most likely get to Europe, 78—The chances of its	
getting there, 79-82—Could it become acclimated there 1, 82.	
Nomenclature	83
The bogus Colorado potato-beetle, Doryphora juncta	85
It has always existed east of the Mississippi, 85—It never attacks	
the cultivated potato, 85—Easily confounded with its potato-	
feeding congener, 86-How the two differ, 86-88.	
OTHER INSECT FOES OF THE POTATO.	
THE STALK-BORER, Gortyna nitela	90
Habits, 90—Remedy, 91.	
THE POTATO STALK-WEEVIL, Trichobaris trinotata	92
Habits, 92—Remedy, 93.	
THE POTATO OR TOMATO WORM, Protoparce celeus	93
Habits, 94—Remedies, 95—Parasites, 96.	
BLISTER-BEETLES	96
The striped blister-beetle, Epicauta vittata	97
The ash-gray blister-beetle, Macrobasis unicolor	98
The black-rat blister-beetle Macrobasis unicolor	99

1558.	RILEY, C. V.—Continued.	
	BLISTER-BEETLES-Continued.	
	The black blister-beetle, Epicauta pennsylvanica	99
	The margined blister-beetle, Epicauta cinerea	99
	Remedies, 100.	
	THE THREE-LINED POTATO-BEETLE, Lema trilineata	100
	Habits, 101, 102—Remedies, 102.	
	THE CUCUMBER FLEA-BEETLE, Crepidodera cucumeris	102
	Habits, 103—Remedies, 103.	
	THE CLUBBED TORTOISE-BERTLE, Coptocycla clarata	103

- 1559. RILEY, C. V. Gall-insects. <Johnson's New Universal Cyclopædia, 1876, v. 2, pp. 412-416, 16 figs. S.-b. No. 10, pp. 76-80. Definition of the term "gall-insects"; classification and habits of the same; mention by name and figures of typical species and galls made by them; dimorphism and metagenesis of Cynips.</p>
- 1560. RILEY, C. V. Gall-nuts. < Johnson's New Universal Cyclopædia, 1876, v. 2, p. 417, 1 fig. S.-b. No. 10, p. 81.

 Definition of "gall-nuts" formed by Cynips gallæ-tinctoriæ on twigs of Quercus infectoria; local origin, chemical composition, use in the arts.
- 1561. RILEY, C. V. Galls. < Johnson's New Universal Cyclopædia, 1876, v. 2, pp. 418-419, 2 figs. S.-b. No. 10, pp. 82-83.

 Definition of "galls;" their variety of form, texture, and location; their nature and source.
- 1562. RILEY, C. V. Locust prospects. <Colman's Rural World, 3

 January, 1877. S.-b. No. 14, pp. 57-58; 58-59.

 Explains means of prognosticating the advent of Calopienus spretus in the spring of 1877; recommends measures of precaution and the procurement

spring of 1877; recommends measures of precaution and the procurement of information concerning means against these insects, and gives assurance that the invasion of the locusts will be but temporary.

- 1563. RILEY, C. V. Bots. <Sci. Amer., 6 January, 1877 [v. 50], n. s.,
 v. 36, pp. 9-10. Reprint: —S.-b. No. 14, pp. 235-236. <Lancaster Farmer, 15 September, 1877, v. 9, p. 142. S.-b. No. 14,
 p. 129. <Colman's Rural World. S.-b. No. 14, p. 100.
 - Habits, ravages, and means against Gastrophilus equi, Cophalomyis [= Estrus] ovis, and Hypoderma bovis.
- 1564. [RILEY, C. V.] Academy of Science. Brilliant and profound address of Prof. C. V. Riley. The splendid record of the Academy for the past year. Reports of officers and committees and election of managers for the ensuing year. <St. Louis Times, 16 January, 1877, v. —. S.-b. 14, pp. 91-96. Reprint: <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 238-254 Proc. Separate: <St. Louis, 1877, 16 pp. Translation: <Anzeiger des Westerns, 16 January, 1877, v. 43, p. 3. S.-b. No. 14, pp. 50-54.
 - Review of progress made in science in North America, and especially by the Academy of Science of St. Louis in 1876; remarks on entomological observations and publications, and especially on Doryphora decembineata, Phylloxera vastatrix, and Caloptenus spretus.

- 1565. RILEY, C. V. Is this a grasshopper year? Prof. Riley's opinion concerning the prospect for bugs. It all depends on the kind of weather we have during February. <8t. Louis Daily Globe Democrat, 7 February, 1877, v. 2, No. 263, p. 3. S.-b. No. 14, pp. 69-70. Reprint: <Industrialist [Manhattan, Kans.] 17 February, 1877, v. 2, pp. 1, 4. S.-b. No. 14, p. 49. See: <Colman's Rural World, 1877. S.-b. No. 1, pp. 59-00.
 - Replies to questions as to the likelihood of the hatching of eggs of Caloptenus spretus in the spring of 1877; the degree of development attained before winter; possibility of the resumption of development after it has once been arrested.
- 1566. [RILEY, C. V.] Are the locusts hatching? Mistaken identity. <Colman's Rural World, 14 February, 1877, 2 figs. S.-b. No. 14, p. 62.
 - Modified extract from 8th Ann. Rept. State Ent. Mo., May, 1876, pp. 149-150, Tragocophala [=Chortophaga] viridifasciata and Tettix granulatus mistaken for Caloptonus spretus; geographical distribution of the first; hibernation and colors of the two former; figures of both.
- 1567. Riley, C. V. Condition of locust eggs: Inquiries answered. <Colman's Rural World, 21 February, 1877. S.-b. No. 14, pp. 67-68; 68.
 - Replies to inquiries as to the degrees of development attained by eggs of Caloptenus spretus submitted for examination.
- 1568. RILEY, C. V. Tarred paper for fruit trees. <Colmau's Rural World, 7 March, 1877. S.-b. No. 14, p. 61.
 - Critical review of article by E. Gaylord; the inclosure of trunks of fruit trees in tarred paper serviceable as a protection from the sun, rabbits, mice, and borers.
- 1569. RILEY, C. V. Insect on the grape. Cardener's Mo. and Hortic.,
 March, 1877, v. 19, p. 90.
 - Varieties of grape attacked by Desmis maculalis; means against the same.
- 1570. RILEY, C. V. Ninth annual report on the noxious, beneficial, and other insects of the State of Missouri, made to the State Board of Agriculture, pursuant to an appropriation for this purpose from the legislature of the State. <12th Ann. Rept. State Board of Agric. for 1876, March, 1877, pp. 7+129+3, 33 figs. Separate: <Jefferson City, Mo., March, 1877, pp. 7+129+3, 33 figs.

NOXIOUS INSECTS.

3

THE GOOSEBERRY SPAN-WORM, Enfitchia ribearia

Its natural history, 3—Most destructive gooseberry insect in Missouri, 3—Generic nomenclature, 3—Characters of the moth, 4—
Description of the egg, 4—Where the eggs are laid, 4—The in-

23

27

29

32

1570. RILEY, C. V.—Continued.

worms, 7.

THE GOOSEBERRY SPAN-WORM—Continued.
sect single-brooded, 5—How it spreads, 5—A native species, 5—
Its past history, 5—It prefers the gooseberry to the currant, 6—
The moth is closely imitated by one which greatly differs structurally, 6—Parasites, 6—Remedies, 7—Other currant span-

The imported current-worm, Nematus ribesis

Belongs to the "false-caterpillars," 7—The different specific names it has received, 7—An imported species, 8—Its introduction and spread, 8—Independently imported at several eastern points, 9—Its natural history, 9—How the eggs are laid, 10—Nature and habits of the worm, 11—Characters of the parent flies, 12—Preventive measures, 13—Remedies, 13—White hellebore the best, 13—How best used, 14-15—The worm is not poisonous, 16—Natural enemies, 17—It furnishes a forcible example of arrenotoky, or the power of producing male offspring without impregnation, 18—Results of parthenogenesis in different insects, 18—It also furnishes an interesting instance of defunctionation of special parts, 19—The saw of the female imperfect compared with others, 20—Evolutionary bearings of this fact, 21—Descriptive, 21—Variation in the antennæ and wing veins, 22.

THE NATIVE CURRANT-WORM, Pristiphora grossulariæ

Wherein it differs from the imported species, 23—Its habits, 24—
Where the eggs are laid, 25—How the winter is passed, 25—Its occurrence in Missouri, 26—Remedies, 26—Descriptive, 26.

A new enemy to wheat, 50—First complaint of it in the East, 51—First appearance in Kansas, 51—Habits and natural history, 52—The egg differs from that of the army-worm, 53—Wherein the worm is distinguished from its destructive congener, 54—Two broods each year, 54—Natural enemies, 54—Remedies, 55—Descriptive, 55.

50

57

THE ROCKY MOUNTAIN LOCUST, Caloptenus spretus.

It continues to interest the people of the West, 57—Previous opin-

ions justified, 57—The invasion of 1876, 59—Few in British America, 59-Condition of things in Montana, 59-In Wyoming, 59in Dakota, 59-In Minnesota, 60; locusts and alkali soil, 61; good done by Governor Pillsbury, of Minnesota, 61-In Colorado, 62-In Iowa, 63-In Nebraska, 64-In Kansas, 65-In Missouri, 66-Flights in opposite directions at the same time, 66-Counties in Missouri that were overrun, 67-Red-legged locust troublesome in east Missouri, 68-Detailed reports from counties in Missouri, 68-Andrew County, 68-Atchison County, 68-Barry County, 68—Barton County, 69—Bates: County, 69—Benton County, 69—Buchanan County, 69—Cass County, 69—Cedar County, 70-Caldwell County, 70-Clay County, 70-Dade County, 70-De Kalb County, 70-Gentry County, 70-Greene County, 71-Harrison County, 71-Henry County, 71-Hickory County, 71-Holt County, 71-Jasper County, 72-Jackson County, 73-Johnson County, 73-Lafayette County, 73-Lawrence County, 73-McDonald County, 74-Newton County, 74-Nodaway County, 74-Pettis County, 74-Platte County, 74-Polk County, 75-Ray County, 75-Saint Clair County, 75-Vernon County, 75-In Indian Territory, 76-In Texas, 76-In Arkansas, 76—Destination of the departing swarms of 1875, 77—They reached into British America, 78—Source of the swarms of 1876, 79—Eastern line reached, 80—Rate at which the insects spread, 80-Direction of flight, 81-Influence of wind in determining the course of locust swarms, 81-Locust flights east of the Mississippi, 81—Geographical range of species, 82—Causes which limit the spread of the Rocky Mountain locust, 83-Flights of Acridium americanum, 84-Does the female of the Rocky Mountain locust lay more than one egg-mass?, 85—How the eggs are laid, 86-Philosophy of the egg-mass, 87-How the young locust escapes from the egg, 88-How it escapes from the ground, 90-Additional natural enemies, 91—Animals which destroy the eggs, 91—The Anthomyia egg-parasite, 92—The common flesh-fly, 95— Other undetermined enemies of the eggs, 96-Insects which destroy the active locusts, 98-Experiments with the eggs and conclusions therefrom, 99-Experiments to test the effects of alternately freezing and thawing, 99-Experiments to test the influence of moisture upon the eggs, 104-Experiments to test the effects of burying at different depths and of pressing the soil, 104—Experiments to test the effects of exposure to the free air, 104- The Omaha conference, 106-Remedies and suggestions, 108-Des ruction of the young or unfledged locusts, 108-Protec-

THE ROCKY MOUNTAIN LOCUST-Continued.

tion of fruit-trees, 110—Legislation, 111—Act passed by the Missouri legislature, 111—Acts passed by the Kansas legislature, 112—Act providing for the destruction of locusts in Minnesota, 114—Area in which eggs were laid, 116—Condition of eggs, 117—Temperature of the winter of 1876—77, 120—Prospects for 1877, 121.

INNOXIOUS INSECTS.

- 1571. RILEY, C. V. Important observations on the Rocky Mountain locust or grasshopper pest of the West. <Sci. Amer., 28
 April, 1877 [v. 50], n. s., v. 36, pp. 260-261, 5 figs.
 - Observations on the eag-laying habits of Caloptenus spretus; figures egg, eggmasses, method of oviposition, and female anal characters.
- 1572. RILEY, C. V. Experiments with locust eggs and conclusions therefrom. <Sci. Amer., 5 May, 1877 [v. 50], n. s., v. 36, pp. 276-277. S.-b. No. 14, p. 101.
 - Experiments upon the vitality of eggs of Caleptenus spretus; effects of alternately freezing and thawing, of exposure to great moisture or to the free air, and of burying them at different depths.
- 1573. RILEY, C. V. Prof. Riley's report to the Governor of Kansas:

 The grasshopper question: Interesting information. <Commonwealth [Topeka, Kans.], 12 May, 1877, No. 2500, p. 2. S.-b.

 No. 14, pp. 63–67. Reprint: <St. Louis Daily Globe-Democrat,
 14 May, 1877, v. 2, No. 359, p. 3. S.-b. No. 14, pp. 70–72.

 Notice: < 1bid., p. 4. S.-b. No. 14, p. 70. See: <Kansas
 Farmer, 16 May, 1877. S.-b. No. 14, p. 59.
 - Reply to letter of Governor G. T. Anthony; area within Kansas in which eggs of Caloptenus spretus were laid in 1876; the degree to which the young from these eggs had perished or were likely to commit ravages in the summer of 1877; causes of the destruction of a great proportion of the young locusts; means of completing this destruction.
- 1574. RILEY, C. V. The strawberry leaf-roller, Anchylopera fragariæ. <Gardener's Mo. and Hortic., May, 1877, v. 19, pp. 143-144, fig. S.-b. No. 14, p. 45.
 - Reprint, with additional introduction and note, from 1st Ann. Rept. State Ent. Mo., March, 1869, pp. 142-143, fig. 80. See No. 1059 for synopsis of contents.

- 1575. [RILEY, C. V., et al.] U.S. Entomological Commission. Circular No. 1. [Riley, Packard, Thomas.] < Washington: 1877, 80, 4 pp.
 - Queries regarding the migrations, appearances, habits, and ravages of Caloptenus spretus; directions for making replies.
- 1576. RILEY, C. V. U. S. Entomological Commission. Circular No. 2. <Washington: 1877, 8°, 4 pp.
 - Plans of work; request for information regarding the natural history, insect enemies, and parasites of *Caloptenus spretus* and other locusts; means against the same.
- 1577. [RILEY, C. V., et al.] Bulletin of the United States Entomological Commission. Destruction of the young or unfledged locusts. No. 1. [Riley, Packard, Thomas.] < Washington: 1877, 8°, 12 pp. See: < Sci. Amer., 2 June, 1877 [v. 50], n. s., v. 26, p. 344.
 - Enumeration of means of destroying young or unfiedged locusts; quotations of laws passed by States of Missouri, Kansas, and Minnesota to provide for the destruction of locusts and their eggs.
 - Supplement to Bulletin No. 1, 2 pp. Summary of means for the destruction of young locusts.
- 1578. [RILEY, C. V., et al.] Bulletin of the United States Entomological Commission. On the natural history of the Rocky Mountain locust, and on the habits of the young or unfledged insects as they occur in the more fertile country in which they will hatch the present year. No. 2. [Riley, Packard, Thomas.] < Washington: May, 1877, 8°, 15 pp., figs. 1-11, map.
 - Description, oviposition, hatching, transformations, and habits of Caloptenus spretus; map of the country that will suffer most severely, showing the eastern limit of injury the present year.
- 1579. RILEY, C. V. The grape leaf-folder. <Journ. and Farmer, 14
 June, 1877. S.-b. No. 14, p. 133.
 - Description of larva, imago, habits, ravages, distribution, seasons, and food-plants of Desmia maculalis.
- 1580. RILEY, C. V. The rascal leaf-crumpler. <Journ. and Farmer, 14 June, 1877. S.-b. No. 14, pp. 133-134.
 - Description of larva, larva-case, imago, habits, ravages of, distribution, seasons, food-plants, and means against Phycita nebulo [= Acrobasis indiginella].
- 1581. RILEY, C. V. The Hessian-fly. <Journ. and Farmer, 14 June, 1877. S.-b. No. 14, p. 135.
 - Description of eggs, larva, puparium, and image of Cocidomyia destructor; civil history, seasons, habits, ravages of and means against it; habits of Semiotellus [= Merisus] destructor.
- 1582. RILEY, C. V. Locust prospects. <Sci. Amer., 16 June, 1877, [v. 50], n. s., v. 36, p. 369.
 - Causes of the destruction of a great proportion of the young of Caloptenus spretus; a general and disastrous invasion improbable for some years.
- 1583. [RILEY, C. V.] Insect enemies. <N. Y. Tribune, 16 June, 1877. S.-b. No. 14, pp. 234-235.
 - Apple-borers; answer to letter of V. B. P.; young apple-trees killed by Xyleborus obesus.

- 1583. RILEY, C. V.—Continued.
 - Rose-chafer; answer to letter of I. M. H.; means against Macrodactylus subspinosus.
 - Bark-louse; answer to letter of J. L. K.; seasons and ravages of and means against Mytilaspis pomicorticis [= pomorum].
 - Flea-beetle; answer to letter of J. E. R.; habits, ravages, and means against Graptodera [= Haltica] chalybea.
- 1584. RILEY, C. V. The Hellgrammite. <Sci. Amer., 23 June, 1873 [v. 50], n. s., v. 36, pp. 392-393, 3 figs.
 - Structure and habits of Corydalus cornutus, description and figures of eggmasses, eggs, larvæ, pupa, and images; adaptations of sexual structure to conditions; figure and description of eggs of Belostoma grande [= americanum].
- 1585. RILEY, C. V. The 'hopper in Iowa. Report by Professor Riley, Chief of the U. S. Entomological Commission. <Chicago Daily Tribune, 4 July, 1877, v. 32, p. 3. S. b. No. 14, pp. 112–114. Extract: <N. Y. Tribune, 4 July, 1877. S. b. No. 14, p. 128.
 - Report to the governor of Iowa as to the observed and anticipated ravages of Caloptenus spretus in southwestern Iowa in 1877; the young locusts mostly destroyed by the weather; their natural enemies and the efforts of man; localities infested; recommends diversification of crops; summary of means against the young locusts; claims for the first recommendation of proper ditching; description of apparatus for catching and killing locusts.
- 1586. RILEY, C. V. Strawberry worm and remedy. <N. Y. Tribune, 18 July, 1877. S. b. No. 14, p. 231.
 - Description of egg, larva, and imago, habits of and means against Emphytus [= Harpiphorus] maculatus.
- 1587. RILEY, C. V. Fighting the Hessian-fly. <N. Y. Tribune, 18
 July, 1877. S.-b. No. 14, pp. 232-233. Reprint: <Colman's
 Rural World, 5 December, 1877. S.-b. No. 14, pp. 268-269.
 - Habits, seasons, parasites, introduction, spread, and ravages of and means against Cooldomyia destructor; description of eggs and larva.
- 1588. RILEY, C. V. The grasshopper. Considered practically and scientifically with a retrospective and prospective glance at his history. <Daily Rocky Mountain News [Denver, Colo.], 1 August, 1877, v. 18, p. 4. S.-b. No. 14, pp. 81-84; 110-112. Reprint: <Colorado Farmer, 2 August, 1877, v. 9, No. 31, p. 4. <Chicago [Ill.] Daily Tribune, 7 August, 1877, v. 32, p. 7. S.-b. No. 14, pp. 84-86.</p>
 - Observations on the past and present conditions of Caloptenus spretus in 1877 in the regions invaded by it; work of its enemies and parasites; complication of the locust problem in Colorado owing to the climate of the State; prospects of future injury; means against the locusts in Colorado; work of the U. S. Entomological Commission.
- 1589. RILEY, C. V. In reference to wheat-worms. < Prairie Farmer, 11 August, 1877. S.-b. No. 14, p. 127; No. 20, p. 86.
 - Critical review of C. Thomas' "A wheat insect" (Prairie Farmer, 21 July, 1877), with correction of some obscurities in regard to Meromyza americana and Gortyna nitela.

- 1590. RILEY, C. V. Locust flights. An appeal from Prof. C. V. Riley. <Manitoba Daily Free Press, 8 September, 1877. S.-b. No. 14, p. 91.
 - Reprint of questions of Circular No. 1 of U. S. Entomological Commission, with request for answers and other information from observers in Manitoba.
- 1591. RILEY, C. V. The locusts in Kansas. <Sci. Amer., 8 September, 1877 [v. 51], n. s., v. 37, p. 164.
 - No danger of locust invasion in Kansas and adjacent States in the fall of 1877; diseased conditions of late swarms; seasons of drought and locust swarms generally followed by rainy seasons and a searcity of locusts.
- 1592. RILEY, C. V. A satisfactory grasshopper-machine. <Sci. Amer., 8 September, 1877 [v. 51], n. s., v. 37, p. 169.

 Description and figure of a machine for the killing of locusts.
- 1593. [RILEY, C. V.] Mistaken identity. <N. Y. Tribune, 12 September, 1877. S.-b. No. 14, p. 233.</p>
 - Answer to letter of Subscriber; Lema trilineata mistaken for Doryphora 10-lineata; first occurrence of the latter in Vermont in 1876.
- 1594. RILEY, C. V. Injured orchard. <N. Y. Tribune, 12 September, 1877. S.-b. No. 14, p. 233.
 - Answer to letter of G. W. T.; occurrence of numerous nests of Formicidae about the roots of orchard trees; probability that the ants are harmless; means against ants.
- 1595. [RILEY, C. V.] The stalk-borer. <N. Y. Tribune, 12 September, 1877. S.-b. No. 14, p. 233.
 - Answer to letter of Wisc.; food-plants of Gortyna mitela; description of larva.
- 1596. RILEY, C. V. Locust prospects in southwest Missouri this fall. <Journ. and Farmer, 27 September, 1877. S.-b. No. 14, pp. 236-237.
 - Ravages of Caloptenus spretus in Missouri in 1876 and prospects of same in 1877; history of locust flights in 1877.
- 1597. RILEY, C. V. The Colorado potato-beetle in Europe. German thoroughness. <Sci. Amer., 29 September, 1877 [v. 51], n. s., v. 37, p. 198. S.-b. No. 14, pp. 123-124.
 - Occurrence of *Doryphora* 10-lineata at Bremen and at Mülheim, in Germany; thorough measures taken to eradicate the pests; economic value of the measures.
- 1598. [RILEY, C. V.] The cussed red-leg. <Chicago [Ill.] Times, 29
 September, 1877. S.-b. No. 14, pp. 119-123. Reprint: <Mo. Rept. Kansas State Board of Agric., 1877, pp. 32-41. S.-b. No. 14, pp. 146-161. Abstract: <Amer. Nat., November, 1877, v. 11, pp. 663-673. <Ca. Nat. and Quart. Jour. Sci., December, 1877, v. 8, pp. 363-374.
 - Distinction between the terms locust and grasshopper; nature of varieties and species; distribution of Caloptonus spretus; laws governing its migrations and distribution; exemplification of these laws by records of flights during 1877; means against locusts; distinctions between Caloptonus spretus and C. femur-rubrum; habite and transformations of Astoma gryllaria (—the young of Trombidium sericeum [—locustarum]); habits of the latter.

- 1599. RILEY, C. V. [White-grub fungus.] <N. Y. Weekly Tribune, 4
 October, 1877. Notice: <Amer. Ent., June, 1880, [v. 3], n. s., v.
 1, p. 140.
- 1600. RILEY, C. V. On the larval characters and habits of the blister-beetles belonging to the genera *Macrobasis* Lec. and *Epicauta* Fabr.; with remarks on other species of the family *Meloidæ*. <Trans. Acad. Sci. St. Louis, 5-16 November, 1877, v. 3, pp. 544-562, figs. 35-39; figs. 1-12 of pl. 5 with 1 p. expl. of pl. Separate: <[St. Louis, Mo., 1877], pp. 1-19, figs. 35-39; figs. 1-12 of pl. 5 with 1 p. expl. of pl. Abstract: <Sci. Amer., 1877 [v. 51], n. s., v. 37; 1 December, p. 346; 15 December, p. 373; 404 il. S.-b. No. 14, pp. 272-275. <Ent. Mo. Mag., January, 1878, v. 14, pp. 169-175. Reprint, with changes and omissions: <Amer. Nat., 1878, v. 12; April [19 March], pp. 213-219, figs. 1-2; May [22 April], pp. 282-290, figs. 3-5, pl. 1 with 1 p. expl. of pl. Separate: <Boston: 1878, pp. 213-219; 282-290; pl. 1 with 1 p. expl. of pl.
 - Views of earlier authors as to the habits of meloid larvæ; nature of the newly hatched larva (triungulin); natural history of Melos and Sitaris, with reference to the literature, descriptions of the triungulina, especially of M. barbarus?; figures of the several stages; nature of hypermetamorphosis; food-habits and breeding habits of Macrobasis and Epicauta, especially of E. vittata; summary of what is known of the larval habits of other genera of Meloidæ; descriptions and figures of all the earlier stages of E. vittata; descriptions of eggs and triungulins of E. cinerea, E. pennsylvanica, and Henous confertus.
- 1601. RILEY, C. V. On a remarkable new genus in *Meloidæ* infesting mason-bee cells in the United States. <Trans. Acad. Sci. St. Louis, 16 November, 1877, v. 3, pp. 563-565, fig. 40; fig. 13 of pl. 5, with 1 p. expl. of pl. Separate: <[St. Louis, Mo., 1877], pp. 20-22, fig. 40; fig. 13 of pl. 5, with 1 p. expl. of pl. Abstract: <Amer. Nat., April [19 March], 1878, v. 12, pp. 218-219.
 - Description of imagos, ultimate stage of second larva, and coarctate larva of Hornia minutipennis, n. g. et n. sp., parasitic in cells of Anthophora sponsa [=abrupta]; figures 3 imago and coarctate larva of the same; illustrates the stages of degradation in tarsal claws of several genera of Meloidæ; synoptic table of the North American genera of Meloini.
- 1602. RILEY, C. V. Additional notes on Megathymus yucca. <Trans. Acad. Sci. St. Louis, 16 November, 1877, v. 3, pp. 566-568. Separate: <[St. Louis, Mo., 1877], pp. 23-25.
 - Supplementary to No. 1465; habits, seasons, and number of molts of the larva; description of larva in second and third stage and of variations in the imago; description of coloradensis n. var.; Egiale cofaqui a variety of Megathymus yuccæ; Erynnis alceæ bores in stems of Malva sylvestris in Europe; Elaphidion tectum and Soyphophorus yuccæ in stems of Yucca.
- 1603. RILEY, C. V. Further remarks on *Pronuba yuccasella* and on the pollination of *Yucca*. <Trans. Acad. Sci. St. Louis, 16 No-

- 1603. RILEY, C. V.—Continued.
 - vember, 1877, v. 3, pp. 568-573. Separate: <[St. Louis, Mo., 1877], pp. 25-30. Translation: <Stett. Ent. Zeit., 1878, Jahrg. 39, pp. 377-382.
 - Critical review of remarks on *Pronuba* by Chambers, Zeller, and Boll; *Hyponomeuta quinquepunotella* mistaken for *Pronuba yuccasella*; variations in the former; uniformity in the latter; behavior of *Pronuba* in flowers of *Yucca*; question of the method of fertilization of *Yucca*.
- 1604. RILEY, C. V. On the differences between Anisopteryx pometaria,
 Harr. and Anisopteryx ascularia W. V., with remarks on the
 genus Paleacrita. <Trans. Acad. Sci. St. Louis, 16 November20 December, 1877, v. 3, pp. 573-577. Separate: <[St. Louis,
 Mo., 1877], pp. 30-34.
 - Characters of Anisopteryx ascularia show that it is congeneric with A. pometaria; criticism of Packard's Monograph of Geometrida; structural characters separating Anisopteryx and Paleacrita.
- 1605. RILEY, C. V. Wheat rust and Hessian fly. <N. Y. Tribune,
 19 December, 1877. S.-b. No. 14, p. 235.
 - Means against Puccinia graminis; P. graminis and Ecidium berberidis are alternate generations of one species.
- 1606. RILEY, C. V. A new oak-gall on acorn cups. <Trans. Acad. Sci. St. Louis, 20 December, 1877, v. 3, pp. 577-578.

 Description of gall of Cymips querous-glandulus n. sp. [p. 578] on cupules of
 - all species of *C. fecundatrix* in Europe, and of *C. querous-frondosa*; occurrence of galls of *C. fecundatrix* in Europe, and of *C. querous-frondosa*; occurrence of galls of *C. [=Amphibolips]* querous-prunus on cupules of acorns.
- 1607. [RILEY, C. V.] [Maggots in sauce.] <N. Y. Tribune, 26 December, 1877. S.-b. No. 14, p. 234.
 - Larvæ of *Drosophila* sp. in canned sauce; *D. nigricornis* common in similar substances.
- 1608. [RILEY, C. V.] [Round-headed apple-tree borer.] <N. Y. Tribune, 26 December, 1877. S.-b. No. 14, p. 234, Ravages of and means against Saperda bivittata [= candida].
- 1609. RILEY, C. V. Entomological notes. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 217-218 Proc. See: <Amer. Nat., October, 1876, v. 10, p. 635.
 - Generic resemblance of Carpocapsa saltitans to C. pomonella; correction of vernacular name of Cicada [= Tibicen] septendecim; occurrence of the same in Virginia in 1876; yearly development of C. [= T.] tredecim; Sericaria mori reared for five years on Maclura aurantiaca with increased vigor and healthfulness; Salix nigra stripped by larvæ of Vanessa antiopa; habits of the same.
- 1610. RILEY, C. V. Entomological notes. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 219 Proc. See: <Colman's Rural World, 1876. S.-b. No. 10, p. 165. <N. Y. Tribune, 1876. S.-b. No. 9, p. 215. <Amer. Nat., October, 1876, v. 10, p. 635.
 - Occurrence and ravages of Leucania albilinea in Kansas; ectoparasitism of Uropoda americana on Doryphora decemlineata; list of vertebrate enemies of

- 1610. PILEY, C. V.—Continued.
 - D. decemlineata; occurrence of D. decemlineata in New Hampshire; its ravages along the Atlantic coast; see No. 1505.
- 1611. RILEY, C. V. Centennial insects. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 220-221 Proc. List and characterization of insects observed injuring exhibits in the Centennial Exposition.
- 1612. RILEY, C. V. Parasites on eggs of Caloptenus spretus. < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 226 Proc.
 - A large proportion of locust eggs throughout the country are being destroyed by parasites, predaceous insects, and other causes; precautions against threatened injury; discovery of four new enemies of locust eggs.
- 1613. RILEY, C. V. Locust flights east of the Mississippi. < Trans.

 Acad. Sci. St. Louis, December, 1877, v. 3, pp. 226-228 Proc.

 Locust swarms in Ohio, Illinois, Georgia, and South Carolinia not composed of Caloptenus spretus; causes which limit the eastward flight of C. spretus.
- 1614. BILEY, C. V. Geographical range of species. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 230-233 Proc.</p>
 Range of species in our time affected only by the interposition of human influence; the limits of this range definitely established by classified knowledge, and impassable, except by man's assistance; Caloptenus spretus can not cleange its habits; the difference in the susceptibility of different
- 1615. RILEY, C. V. [Anticipated locust injury next summer.] < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 236 Proc.

 Abundance and advanced state of development of eggs of Caloptenus spretus in the States invaded in 1876; probable abundance and ravages in 1877.

species to change inexplicable.

- 1616. RILEY, C. V. [Japanese mode of packing silk-worm eggs.]

 <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 236

 Proc.
 - Mode in which the eggs of Sericaria mori are packed in Japan for transportation
- 1617. RILEY, C. V. Locust experience. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 267 Proc. See: <Colman's Rural World, 1877. S.-b. No. 14, p. 265.
 - Northern limit of the permanent breeding-grounds of Caloptenus spretus; confirmation of views regarding laws governing locust invasions; discovery of a new law.
- 1618. RILEY, C. V. Mite transformations. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 267-268 Proc. See: <Colman's Rural World, 1877. S.-b. No. 14, p. 265.</p>
 - Habits and transformations of Trombidium sericeum [= T. locustarum]; Astoma gryllaria an immature form of the same.
- 1619. RILEY, C. V. [Mygale hentzii and Pepsis formosa.] < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 269 Proc.
 - The bite of Mygale hentsii not so deadly as is generally supposed; habits of Popsis formosa in preying on the spider.

1620. RILEY, C. V. [Ravages of Termes flavipes.] < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 269 Proc.

Note to communication of R. D. Grant on some ravages of Tormes flavipes.

1621. RILEY, C. V. On the oviposition of Saperda bivittata Say. < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 269-270 Proc. See: < Colman's Rural World, 28 November, 1877. S.-b. No. 20, p. 6.

Description of egg and manner of oviposition of Saperda bicittata [= candida].

1622. RILEY, C. V. On migratory butterflies. <Trans. Acad. Sci. St. Louis, December, 1877, v. 3, pp. 273–274 Proc. See: <Valley Naturalist, January, 1878, v. 1, p. 2.

Phenomena and causes of the migrations of butterflies, especially of *Danais* archippus.

- 1623. RILEY, C. V. [Phylloxera and grape-rot.] < Trans. Acad. Sci. St. Louis, December, 1877, v. 3, p. 275 Proc. See: < Colman's Rural World, 28 November, 1877. S.-b. No. 14, p. 268.
 - Disagreement with views of A. J. Cook as to there being any connection between the work of *Phylloxera vastatrix* and the ordinary grape-rot.
- 1624. RILEY, C. V. The periodical Cicada. <Western Farmer's Almanac for 1878, 1877, p. 48. S.-b. No. 14, p. 138. Reprint: <Colman's Rural World, 28 November, 1877. S.-b. No. 14, pp. 265-266.

Regular periodicity of the appearance of Cicada [= Tibicon] septendecim and C. [= T.] tredecim; popular description and natural history of the same; chronology of twenty-two different broods, geographically classified.

1625. RILEY, C. V. The locust plague in the United States: being more particularly a treatise on the Rocky Mountain locust, or so-called grasshopper, as it occurs east of the Rocky Mountains, with practical recommendations for its destruction. <Chicago: Rand, McNally & Co., 1877, 236 pp., 3 pl., 42 figs. Extract: <Manitoba Standard, 1877. S.-b. No. 14, pp. 97-98. Notice: <Cultivator and Country Gentl., 30 August, 1877, v. 42, p. 557. <Gardener's Mo. and Hortic., October, 1877, v. 19, p. 317. <Ent. Mo. Mag., October, 1877, v. 14, p. 118.

TABLE OF CONTENTS.

27—Purely an American insect, 28.

Introduction	
Chapter I:	
Characters of the species	1
Classificatory position of the Rocky Mountain locust, 13—How it	
differs from other species, 14—Easily confounded with the com-	
mon red-legged locust, 14—Detailed descriptions of both, 15—	
Its still closer resemblance to the Atlantic migratory locust, 22—	
Characters of this last, 22—Species vs. variety and race, 23—Com-	
parisons of these three closely-allied species in their early stages,	
26-A green variety of the Rocky Mountain locust not infrequent,	

4 400	RILEY.	\sim	**	\sim	 •

	•		
CTT		TT.	

Chronological histery

The locust plague in the "Old World," 29—Extent of its injuries, 30—Migratory species in Europe, Asia, and Africa, 30—The ravages of the locust in America, 31—Its earliest visitations, 31—Injuries on the Pacific coast, 32—Injuries east of the Rocky Mountains, 33—Invasions of 1818—'19, 33—Of 1845—'49, 34—Of 1855, 34—Of 1856, 35—Of 1857—'67, 35, 36—Of 1866, 36—Damage the following year, 37—The invasion of 1873, 38—That of 1874, 39—Why so disastrous, 41—General outlook in the spring of 1875, 42—Severity of the injuries from the young insects that year, 43—Destitution that prevailed, 44—Amount of loss sustained, 45—Destination of departing swarms of 1875, 47—The invasion of 1876, 49—Eastern

CHAPTER III:

Native home and geographical range of the species east of the mountains

limit reached, 53-Omaha conference, 53.

Source of the devastating swarms that reach into the Mississippi Valley, 55—Their origin in the extreme Northwest country lying east of the mountains, 56—Cause of their emigration, 57—Difference between summer and fall swarms, 58—The species not at home in the Mississippi Valley, 62—Not permanent or able to perpetuate itself there, 62—Conditions which prevent such permanence, 63—Not likely to do serious harm east of the ninety-fourth meridian, 65—Reasons why, 65.

CHAPTER IV:

CHAPTER V:

CHAPTER VI:

summer and fall, 108—No evil without some compensating good, 108—Changes that follow the locusts, 109—The prevalence of large green worms, 110—The sudden appearance of a peculiar grass, 110.

55

85

69

CHAPTER VII:

Natural enemies of the Rocky Mountain locust Birds and other vertebrate animals, 113—The good offices of birds, 113-Invertebrate animals, 114-Animals that attack the eggs, 115-The silky mite, 115-The Anthomyia egg-parasite, 118-The common flesh-fly, 123-Undetermined larvæ, 123-Ichneumonflies, ground-beetles, click-beetles, and myriapods, 126-Insects that destroy the active locust, 126-The locust-mite, 128-The anonymous Tachina-fly, 131-Yellow-tailed Tachina-fly, 134-Flesh-fly, 135.

CHAPTER VIII:

Practical considerations How best to prevent locust injuries, 139—Encouragement of natural enemies, 139-Destruction of the eggs, 139-Experiments , with the eggs and conclusions drawn therefrom, 140—Effects of alternately freezing and thawing, 141-Influence of moisture, 143-Exposure to the free air, 148-Burying at different depths, and pressure of the soil, 149-Table of temperatures, 152-Harrowing in the fall, 153-Collecting the eggs, 153-Plowing, how most effectual, 153-Irrigation, tramping, 154-Destruction of the young or unfledged locusts, 155-Burning, 155-Crushing, 157-Trapping, 157-Ditching and trenching, 161-Catching, 161-Different contrivances for this purpose, 162-Use of destructive agents, 162-Coal-oil pans, 163-The use of coal-tar, 164-The protection of plants by special applications, 165-The best means of protecting fruit and shade trees, 166-Sulphur fumes and smudges, 167-Destruction of the winged locusts, 167-Preventive measures, 169-Suggestions that may prove of service, 170-Use of hogs and poultry, 171-The Signal Service, 172-Military aid, 173-Diversified agriculture, 174-Organized effort, 176-State legislation, 176-Missouri locust law, 177-Kansas locust laws, 178, 179-Minnesota locust law, 180-Nebraska locust law, 184—How to avert locust invasions, 186.

CHAPTER IX:

Ravages of locusts east of the Mississippi..... Migratory locusts in the eastern States, 187-The Atlantic locust, 187-Injury from other non-migratory species, 190-Often a serious matter during hot, dry seasons, 193-Locust flights in Illinois in 1875, 195-The species of which the swarms were composed, 197-Locust flights east of the Mississippi not composed of the Rocky Mountain species, 201-Necessity of discriminating between species, 201.

CHAPTER X:

General considerations Nomenclature, 207-Locust vs. grasshopper, 207-Prairie fires vs. locust ravages, 209-Fasting and prayer, 213-Not a divine visitation, 216-Influence of the wind in determining the course of locust flights, 216-Locusts as food for man, 217.-Unnecessary

1626. RILEY, C. V. On an extensile penetrating organ in a gamasid mite. < Proc. Amer. Assoc. Adv. Sci. for 1876, 1877, v. 25, pp. 273-275, 1 fig. See: <Ca. Ent., September [October], 1876, v. 8, p. 180.

alarm caused by comparatively harmless species, 227.

139

113

187

- 1626. RILEY, C. V.—Continued.
 - Habits and description of *Uropoda regetans* and *U. americana* n. sp.; nature of the adhesive filament of these mites; structure of the supposed maxillary penetrating organ in *Uropoda*.
- 1627. RILEY, C. V. On the curious egg-mass of Corydalus cornutus, Linn., and on the eggs that have hitherto been referred to that species. <Proc. Amer. Assoc. Adv. Sci. for 1876, 1877, v. 25, pp. 275-279, 1 fig. Reprint: <9th Ann. Rept. State Ent. Mo., March, 1877, pp. 125-129, figs. 30-33. See: <Ca. Ent., October [November], 1876, v. 8, pp. 181-182. See No. 1570 for synopsis of contents.
- 1628. RILEY, C. V. Phylloxera and grape-rot. <N. Y. Tribune, January, 1878. S.-b. No. 19, pp. 193; 224. Reprint: <Colman's Rural World, 1878. S.-b. No. 14, pp. 266-267. Extract: <Gardener's Mo. and Hortic., July, 1879, v. 21, pp. 213-214. S.-b. No. 23, pp. 118; 143-145. <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, pp. 237-238. See: <Gardener's Chronicle, 9 August, 1879. S.-b. No. 26, p. 15. See No. 1721 for synopsis of contents.
- 1629. RILEY, C. V. Address by Prof. C. V. Riley, retiring, before the St. Louis Academy of Science. <St. Louis Republican, 8 January, 1878.
 S.-b. No. 14, p. 269. Reprint: <Trans. Acad. Sci. St. Louis, 1880, v. 4, pp. 1-6 Proc.
 - Review of progress made in science in 1876; success of the investigation into the Rocky Mountain locust scourge.
- 1630. RILEY, C. V. New facts about the round-head apple-tree borer.
 <N. Y. Weekly Tribune, 20 February, 1878. S.-b. No. 19, pp. 193-194. Reprint: <Colman's Rural World, 20 March, 1878.
 S.-b. No. 14, pp. 267-268. Reprint, with slight omission: <Ihid., 12 March, 1879. S.-b. No. 23, pp. 101-102. See: <Mirror and Farmer, 6 April, 1878. S.-b. No. 20, p. 3.
 - Description of egg and manner of oviposition of Saperda bivittata [= can-dida]; seasons of oviposition; means against the same.
- 1631. [RILEY, C. V.] Buggy beans. <N. Y. Tribune, 20 February, 1878. S.-b. No. 14, p. 234.
 - Habits and distribution of and means against Bruchus fabæ; edibility of the same.
- 1632. RILEY, C. V. On the transformations of the red mites. <Amer. Nat., March [23 February], 1878, v. 12, pp. 139-146, figs. 1-6. Habits, description, and figures of Trombidium locustarum [p. 142], T. giganteum [p. 143], T. muscarum [p. 144], and Hydrachna belostomæ [p. 146], n. sp.; figures and description of most of the stages of the same; definition of Trombidium. [Advance extract from No. 1643.]
- 1633. RILEY, C. V.] Clothes moths. <Sci. Amer., 23 March, 1878 [v. 52], n. s., v. 38, p. 177. S.-b. No. 19, p. 238. Reprint: <Kansas Farmer, 17 April, 1878. S.-b. No. 19, p. 225. <Colman's Rural World, 3 April, 1878. S.-b. No. 19, pp. 198–199.

- 1633. RILEY, C. V.—Continued.
 - Indiscriminate use of names and habits of several species of Times injurious to woolen goods, fur, hair, and similar substances; means against the moths.
- 1634. RILEY, C. V. Pieris vernalis and P. protodice. <Ca. Ent., February [March], 1878, v. 10, p. 39.</p>
 Conformation of T. E. Bean's conclusion that P. vernalis is the spring form of P. protodice.
- 1635. RILEY, C. V. Migratory butterflies. <Sci. Amer., 6 April, 1878 [v. 52], n. s., v. 38, p. 215, fig. S. b. No. 19, p. 250. Extract: <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 102, fig. 34.
 - Occurrence of migratory swarms of Danais erchippus in central southern
 United States in the autumn of 1877, and of return migrations in spring;
 explanation of these migrations; mentions other migratory Rhopalocera;
 distribution of Danais archippus and Cynthia [= Pyrameis] cardui; figure of
 Danais archippus.
- 1636. RILEY, C. V. The horn-bug. <Sci. Amer., 20 April, 1878 [v. 52], n. s., v. 38, p. 249, 1 fig. S.-b. No. 19, p. 241.

 Natural history, description of egg, larva and pupa, and figures of larva, pupa, and image of Passalus cornutus.
- 1637. RILEY, C. V. Egg-feeding mites. <Ca. Ent., March [April], 1878, v. 10, pp. 58-59.

 Citation of records of Acari feeding on eggs.
- 1638. [RILEY, C. V.] Bad work of the grain Aphis <N. Y. Tribune, 26 June, 1878. S.-b. No. 19, p. 224.

 Habits and ravages of and means against Aphis avena [= Nectarophora granaria].
- 1639. [RILEY, C. V.] Of Doryphora. <N. Y. Tribune, 26 June, 1878.
 S. b. No. 19, p. 243.
 Natural history, enemies, and spread of Doryphora decombinate; description
- 1640. RILEY, C. V. The apple-tree borer. <N. Y. Tribune, 24 July, 1878. S.-b. No. 19, p. 195.

of larva and imago; methods of using Paris green.

- Means against Chrysobothris femorata.

 1641. [RILEY, C. V.] The raspberry saw-fly. <N. Y. Tribune, 24 July,
- 1878. S.-b. No. 19, pp. 197; 226.
 Habits of and means against Selandria [= Monophadnus] rubi.
- 1642. [RILEY, C. V.] Inquiring friends. <N. Y. Tribune, 24 July, 1878.
 S.-b. No. 19, p. 198.
 Means against Egeria tipuliformis.
- 1643. [RILEY, C. V., et al.] First annual report of the United States Entomological Commission for the year 1877, relating to the Bocky Mountain locust and the best methods of preventing its injuries and of guarding against its invasions, in pursuance of an appropriation made by Congress for this purpose. With map and illustrations. < Washington, 29 July, 1878, pp. 16+

477+ 295, 111 figs., 5 pls., 2 maps. See: <Sci. Amer., 5 July, 1879 [v. 55], n. s., v. 41, p. 8. S. b. No. 23, p. 121.

TABLE OF CONTENTS.

•	
LETTER OF TRANSMITTAL	X1
LETTER OF SUBMITTAL	XIII
PREFACE INTRODUCTION: By C. V. Riley.	XV
	1
Introductory remarks	•
1—Circulars sent out, 2-6—Area over which eggs were laid in	
1876, 6—Outlook in spring in more southern States, 7—Letter to	
Governor Anthony, of Kansas, on the condition of things in May,	
9-10—State of things in Minnesota in spring, 11—Outlook in Ne-	
brasks in June, 13—Second meeting of the Commission, 14—	
Outlook in Iowa in June, 15—Visit to Colorado, Utah, and Mon-	
tans in June, 17—Visit to Colorado in July, 17-20—Third meet-	
ing of the Commission, 21—Trip to the Pacific coast, 21—Trip to	
British America, 22—List of chapters, 23—Character and synop-	
sis of chapters, 24-29—Prospects for 1877, 29.	
CHAPTER I: By C. Thomas.	
Classification and nomenclature; characters of the species	13
Families of the Orthoptera, 32—Locust vs. grasshopper, 33—Ge-	
neric nomenclature, 37—Generic diagnosis, 40—Species of the	
genus Caloptenus, 42—Full definition of Caloptenus spretus and	
of its nearest congeners, 43–52.	
CHAPTER II: By C. Thomas.	
Chronological history of locust injuries	53
Compared with locust ravages in the Eastern Hemisphere, 53—In-	•
jury in the Northwest early in the present century, 54—Brief	
reviews from 1820 to 1877, 54-56—Locust history in Texas, 57-62;	
in Indian Territory, 63; in Arkansas, 64; in Missouri, 64-68; in	
Kansas, 68-74; in Nebraska, 74-77; in Iowa, 77-80; in Minne-	
sota, 80-87; in Dakota, 88-92; in Montana, 92-96; in Idaho, 96;	
in Wyoming, 97; in Colorado, 99-102; in Utah, 102-104; in New	
Mexico and Arizona, 105; in Nevada, 105; in Oregon and Wash-	
ington Territory, 106; in British North America, 108-112-Tabu-	
lar view of locust years, 113.	
CHAPTER III: By C. Thomas.	
Statistics of losses	114
Difficulty of obtaining reliable data, 114—Estimates in Kansas,	
115; in Minnesota, 116; in Missouri, 117-Loss in Kansas, Ne-	
braska, Iowa, and Missouri in 1874, 118-Loss to different crops,	
120-Loss to Missouri in 1875, 121-Total loss during the years	
1874-77, \$200,000,000, 122.	
CHAPTER IV: By C. Thomas.	
Agricultural bearing of the locust problem	123
Drawback to the settling of the West, 124-What is likely to be	
the effect in the future, 125-Modification and settlement of the	
Western plains, 129-Crops which suffer most, and those which	
suffer least, 130—Small grains not affected by invading swarms,	
128—Need of judgment in planting, 129.	

CHAPTER V: By A. S. Pachard, jr.

Permanent breeding-grounds of the Rocky Mountain locust...... Permanent breeding-grounds, 131-Definition of the permanent region, 133—Its character and extent, 134—The Rocky Mountain locust a sub-boreal insect, 135-The sub-permanent region, 136-The temporary region, 136.

CHAPTER VI: By C. Thomas and A. S. Pachard, jr.

Geographical distribution Eastern limits of spread, 137—Northern limits, 139—Western limits, 140-Southern limits, 141-Sub-permanent region, 142.

CHAPTER VII: By A. S. Pachard, jr., and C. Thomas.

Migrations..... 143

Classification of flights, 143-Invading swarms, 143-Returning swarms, 143—Local flights, 143—Height at which swarms move, 144-Effect of change of wind and weather on flights, 145-Flight at night, 147-Migrations previous to 1877 east of the Rocky Mountain Plateau, 148-Direction of invading swarms prior to 1877, 149-Direction of flight in 1876, 151-Migrations within the permanent region, 153-158; in Montana, 153; in Wyoming, 156; in Colorado, 157; in Eastern Idaho and Utah, 158; in the lower Snake Valley, 158—Return migrations from the temporary region previous to 1877, 159-162—Return migrations in 1877, 162-165-Local flights in 1877, 165-Record of flights for July 3 and July 20, 1877, 165-169—Summary of flights by States, 170-174-Southward flights in 1877, 175-Movements in different directions at one time, 176-Destination of return swarms, 177.

CHAPTER VIII: By C. V. Riley.

Habits and natural history..... 212

Destructive powers of locusts, 212-215—Stoppage of railroad trains, 215-Rate at which locust swarms move, 215-Velocity of flight, 216-Direction of invading swarms, 217-Time of appearance of invading swarms, 217-Flight at night, 218-Height of flight, 219—General habits at night, 219—Where the eggs are laid, 222— Manner in which the eggs are laid, 223—Philosophy of the eggmass, 225—The female lays more than one egg-mass, 226—Interval between different egg-layings, 227-Number of eggs laid, 228-The hatching process, 228-231-Where and under what conditions of soil the young hatch most freely, 231-Time of hatching, 231-Habits of the young or unfledged locusts in the temporary region, 232-Directions in which the young travel, 234-Rate at which the young travel, 235—They reach but a few miles east of where they hatch, 235-Not led by kings and queens, 236-Time of year when wings are acquired, 237—Direction taken by swarms departing from the temporary region, 238-Destination of departing swarms, 238—Do the return swarms breed? 239—Do return swarms from the temporary region retrace their course? 240-The species essentially single-brooded, 240-243-Reasons why it cannot produce two generations annually, 243-The species can not permanently dwell in the temporary region, 244-The insects which hatch there do not remain, 246—Extensive and thick egg-laying seldom occurs twice consecutively in the same locality, 247-Reasons why it does not, 248-Causes of migration. 249 - Food-plants, 251-Crops and plants most liked and those feast liked, 252-254-Unnecessary alarm caused by comparatively harmless species, 255.

131

164 3.	RILEY, C. V.—Continued.	
	CHAPTER IX: By A. S. Pachard, jr.	
	Anatomy and embryology	257
	External anatomy, 257—Divisions of the body, 258—Sexual differences, 259—Internal anatomy, 261—The digestive system, 262—264.—The nervous system, 264—The heart, 266—The trachese and	
	dilated air-sacs, 267—Mode of breathing, 269—The locust an aëronaut, 270—The male reproductive system, 270—The female	
	reproductive system, 271—Organs of special sense, 272—The fine anatomy of the locust, 273—Embryology of the Rocky Mountain	
	locust, 277—Different egg-layers, 278—How the embryo lies within the egg, 278—How it bursts the egg, 279.	•
	CHAPTER X: By C. V. Riley.	
	Metamorphoses	279
	The six stages of growth, 279—Variability in the depth of coloring, 281—The process of molting, 281—Time required for it, 283—Differences in the immature stages between the Rocky Mountain, the lesser, and the red-legged locusts, 263.	
	CHAPTER XI: By C. V. Riley.	
	Invertebrate enemies	284
	Value of the locust's minute enemies, 284—Animals that destroy the eggs, 285—The Anthomyia egg-parasite, 285—The common	
	flesh-fly, 289—Ground-beetles and their larvæ, 289—Harpalus larvæ, 289—The egg-feeding Amara, 291—Blister-beetle larvæ, 292—Their character and locust egg-feeding habits, 293—History of the oil-beetle, 294—History of Sitaris, 295—History of Hornia,	
	296—History of Epicauta, 237—Macrobasis and Henous, 301— Other meloid genera, 302—Soldier-beetle larvæ, 302—Asilid	
	larvæ, 303—Click-beetle larvæ, 304—Miscellaneous species, 305—Chalcid-fly, 306—Animals that prey on the locust after it is born, 306—The locust-mite, 306—The efficacy of its work, 308—Its transformations, 309—Other mites, 312, 313—Ground-beetles,	
	313—Tiger-beetles, 314—Asilus-flies, 317—Digger-wasps, 317—Tachina-flies, 319—Their efficacy in destroying locusts, 321—Florb flies, 322 Jahannan flies, 324 Hair manne, 325 Their	
	Flesh-flies, 323—Ichneumon-flies, 324—Hair-worms, 326—Their curious life-history, 327—332—Insects attacked by hair-worms, 327—How hair-worms get into locusts, 332—Miscellaneous locust enemies, 334.	
	CHAPTER XII: By C. Thomas.	
	Vertebrate enemies	334
	Good offices of birds probably underrated, 334—Experience of correspondents, 336—Some of the most useful birds, 338—Paper by Professor Aughey on the beneficial work of birds, 338—Enormous number of birds destroyed for market, 346—Damage done to insectivorous birds by birds of prey, 348—The English sparrow,	
	349—What public sentiment needs, 349.	
	CHAPTER XIII: By C. V. Riley.	250
	Remedies and devices for destruction Encouragement of natural agencies, 351—Destruction of the eggs,	350
	351—Harrowing in the autumn, 353—Experiments to test the effect of alternately freezing and thawing, 353—Experiments to test the effects of exposure to air, 354—Experience with harrowing, 355—Plowing, 356—Experiments to test the effects of bury-	
	ing at different depths, 356—Experience in plowing, 358—Irrigation, 359—Experiments to test the effects of moisture on the	
	13 ENT——15	

CHAPTER XIII-Continued.

eggs, 359-Tramping, 361-Collecting, 361-Destruction of the young or unfledged locusts, 362-Burning, 363-The burning of prairies, 363-The Hetzel burning machine, 363-The Horner burning contrivance, 363—Hand burners, 364—The Atwood machine, 364-Use of wire and kerosene, 364-Crushing, 364-The Drum locust-crusher, 365—The Simpson locust-crusher, 366— The Hoos locust-crusher, 367-The Hansberry locust-crusher, 368—The Kenworthy locust machine, 370—J. C. Melcher's machine, 371—The Peteler machine, 371—The King suction-machine, 374—The Flory locust-machine, 376—Trapping locusts, 377—Nets and seines, 377-Ditching and trenching, 378-Protection by barriers, 381-Coal-oil, 381-Coal-oil pans, 383-The Canfield pan, 384—The Adams pan, 385—The Anderson coal-oil contrivance, 386-Mr. Long's contrivance, 386-Mr. Watrous's contrivance, 387-Mr. Swearingen's contrivance, 387-Use of coaltar, 387—The Robbins pan, 388, 390—Use of coal-tar previous to 1877, 386-Other machines for the use of coal-tar, 391-Catching and bagging, 391-Principles and facts to be borne in mind in catching locusts, 391-The Riley machine, 392-Mr. Thompson's net, 392-The Elliot catcher, 394-The Wilson-Rhode catcher, 395-Contrivance for catching the pupe, 394-The Godard catcher, 395-The Benson catcher, 396-The Hutchins catcher, 396—The Sylvester catcher, 397—The Hero'hopper-catcher, 398— The Belt device, 398-Hand-nets, 399-Use of destructive agents, 399—Buhach or Persian insect-powder, 400—Veith's insect-fluid, 400-Sulpho-carbonate of potassium, 401-Naphthaline, 401-Paris green, 401—The protection of fruit trees, 403—Destruction of the winged insects, 404-Fumigation, 405-The effect of concussion, 405-Diversified agriculture, 406-Legislation, 407-Bounty laws, 409-Missouri locust act, 409-Kansas locust acts, 410-Minnesota locust acts, 412-Nebraska locust act, 413-Suggestions that may be of service, 414-More attention needed to the growth of root-crops, 415—The benefits of irrigation, 415— Hogs and poultry, 415-Non-planting, 415-Use of soldiers, 416-Preventive measures against the winged insects, 417—Further investigation needed, 417-Means that have been suggested against the winged insects, 418-Systematic burning of young, 418—Co-operation with Dominion Government, 419—Protection by smoke, 419—Farmers should receive the locust probabilities, 420.

CHAPTER XIV: By C. V. Riley.

421

423

CHAPTER XV: By C. V. Riley.

Influence of weather on the species.....

Meteorological conditions affecting the life of the locust, 423—
Effects of weather on the young locusts, 424—Effects of weather on the eggs, 424—Possibility of predicting consequences upon meteorological grounds, 424—The temperature of the soil, 425—
Thermal constant for the hatching of locust eggs, 426—Number of hours which eggs must be exposed to hatch at a temperature of 60° F., 431—Number of hours required in 1875.

1643. RILEY, C. V.—Continued.	
CHAPTER XVI: By C. V. Riley.	
Effects that generally follow severe locust injury	432
manently injured, wild grasses not, 435—Permanent effects on different plants, 436—Injury to fruit trees, 436. CHAPTER XVII: By A. S. Pachard, jr.	
Uses to which locusts may be put	437
CHAPTER XVIII: By C. V. Riley.	
Ravages of other locusts in the United States	443
CHAPTER XIX: By A. S. Pachard, jr.	
Ravages of locusts in other countries	460
Locust injuries in Central America, 460—Great destruction in Honduras and Guatemala, 462—The locusts in South America, 465—The locusts in the Old World, 467—Injuries in Germany, 468—Habita of the European species, 469—Injuries in Russia, 470—Different species affecting the Old World, 471—Geographical distribution of the European migratory locust, 472—476—Notes on Algerian locusts, 476—Injury in China, 477—In southern Australia, 477.	
APPENDICES.	
APPENDIX I: Mr. Allen Whitman's report from Minnesota	[3]
State of things in spring, [4]—Exertions of farmers, [4]—Different rieans employed to destroy the eggs, [4]—Effects of temperature and rain-fall on the eggs, [5]—Dates of hatching, [6]—Late hatching, [7]—Progress during spring, [8]—First winged, [10]—Statistics of damage, [11]. Appendix II:	
Aughey on locust-feeding birds	[13]
Letter of transmittal, [13]—Examinations of the contents of the stomachs of birds of Nebraska, giving the number of locusts, number of other insects, and number of seeds of the different birds considered in their proper classificatory position, [14]-[62].	[10]

1643. RILEY, C. V.—Continued.	
APPENDIX III: Texas data for 1877	
APPENDIX IV: Missouri data for 1877	
APPENDIX V: Kansas data for 1877 Report by George F. Gaumer, special assistant, [85]—Report by N. Godfrey, special assistant, [88]—Reports from corresponder and miscellaneous matter, chronologically arranged, [90]-[10	A. its
APPENDIX VI: Iowa data for 1877 Detailed data from correspondents, and from other sources, chronically arranged, [104]—[110].	
APPENDIX VII: Colorado data for 1877 Report from William Holly, special assistant, [111]—Data from other sources, [113]—[116].	
APPENDIX VIII: Nebraska data for 1877	ly
APPENDIX IX: Narrative of the first journey made in the summer of 1877, by I Packard	[134]
APPENDIX X: Narrative of a second journey in the summer of 1877, by Mr. Packas From Chicago to Utah, [139]—Observations in Oregon and Waington Territory, [140]—From Vancouver Island to Californ [141]—Variations in Caloptonus spretus, atlants, and femur-rubra [143].	h- ia,
APPENDIX XI: British-American data Characteristics and prevailing winds of Manitoba, [145]—Do from the Cypress Hills region, [146]—The locust breeds permontly in the third prairie steppe, [146]—No damage ever do in the Peace River country, [146]—Ravages in the Northwest the early part of the century, [147]—Records by the Hon. Do ald Gunn, [148].	ita na- no in
APPENDIX XII: Autumn flights in 1877	

	•	
1643.	RILEY, C. V.—Continued.	
	APPENDIX XIII: Flights and migrations in 1877 Locust movements in the more southern portion of the temporary region, [156]-[164]Flights in the more northern parts of the locust country, [164]-[197].	[156]
	APPENDIX XIV:	
	Answers to the question: Did any locusts remain in the temporary region after the departure of the return swarms?	
	APPENDIX XV:	
	Prevailing direction in which the young insects travel	
	APPENDIX XVI:	
	Time of year when the bulk of the insects become winged Data from various States and Territories, [205]-[206].	[205]
	APPENDIX XVII:	
	Habits of the insects at night Nocturnal habits of locusts in Iowa, Dakota, and Nebraska, [207]— In Minnesota and elsewhere, [208]–[210].	[907]
	APPENDIX XVIII:	
	Facts relating to the eggs	
	APPENDIX XIX:	
	Means of destruction	[218]
	APPENDIX XX:	
	Injury: crops affected	[222]
	APPENDIX XXI: Previous visitatious—Aid of animals Data from Nebraska and Iowa, [226]—From Minnesota, [227]— From Dakota, [229].	[226]
	APPENDIX XXII:	
	Minnesota data	[230]
	APPENDIX XXIII:	
	Are the eggs ever laid thickly for two consecutive years in the same ground ?	[236]
	Answers to the above question from Arkansas, Colorado, and Iowa, [236]—From Kansas, [237]—From Missouri, [238]—From Minnesota, Nebraska, and Texas, [239].	- •
	APPENDIX XXIV:	
	Miscellaneous data	[240]

1643. RILEY, C. V.—Continued.

APPENDIX XXV:

From Dakota, [243]-[247]-From Montana, 248-From Utah,

Data from Dakota, Montana, Utah, and New Mexico...

[253]—From New Mexico, [259].

[243]

	APPENDIX XXVI:
	List of correspondents
	In Arkansas, British America, and California, [261]—In Colorado and Dakota, [262]—In Idaho and Iowa, [263]—In Kansas, [264]—
	In Minnesota, [265]—In Missouri, [267]—In Montana, [269]— In Nebraska, Nevada, and New Mexico, [270]—In Texas, Utah,
	and Washington Territory, [271]—In Wyoming, [272].
	APPENDIX XXVII:
	Bibliography on the locusts of America. By B. P. Mann
1644.	RILEY, C. V. Attractive but untrue. <n. 1878.="" 19,="" 198.<="" 31="" july,="" no.="" p.="" sb.="" td="" tribune,="" y.=""></n.>
	Absurdity of statement that "a parasite of the strawberry plant has been found to wage a war of extermination against the Phylloxera."
1645.	RILEY, C. V. Locusts eat the castor bean. <n. 14<br="" tribune,="" y.="">August, 1878. Sb. No. 19, pp. 198, 227-228.</n.>
	Acridida relish Fagopyrum and Linum but eat Ricinus communis with reluctance, though with impunity.
16 4 6.	[RILEY, C. V.] The stalk-borer. <n. 1878,="" 19,="" 21="" 227;="" 242-243;="" 244.<="" 37.="" august,="" no.="" pp.="" sb.="" td="" tribune,="" v.="" weekly="" y.=""></n.>
	Description, food-plants, and habits of larva and description of image of Gortyna nitela; G. nebris a variety of G. nitela.
1647.	RILEY, C. V. That hundred and fifty million dollars. <sci. 117.="" 1878="" 19,="" 24="" 245.<="" 39,="" 53],="" [v.="" amer.,="" august,="" n.="" no.="" p.="" s.,="" sb.="" td="" v.=""></sci.>
	Critical review of sensational reports in current agricultural papers; an appropriation of \$5,000 made to the U.S. Department of Agriculture for cutton insect investigation.
164 8.	RILEY, C. V. Silk-worm breeding. <sci. 1878<="" 24="" amer.,="" august,="" td=""></sci.>
•	[v. 53], n. s., v. 39, p. 119. S. b. No. 19, p. 244. Number of annual generations of <i>Sericaria mori</i> ; inferiority of races breeding more than once annually.
1649.	RILEY, C. V. Cotton-worm. < Daily Constitution [Atlanta, Ga.], 8 September, 1878, v. 11, No. 73, p. 1.
	Alstia argillacea [=xylina] feeds from the extra-floral nectar-glands of the cotton-plant.
1650.	RILEY, C. V. That "fatherless and motherless race." The bas-
	ket-worm, alias drop-worm, alias bag-worm, Thyridopteryz ephemeræformis. <sci. 1878,<="" 28="" amer.="" september,="" suppl.,="" td=""></sci.>
	fig. Sb. No. 19, pp. 245-246.

worm," with additions; degradation and breeding habits of female Psychidæ; method of imagination, fecundation, and oviposition of Thyridop-

- 1650. RILEY, C. V.—Continued.
 - teryx ephemeræformis; description of eggs and figures of all other stages; down in which the eggs are enveloped is extruded from the abdomen with the eggs; food-plants, parasites, and means against the species; its follicles available for silk production; this species proterogynous but most insects proterandrous.
- 1651. RILEY, C. V. Notes on the life-history of the blister-beetles, and on the structure and development of *Hornia*. <Kansas City Review of Science and Industry, September, 1878, vol. 2, No. 6, p. 353. S.-b. No. 19, p. 209. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1878 [14 July], 1879, v. 27, pp. 284-285. See: <Ca. Ent. September | October|, 1878, v. 10, pp. 177-178.
 - Occurrence in *Epicauta* and *Macrobasis* of the hypermetamorphoses characteristic of *Meloida*; specification of the stages of the same; summary of the life-history of *Hornia minutipennis* parasitic on *Anthophora abrupta*.
- 1652. RILEY, C. V. On the larval characteristics of Corydalus and Chauliodes, and on the development of Corydalus cornutus.

 <Kansas City Review of Science and Industry, September, 1878, v. 2, No. 6, p. 354. S.-b. No. 19, pp. 210-211. Reprint:

 <Ca. Ent., May, 1879, v. 11, pp. 96-98. <Proc. Amer. Assoc. Adv. Sci. for 1878, [14 July], 1879, v. 27, pp. 285-287. See:

 <Ca. Ent., September [October], 1878, v. 10, p. 178.
 - Characters and economic value of larva of Corydalus cornutus; characters in brief of the imago; characters of larva of Chauliodes; description of eggs of both genera; situation of egg-masses of Corydalus and physical character of their covering; description of respiratory apparatus and method of respiration of larvæ of Corydalus.
- 1653. RILEY, C. V. Biological notes on the gall-making *Pemphigina*.
 Kansas City Review of Science and Industry, September, 1878, v. 2, No. 6, p. 380. S.-b. No. 19, pp. 211-212: 212. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1878, [14 July], 1879, v. 27, pp. 288-289. Abstract: <Nature, 28 November, 1878, v. 19, p. 75. Extract: <Sci. Amer., 26 October, 1878 [v. 53], n. s., v. 39, p. 266. S.-b. No. 19, p. 244.
 - Method of hibernation of gall-making Pemphigina on Ulmus.
 - NOTE. -Nos. 1651-1653 and 1668-1669 were reprinted from Proc. Amer. Assoc. Adv. Sci., v. 27, under a separate cover. Salem, February, 1879.
- 1654. [RILEY, C. V.] The cottonwood killer. <N. Y. Weekly Tribune,
 9 October, 1878. S.-b. No. 19, p. 242. Extract: <Amer. Ent.,
 July, 1880 [v. 3], n. s., v. 1, pp. 159-160.
 - Habits, ravages, and food-plants of *Lina scripta*; description of and means against the same; *Trypeta pomonella* an example of the acquisition of new habits in insects.
- 1655. [RILEY, C. V.] A new insect foe to green corn. <N. Y. Tribune, 9 October, 1878. S.-b. No. 19, pp. 234; 244. Ravages of Cetonia [= Euphoria] inda.
- 1656. [RILEY, C. V.] Inquiring friends. <N. Y. Tribune, 9 October, 1878. S.-b. No. 19, p. 245.
 - Ravages and food-plants of Hylosiaus opaculus; impracticability of means against bark-borers in large trees.

1657. RILEY, C. V. Notes from the South. Facts about the cottonworm. <Sci. Amer., 16 November, 1878 [v. 53], n. s., v. 39, pp. 312–313.
S.-b. No. 19, pp. 213–214; 239–241.
See: <Amer. Rural Home, 27 September, 1878.
S.-b. No. 23, p. 128.

Note of a trip in the southern United States in 1878; commission to investigate insects injurious to the cotton-plant; food-plants and habits of the larva and image of Aletia argillacea [=xylina]; use of poisonous baits and of Paris green; Anomis exacta injurious in certain regions.

1658. RILEY, C. V. Some further facts regarding that "fatherless race." <Sci. Amer. Suppl., 30 November, 1878. S.-b. No. 19, pp. 238-239.

Description and figures of male copulatory organs of Thyridoptoryx ophemoraformis; description of method of copulation; figures larva, pupa, imagos, and follicles of the same; "parthenogenesis, though not improbable, seldom takes place in this species;" nature of the down intermingled with the eggs; previous error concerning this down.

1659. [RILEY, C. V.] "The carpet bug." <N. Y. Tribune, 1 December, 1878. S.-b. No. 19, p. 236.

Ravages of and means against Anthronus scrophulariæ; figures larva, pupa and imago.

1660. [RILEY, C. V.] A bug that eats bees. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, p. 237. Reprint: <Prairie Farmer, 4 January, 1879, v. 50, p. 3. S.-b. No. 23, p. 110. Description and habits of Phymata erosa.</p>

1661. [RILEY, C. V.] Tomato worm. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, p. 237.

Excessive abundance of larva of Sphinx quinquemaculata [= Protoparce celcus] near Port Hope, Ontario, in 1878.

1662. [RILEY, C. V.] Worm snake. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, p. 237.

Habits and abode of larvæ of Sciara sp. at Orange, Conn.

1663. [RILEY, C. V.] Carpet pests. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, p. 237.

Food habits, spread of, and means against Anthrenus scrophularia.

1664. RILEY, C. V. Corn worm. <N. Y. Tribune, 4 December, 1878.
 S.-b. No. 19, pp. 235; 237.

Food-plants and means against Heliothis armigera.

1665. [RILEY, C. V.] Hessian-fly. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, pp. 235; 237.

Indications of the ravages of and food-plants of Cecidomyia destructor.

1666. [RILEY, C. V.] Apple-worm. <N. Y. Tribune, 4 December, 1878. S.-b. No. 19, pp. 235; 237.

Baldwin apples more exempt than other varieties from the attacks of the second brood of Carpocapsa pomonells.

1667. RILEY, C. V. Anent the English sparrow. < Evening Star [Washington, D. C.], 28 December, 1878, v. 52, No. 8030, p. 1. S.-b. No. 19, pp. 214-215; No. 23, p. 151.

Value of Passer domesticus as a destroyer of insects and in other respects; objections to the special protection of the same.

- 1668. RILEY, C. V. Silk-culture; a new source of wealth to the United States. <Kansas City Review of Science and Industry, 1878, v. 2, pp. 419-423. S.-b. No. 19, pp. 216-218. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1878, [14 July], 1879, v. 27, pp. 277-283. Extract: <Kansas Farmer, 1879: 26 February, 12 November. S.-b. No. 23, pp. 105-106; No. 39, pp. 91-92. See: <Ca. Ent., September [October], 1878, v. 10, p. 178. <St. Louis Globe-Democrat, 28 August, 1878. S.-b. No. 19, p. 203. Practicability and desirability of the extensive establishment of silk-culture
 - Practicability and desirability of the extensive establishment of silk-culture in the United States; sketch of efforts made towards such establishment; requisites to its success.
- 1669. RILEY, C. V. The philosophy of the movements of the Rocky Mountain locust. <Kansas City Review of Science and Industry, 1878, v. 2, pp. 424-427. S.-b. No. 19, pp. 218-220. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1878, 14 July, 1879, v. 27, pp. 271-277. See: <St. Louis Daily Globe-Democrat, 24 August, 1878, v. 4, No. 95.
 - Limits of breeding-grounds of and region invaded by Caloptenus spretus; causes, periods, and directions of flights of the same; causes of the limitation of the flights; ravages; generalizations regarding locust invasions; work of the U. S. Entomological Commission.
- 1670. RILEY, C. V. A complete life-history of the army-worm, Leucania unipuncta, and its parasites. <25th Ann. Rept. Secr. Mass. Board Agric. for 1877, 1878, pp. 243-253, figs.
 - Early history, synonymy, distribution, seasons, enemies, and parasites of Leucania unipuncia; habits of larvæ and of female imagos; descriptions of all stages; figures of all stages except the egg; figures of two unnamed parasitic Ichneumons. Parasitic habits of Exorista [= Nemoræa] leucaniæ, E. flavicauda, Microgaster [= Apanteles] militaris, Pesomachus minimus, and Ophion purgatus; descriptions of Microgaster [= A.] militaris, Mesochorus vitreus, and Ophion purgatum. Ichneumon leucaniæ also a parasite.
- 1671. RILEY, C. V. Phylloxera. < Johnson's New Universal Cyclopædia, 1878, v. 3, pp. 1241-1243, 8 figs. S.-b. No. 10, pp. 192-194. Definition of the genus Phylloxera; list of the sixteen (16) described United States species; descriptions of their galls; distribution, history, description, and figures of various stages and ravages of and means against Phylloxera vastatrix.</p>
- 1672. RILEY, C. V. Potato-bug. < Johnson's New Universal Cyclopædia, 1878, v. 3, pp. 1361-1364, 9 figs. S.-b. No. 10, pp. 195-198.

 List of principal enemies of potato-plant; spread, natural history, food-plants, enemies of, and means against Doryphora decemlineata; figures of the same in its different stages, and of Lydella [= Exorista] doryphora, Calosoma calidum, Hippodamia convergens, Mysia [= Anatis] 15-punctata, Arma [= Podisus] spinosus, Harpactor [= Milyas] cinctus, Perillus circumoinctus, and Doryphora juncta.
- 1673. RILEY, C. V. Weevil. < Johnson's New Universal Cyclopædia, 1878, v. 4, p. 1338, fig. S.-b. No. 14, p. 46.</p>
 Definition of "weevil;" mention of numerous species, with statement of their food-plants and manner of obtaining food; habits and means against

Sitophilus [= Calandra] granaria and S. [= C.] oryzæ.

- 1674. RILEY, C. V. The locust swarms that devastate the trans-Mississippi country; their source, movements, and eastern limit.

 <Western Farmer's Almanac for 1879, 1878, pp. 48-50. S.-b.
 No. 19, pp. 247-248.
 - Limits of the breeding-grounds of and of regions invaded by Caloptenus spretus; causes, periods, and directions of their flights; causes of the limitations of the same physical and principally atmospheric; amount of ravages.
- 1675. [RILEY, C. V.] Prof. Riley in favor of the birds. <Evening Star [Washington, D. C.], 9 January, 1879, v. 53, No. 8039, p. 3. S. b. No. 19, pp. 223-224; No. 23, p. 94.
 - From St. Louis [Mo.] Globe-Democrat. List of some birds which are beneficial to horticulture and some which are noxious.
- 1676. RILEY, C. V. Letter from Prof. C. V. Riley. <Colman's Rural World, 15 January, 1879. S.-b. No. 23, pp. 107-108.
 - Letter to N. J. Colman transmitting report as former treasurer of the Missouri State Horticultural Society; transmission of documents for distribution; proposal for a revised and condensed edition of author's reports as State entomologist of Missouri.
- 1677. [RILEY, C. V.] Michigan apples and codling-moth. <N. Y. Tribune, 15 January, 1879. S. b. No. 23, p. 109.
 - Freedom of apples from Carpocapsa pomonella in 1878 in Michigan due to the smallness of the apple crop in 1877 and to the efforts made by the orchardists to exterminate the insects.
- 1678. RILEY, C. V. Notes on the Aphididæ of the United States, with descriptions of species occurring west of the Mississippi. <Bull. U. S. Geol. and Geog. Surv. Terr., 28 February, 1879, v. 5, pp. 1-32, pl. 1-2, with 2 pp. expl. of pl. Separate: <Washington, 22 January, 1879, 32 pp., 2 pl., with 2 pp. expl. of pl. Abstract: <Sci. News, 15 April, 1879, v. 1, pp. 184-186.
 - Part 1, pp. 1-17, by C. V. Riley, entitled "Biological notes on the Pomphigina, with descriptions of new species;" history and description of Schizoneura americana n. sp. [p. 4], Colopha ulmicola, Pemphigus populi-monilis n. sp. [p. 13], P. populi-transversus n. sp. [p. 15], P. p.-ramulorum n. sp. [p. 16], P. accrifolii n. sp. [p. 16], P. fraxinifolii n. sp. [p. 17], and Hormaphis spinosus.
 - Part 2, pp. 18-32, by J. Monell, entitled "Notes on Aphidina, with descriptions of new species." Describes nine (9) new species of Siphonophora, five (5) of Aphis, two (2) of Rhopalosiphum, nine (9) of Callipterus, three (3) of Chaitophorus; description of other species of these genera and of Drepanosiphum and the genera above named; synoptic tables of some of the species.
- 1679. [RILEY, C. V.] [Plant-lice on potatoes.] <N. Y. Tribune, 12 February, 1879. S.-b. No. 23, p. 109.

 Aphididæ reputed injurious to potato plants, probably Aphis solani.
- 1680. RILEY, C. V. Missouri entomological reports. < Colman's Rural World, 19 February, 1879. S.-b. No. 23, p. 102.
 - Proposal for republication of the reports of the State entomologist of Missouri; reasons for the non-publication of a tenth report; sketch of what that report would have contained.

- 1681. RILEY, C. V. Entomological notes. The chinch-bug. <Farmer's Review. February, 1879, 2 figs. S.-b. No. 23, pp. 111-112.
 - Figures of all stages, ravages, natural history, and means against Blissus loucopterus; its method of hibernation, and the effect of weather upon it.
- 1682. RILEY, C. V. Notes on the apple-worm. <Colman's Rural World, 5 March, 1879. S. b. No. 23, p. 102. Reprint with slight changes: <Amer. Nat., August [July], 1879, v. 13, pp. 523-524. <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, pp. 238-239.

See No. 1721 for synopsis of contents.

- 1683. RILEY, C. V. The rice weevil. Serious injury to stored and to cribbed corn. <Farmer's Review, March, 1879, fig. S.-b. No. 23, p. 99; 112.
 - Figures larva, pupa, and imago of Calandra oryzæ and imago of C. granaria; habits, ravages, and means against the same; description of egg and method of oviposition of C. oryzæ; use of carbon bisulphide against these insects; poisonous effects of eating the comminuted beetles.
- 1684. RILEY, C. V. [Letter to W. G. Le Duc.] < Boston Daily Herald, 4 April, 1879. < Psyche Advertiser, 11 April, 1879, v. 2, p. 9. < Ent. Nach., 1 July, 1879, jahrg. 5, p. 177.
 - The reason for which the author resigned as entomologist of the U. S. Department of Agriculture was not ill-health.
- 1685. RILEY, C. V. Preventing rot in plums. <N. Y. Tribune, 9 April, 1879. S.-b. No. 23, p. 110; 129.
 - Trapping and jarring recommended as a means against Conotrachelus nenuphar.
- 1686. [RILEY, C. V.] A new insect pest. <Colorado Farmer, 10 April, 1879, v. 12, No. 15, p. 6. S.-b. No. 23, p. 108.
 - Injury to orchards and ornamental trees in California by *Dorthesia* [characias *], introduced from Australia; ravages of *D. characias* in southern Africa; prospective ravages in California; means against it.
- 1687. RILEY, C. V. The nervous system and salivary glands of Phylloxera. <Psyche, 11 April, 1879, v. 2, pp. 225-226.
 - Correctness of E. L. Mark's conclusion that the author had mistaken nerve cords for tracheæ; dissent from M. Cornu's view that the root swellings caused by *Phylloxera* are due to mechanical action of the puncture and the subsequent absorption of liquids; swelling considered to be caused by the introduction of a salivary secretion.
- 1688. RILEY, C. V. Mr. Henderson's experiments. < Gardener's Mo. and Hortic., April, 1879, v. 21, pp. 120-121. S.-b. No. 23, p. 107.
 - Review of P. Henderson's "Carnivorous plants;" believes that Dionæa and Drosera are nourished by the insects digested by their leaves.
- 1689. RILEY, C. V. The migrations and hibernation of Aletia argillacea. <Washington [D. C.] World, 10 May, 1879. <Science News, 1 June, 1879, v. 1, pp. 230-232. S.-b. No. 23, pp. 119-120. <Sci. Amer., 14 June, 1879 [v. 54], n. s., v. 40, p. 375. <Galveston [Tex.] Daily News, 24 October, 1879, v. 38, p. 2.

- 1689. RILEY, C. V.—Continued.
 - S.-b. No. 23, pp. 166–167, No. 42, pp. 38–39. Abstract: < Amer. Nat., November [25 October], 1879, v. 13, p. 726. < Farmer's Review, September, 1879. S.-b. No. 23, p. 108.
 - Ravages of insects, particularly of Aletia argillacen [= xylina] in the United States; hibernation theories; author's belief in the probable hibernation of the moth in the southern parts of the cotton belt; species mistaken for Aletia; description and food-plants of Aspila virescens.
- 1690. RILEY, C. V. Insects affecting clover. <N. Y. Tribune, 14 May, 1879. S.-b. No. 23, p. 90; 97; 129.
 - Habits, ravages of and means against Hylesinus trifolii and Cecidomyia leguminicola; prior observations upon the same; characters distinguishing C. leguminicola from C. [= Diplosis] tritici and C. destructor.
- 1691. RILEY, C. V. [*Œcanthus niveus.*] <N. Y. Tribune, 14 May, 1879.
 S.-b. No. 23, p. 95; 98.
 - Occurrence of eggs of *Ecanthus* niceus in twigs of apple, raspberry, grape, and other plants; slight injuries to be expected from the same; means against them.
- .1692. [RILEY, C. V.] Insect powders and their use. <N. Y. Tribune, 14 May, 1879. S.-b. No. 23, p. 95; 96; 130-131.
 - Abstract of W. Saunders' "Insect powder," with additions; differences between Persian and Dalmatian insect powder; methods of using them against insects.
- 1693. [RILEY, C. V.] The grasshopper prospect. <N. Y. Tribune, 14 May, 1879. S.-b. No. 23, p. 98.
 - Prospect of injuries by Caloptenus spretus in 1879; a new species injurious in the Sierra Valley of California.
- 1694. RILEY, C. V. The westward progress of the imported cabbageworm. <Amer. Nat., June [20 May], 1879, v. 13, p. 393.
 See: <Colorado Farmer, 10 April, 1879. S.-b. No. 23, p. 108.
 <Farmer's Review, April, 1879. S.-b. No. 22, p. 64. <Colman's Rural World, 1879. S.-b. No. 23, p. 108. <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, p. 239.
 Spread of Pieris rapæ into Illinois and Missouri; means against the same.
- 1695. [RILEY, C. V.] Lures for moths. <N. Y. Tribune, 28 May, 1879. S.-b. No. 23, p. 95.
 - Trapping Carpocapsa pomonella in shallow dishes of sweetened liquid more harmful than beneficial, as many useful insects are destroyed; the method serviceable against Heliothis armigera.
- 1696. [RILEY, C. V.] The current-worm. <N. Y. Tribune, 11 June, 1879. S.-b. No. 23, p. 106.
 - Description, habits, distribution, and spread of and means against Nematus ventricosus [=ribesii]; methods of using hellebore.
- 1697. [RILEY, C. V.] Sweet-potato beetle. <N. Y. Tribune, 11 June, 1879. S.-b. No. 23, pp. 106—107; 131.
 - Description of all stages, habits, food plants, and means against Coptocycla [= Cassida] nigripes.

- 1698. RILEY, O. V. [The seventeen year Oicada.] <N. Y. Tribune, 1879. Reprint: <Colman's Rural World, 25 June, 1879. S.-b. No. 23, p. 152.
 - Boundaries of the areas in which Cicada [= Tibican] septendecim is expected to occur in 1879; request for information of its appearance.
- 1699. RILEY, C. V. The imported carpet beetle, Anthrenus sorophularia, L. <Farmer's Review, 1879, fig. S.-b. No. 23, pp. 96–98. Extract: <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 54.
 Letter of J. H. Parsons, with answer; description of larva and image of Anthrenus scrophularia; its history, habits, ravages, and means against them; figures larva, pupa, and image.
- 1700. [RILEY, C. V.] The cheese-skipper. <N. Y. Tribune, 2 July, 1879. S.-b. No. 23, p. 93. Notice: <Farmer's Review, 7 September, 1879. S.-b. No. 23, p. 103.
 - Description of larva, pupa, and imago, habits and means against Ptophila caset.
- 1701. BILEY, C. V. The thick-thighed walking-stick. <Sci. Amer., 5
 July, 1879 [v. 55], n. s., v. 41, pp. 7-8, fig. S.-b. No. 23, p. 121.
 Reprint, with slight changes: <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, pp. 241-245, pl. 3.
 Treats of Diapheromera femorata. See No. 1721 for synopsis of contents.
- 1702. RILEY, C. V. The cotton-worm. < Mobile Register, 9 July, 1879. S.-b. No. 68, p. 204. See: < Colorado Citizen, 17 July, 1879. S.-b. No. 23, p. 91. < Galveston News, 19 July, 1879. S.-b. No. 23, p. 93.
 - Remarks at the Cotton Exchange about prospective injuries from Aletia argillacea [=xylina]; sudden appearance and means against the same; more injurious in wet than dry weather.
- 1703. RILEY, C. V. Other insects affecting cheese. <N. Y. Tribune, 9 July, 1879. S.-b. No. 23, p. 92. Reprint: <Amer. Dairyman. <Western Rural, 9 August, 1879, v. 17, No. 32, p. 250. S.-b. No. 23, p. 92.
 - Habits and transformations of Tyroglyphus siro, Corynetes [= Necrobia] rufipes, and Dermestes lardarius; tenacity of life of the Tyroglyphus and assumption of Hypopus form by the same; anecdote of Latreille; injury done in cheese by Musca corvina.
- 1704. RILEY, C. V. Pupation of the Nymphalidæ. Abstract: <Psyche, 11 July, 1879, v. 2, pp. 249-251.
 - Notice of prevalent explanations of the manner in which suspended pupa of Lepidoptera support themselves while shedding the larval skin; explanation of the process in *Vanessa antiopa* and *Paphia glycerium*.
- 1705. RILEY, C. V. Fire-flies. <Sci. Amer., 26 July, 1879 [v. 55], n. s., v. 41, p. 49. S.-b. No. 23, p. 104.
 - Definition of "Fire-flies;" description of larva, pupa, and image of *Photinus pyralis*; *Photuris pennsylvanicus*, the commonest species in the more northern states; absence or imperfect development of wings in certain female *Lampyridæ*.

1706. RILEY, C. V. Grape-scale insect, new species. <Pacific Bural Press, 16 August, 1879. S.-b. No. 32, p. 66.

Answer to inquiry of editor of Press; a new coccid, allied to Pulvinaria vitis.

- 1707. RILEY, C. V. Dragon flies. <Sci. Amer., 23 August, 1879 [v. 55],
 n. s., v. 41, p. 113. S. b. No. 23, p. 100.
 Description and natural history of *Libellulidæ*.
- 1708. RILEY, C. V. Failure of tea roses. Habits of Fuller's rose beetle, Aramigus fulleri Horn. <Sci. Amer., 30 August, 1879 [v. 55], n. s., v. 41, p. 129, fig. S.-b. No. 23, pp. 114-116. Reprint: <Gardener's Mo. and Hortic., October, 1879, v. 21, pp. 310-311, fig. S.-b. No. 23, pp. 126-127. Reprint, with additions: <Ann. Rept. [U. S.] Commissioner Agric. for 1878, 22 November, 1879, pp. 255-257, pl. 17, f. 2. See No. 1721 for synopsis of contents.
- 1709. RILEY, C. V. The "Devil's darning needle." <Sci. Amer., 6
 September, 1879 [v. 55], n. s., v. 41, p. 148. S. b. No. 23, p.
 104. Review: <Sci. Amer., 27 September, 1879 [v. 55], n. s.,
 v. 41, p. 194. S. b. No. 23, p. 103½.

Diapheromera femorata winters in the egg-state; W. J. McGee confounds the above with a species of Ranatra; the term "Devil's darning needle" mostly associated in the popular mind with the dragon-flies (Libellulidæ).

- 1710. [RILEY, C. V.] The red-spider on roses. <Sci. Amer., 13 September, 1879 [v. 55], n. s., v. 41, p. 161. S.-b. No. 23, pp. 116-117.
 - Nature, ravages, and means against Tetranychus telarius; habits and transformations of Acarina.
- 1711. RILEY, C. V. Philosophy of the pupation of butterflies. <Sci. Amer. Suppl. No. 193, 13 September, 1879, p. 3069, 3 figs. S.-b. No. 23, p. 125. Reprint: <Science News, 15 September, 1879, v. 1, pp. 346-350. Reprint with changes: <Nature, 16 October, 1879, v. 20, pp. 594-595. <Kosmos, January, 1880, bd. 6, pp. 313-318. <Proc. Amer. Assoc. Adv. Sci. for 1879, 1880, v. 28, pp. 455-463, figs. <Separate: Salem, Mass., July, 1880, 9 pp., 6 figs. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 162-167, figs. 65-70. Abstract: <The Field Nat. and Sci. Student, January, 1883, pp. 179-180. S.-b. No. 42, p. 1. <Bull. Philos. Soc. Wash., 1879, v. 3, pp. 41-43.
 - Illustrates the prevalent explanation of the manner in which suspended chrysalids become attached to their supports; describes and illustrates the more correct process; structure of the terminal joint of the chrysalis especially in Vanessa antiopa, Terias sp., Danais archippus, and Paphia glycerium.
- 1712. RILEY, C. V. Parasites of the cotton-worm. < Ca. Ent., September, 1879, v. 11, pp. 161--162.
 - List of parasites of Aletia argillacea [=xylina]; descriptions of Trichogramma pretiosa n. sp. [p. 161], Cirrospilus [= Tetrastichus] esurus n. sp. [p. 162], and Tachina aletia n. sp. [p. 162].

- 1713. RILEY, C. V. [Insect injurious to junipers.] <Ca. Ent., September, 1879, v. 11, p. 177.

 Junipers on Long Island injured by Dapsilia rutilana.
- 1714. RILEY, C. V. The imported cabbage-worm in the South. <Farmer's Review, September, 1879. S.-b. No. 23, p. 106.

 Spread of *Pieris rapa* westward and southward; its present distribution; southern limit of distribution of *Deryphora* 10-linests.
- 1715. BILEY, C. V. The shedding of the tracheæ and double cocoons. <Amer. Nat., October, 1879, v. 13, p. 652.

 Review of two notes by E. Potts.
- 1716. RILEY, C. V. Leaf-galls on the grape-vine. <N. Y. Tribune, 1
 October, 1879. S.-b. No. 23, pp. 117; 173. See: <Kansas
 Farmer, 26 November, 1879. S.-b. No. 26, p. 9. <West. Rural,
 10 December, 1879.
 - Habits, ravages, reproduction, and distribution of Phylloxera vastatrix.
- 1717. RILEY, C. V. The Croton bug as a library pest. <Library Journal, September-October, 1879, v. 4, p. 376.

 Letter to Weston Flint; ravages of and means against Blatta [= Ectobia] germanica.
- 1718. RILEY, C. V. The Ailanthus silk-worm, Attacus (Samia) cynthia.

 <Science News, 15 October, 1879, v. 1, pp. 377-383. Extract:

 <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, pp. 56-58, fig. 16.

 <Farmer's Review, 8 January, 1880, v. 4. S.-b. No. 23, pp. 167-168.
 - Description of eggs, larvæ, and images of Samia [=Attacus] cynthia; history, food-plants, acclimation, and artificial rearing of the same, and of Samia [=Attacus] ricini; differences between the eggs, larvæ, and cocoons of the two species; availability of these and other species for silk-culture; superiority of Sericaria mori for this purpose; occurrence of parthenogenesis and of retardation of development in Bombycidæ.
- 1719. RILEY, C. V. The cotton-worm. Letter from Prof. C. V. Riley on some recent cotton-worm articles in the News. <Galveston [Tex.| Daily News, 24 October, 1879, v. 38, No. 185, p. 2. S.-b. No. 23, pp. 164–165. Reply: <Ibid., 31 October, 1879, No. 191, p. 4.
 - Critical review of W. J. Jones's "The cotton caterpillar," and of Investigator's "The cotton-worm;" denies that Aletia argillacea [=xylina] hibernates in the pupa state; author not interested in entomological discoveries for mercenary ends.
- 1720. RILEY, C. V. [Spread of *Pieris rapæ* into Alabama.] < Ca. Ent., October, 1879, v. 11, p. 196.
 - Pieris rapæ now quite common in Alabama; it has reached Selma but not Mobile.
- 1721. RILEY, C. V. Report of the entomologist. <Ann. Rept. [U. S.]

 Commissioner Agric. for 1878, 22 November, 1879, pp. 207-257,

 + 1 p. expl. of pl., pls. 1-7. Separate: <Washington, Octo-

1721.	RILEY, C. V.—Continued.	
	ber, 1879, pp. 52 + 1 p. expl. of pl., 7 pl. Notice: <garden 1879,="" 21,="" 349-350.="" and="" hortic.,="" mo.="" november,="" pp.="" sb.<="" td="" v.=""><td></td></garden>	
	23, p. 29.	NU
	LETTER OF SUBMITTAL	20
	Insects affecting the cotton plant Report of special investigations carried on in 1878, 210—Organization and personnel of the commission of inquiry, 210—Circular letter of inquiries, 210—Report of A. R. Grote, 213—Seasons, hibernation, and migrations of Aletia argillacea [= xylina], 213—Discovery of parasites, 214—Food-plants of the imago, 214.	210
	THE SILK WORM: A BRIEF MANUAL OF INSTRUCTIONS FOR THE PRO-	
	DUCTION OF SILK. [A reprint of special report No. 11.] Culture of Sericaria mori in the United States, 215—Profits of producing cocoons, 217—Profits of reeling, 217—Profits in raising eggs, 218—Nature of the silkworm, 218—The egg, 219—The larva, 219—The cocoon, 220—The chrysalis, 221—The moth, 221—Enemies and diseases, 221—Muscardine, 221—Pébrine, 222—Other diseases, 223—Varieties or races, 223—Wintering and hatching the eggs, 224—Feeding and rearing the worms, 225—Preparation for spinning, 228—Gathering the cocoons, 229—Choking the chrysalis, 229—Egg-laying; reproduction, 230—Reeling, 232—Food-plants, 235—Glossary of terms used, 236.	215
	THE GRAPE PHYLLOXERA Importance of American grape-culture, 237—Grafting of vines, 237—Distribution of Phylloxera, 238—Hopleseness of means against, 238.	237
	Notes on the apple-worm	238
	THE WESTWARD PROGRESS OF THE IMPORTED CABBAGE-WORM Spread of <i>Pieris rapæ</i> , 239—Means against the same, 239.	239
	CHAPIN'S APPLE-LEAF SEWER, Phozopteris nubeculana Clem	239
	THE THICK-THIGHED WALKING-STICK, Diapheromera femorata Say Vernacular names, 241—Nomenclature, 241—Characters, 241—Destructive powers, 241—Habits and natural history, 243—Its appearance every alternate year, 244—Natural enemies, 244—Remedies, 245—Description of egg, 245—The larva, 245.	241
;	THE GREAT ELM-LEAF BEETLE, Monocesta coryli, Say*	245
	THE JUNIPER WEB-WORM, Dapsilia rutilana, Hitbn Importation, 247—Distribution in Europe, 247—Habits, 247—Remedies, 248—Description of larva, 248; of pupa, 248; of imago, 248.	247

^{*}Translation: \langle Acker- und Gartenbau-Zeitung, 1 June, 1880, v. 11, p. 87, fig. 8.-b. No. 23, p. —.

1721. RILEY, C. V.—Continued.	
THE CLOVER-ROOT BORER, Hylesinus trifolii, Müller *	248
ters of <i>H. opaculus</i> and <i>H. trifolii</i> , 250—Enemies, 250—Description of larva, 250; of pupa, 250.	
THE CLOVER-SEED MIDGE, Cecidomyia leguminicola, Lintner	250
THE APPLE COLEOPHORA, Coleophora malivorella, n. sp.†	253
FULLER'S ROSE-BEETLE, Aramigus fulleri Horn History, 255—Distribution, 255—Ravages, 255—Habits and characters of imago, 256—Remedies, 257—Description of egg, larva, and pupa, 257.	258

1722. RILEY, C. V. [Parasites bred from the cotton-worm.] <Ca. Ent., November, 1879, v. 11, p. 205.

Nine species of parasites have been bred from Aletia argillacea [=xylina].

1723. RILEY, C. V. The bee-moth. <N. Y. Tribune, 1879. Reprint: <Farmer's Review, 3 January, 1880. S.-b. No. 23, p. 158.

Description of larva and imago, habits, ravages, and means against Galleria

1724. RILEY, O. V. The silk-worm; being a brief manual of instructions for the production of silk. <Special Report No. 11 [U. S.], Dept. Agric., Washington, 1879, pp. 31, 8 figs. Second edition: <Washington, 1882, 37 pp., 8 figs. Third edition: <Washington, 1883, 37 pp., 8 figs. Fourth edition: Fifth edition: Sixth edition: <Bull. No. 9 Div. Ent. U. S. Dept. Agric., 11 July, 1886, 65 pp., 29 figs., 2 pl. Seventh edition: <Ibid., April, 1888.

Treats of Sericaria mori. See No. 1721 for synopsis of contents.

1725. RILEY, C. V. London purple as an insecticide. <Farmer's Review, 29 January, 1880, v. 4, No. 5, p. —.

1726. [RILEY, C. V.] [The food-habits of thrushes.] < Amer. Ent., January, 1880 [v. 3], n. s., v. 1, pp. 2-3.

Notice of S. A. Forbes' "The food-habits of thrushes;" incompleteness of our knowledge of the food-habits of birds and of *Carabida*; author's opinion favorable to the birds.

1727. [BILEY, C. V.] The grape Phylloxera in California. <Amer. Ent., January, 1880 [v. 3], n. s., v. 1, p. 3.

Phylloxera vastatrix as destructive to Vitis vinifera in California as in Europe; the winged female supposed not to have appeared yet in California; speculations upon this modification of habit, if true.

oereana.

^{*}Translation: <Acker- und Gartenbau-Zeitung, 1 August, 1880, v. 11, p. 119, 2 figs. S.-b. No. 23, p. —. Extract: <Prairie Farmer, 22 November, 1879, v. 50, p. 370. S.-b. No. 23, pp. 168-169. <Amer. Rural Farmer, 18 December, 1879. S.-b. No. 45, pp. 23-24.

[†]Translation: <Acker- und Gartenbau-Zeitung, 1 September, 1880, v. 11, p. 135, fig. S.-b. No. 23, p. 182.

- 1728. [RILEY, C. V.] On the hibernation of the cotton-worm, Aletis argillacea Hübn. <Amer. Ent., January, 1880 [v. 3], n. s., v. 1, pp. 6-11, figs. 3-5; p. 15.
 - Advance print from Bull. No. 3 U. S. Ent. Commission, pp. 24-31. See No. 1736 for synopsis of contents.
- 1729. [RILEY, C. V.] [Inquilines in galleries of the common white-ant.] <Amer. Ent., January, 1880, [v. 3], n. s., v. 1, p. 15.
 - Trichopsenius depressus and three undescribed Aleocharini found inquilinous in galleries of Termes flavipes in Texas by E. A. Schwarz; the Trichopsenius previously known by a single specimen from Georgia.
- 1730. [RILEY, C. V.] Large white scale on acacias, etc. < Amer. Ent. January, 1880 [v. 3], n. s., v. 1, p. 20.

Occurrence and ravages of *Dorthesia characias*? in California and southern Africa; characters of the insect.

- 1731. [RILEY, C. V.] [Ravages of moths in cushions.] < Amer. Ent., January, 1880 [v. 3], n. s., v. 1, p. 20.
 - Conclusions of discussion by German Society of Railway Companies as to the best method of preventing the ravages of moths in the cushions of railway carriages.
- 1732. [RILEY, C. V.] [Grape Phylloxera in Geelong, Victoria.] < Amer. Ent., January, 1880 [v. 3], n. s., v. 1, p. 20.

 Occurrence of Phylloxera vastatrix in Geelong, Victoria.
- 1733. RILEY, C. V. Borers in black-ash: Fall web-worm: Apple-tree insects. <Amer. Ent., January, 1880 [v. 3], n. s., v. 1, pp. 22-23.
 - Black-ash trees in New York killed by the borings of Parandra brunnea; ash trees bored also by Trochilium [= Fatua] denudata; the leaves of the same and other trees destroyed by Hyphantria textor [= cunea]; apples injured by the "gimlet-worm"; habits, ravages of, and means against Bucculatrix pomifoliella; figure of cocoons and image of this moth.
- 1734. [RILEY, C. V.] Skippers injuring smoked hams. <Amer. Ent.,

 January, 1880, [v. 3], n. s., v. 1, pp. 23-24.

 Answer to communication from E. A. Smith; hams injured by larvæ of Fiephila case;; means against these and other insects injuring hams.
- 1735. [RILEY, C. V.] Tipula eggs in the stomach of the cat-bird. <Amer. Ent., January, 1880 [v. 3], n. s., v. 1, p. 24.

 Answer to communication from S. A. Forbes; description of eggs and method of oviposition of Tipula trivittata?
- 1736. RILEY, C. V. The cotton-worm. Summary of its natural history, with an account of its enemies and the best means of controlling it; being a report of progress of the work of the commission. <Bull. No. 3, U.S. Ent. Commission, 28 January, 1880, 144 pp., 1 pl., 84 figs. Extract: <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, pp. 67-68, figs. 19-20. <Farmer's Review, January, 1880, 2 figs. S. b. No. 23, pp. 163-164.

	,	
•	Text of circular sent out by the commission, 1—History of the investigation, 1—Need thereof, 2—Early work of Townend Glover, Thomas Affleck, and D. L. Phares, 2—Special report ordered by Congress, 3—Character of the Bulletin, 3—Division of the cotton-belt in the southern or hibernating and the northern or non-hibernating portions, 4—Practical aim of the commission, 4—Persons who have assisted, 4.	
	1. DESTRUCTIVENESS OF THE WORM Losses occasioned by it Increase of injury westward, 7—Least injury in northern portion of belt, 7—Tabular statement of loss for past 15 years, 8.	7
	2. POPULAR AND SCIENTIFIC NAMES FOR THE INSECT	8
	3. CHARACTERS AND STAGES OF THE INSECT The egg The worm always hatched from an egg, 9—Where the egg is laid, 9—Number of eggs to a leaf, 10—All eggs perish during winter, 10—Other eggs mistaken for those of Aletia, 10.	9 9
	The worm or larva	10
•	The chrysalis	12
•	The moth or imago	13
	4. Time elapsing from one generation to another It varies greatly, but averages about one month, 16.	16
	5. TIME OF YEAR WHEN THE FIRST WORMS APPEAR Prevailing opinions erroneous, 16—The worms hatch in April, 16—First worms always in small numbers, 17—The third generation usually called the first, 17.	16
	6. CONDITION OF SOIL AND PLANT CONNECTED WITH THE APPEARANCE OF THE FIRST WORMS The worms appear first on low, moist, and rich lands, 18—Few exceptions to the rule, 18—The plant must be well advanced and luxuriant, 18—Some shelter generally near, 19—Reoccurrence year after year in the same spots, 19—Theory explaining first appearance under the circumstances observed, 19.	18
	7. WET WEATHER FAVORS THE DEVELOPMENT OF THE WORMS Humid and hot atmosphere most favorable, 20—Heavy storms destroy, 20—Effect of dryness, 21—Indirect influences of wet weather, 21—It prevents the working of the worm's natural enemies, 21.	20

1736. RILEY, C. V.—Continued.	
8. Number of annual generations	21
9. MIGRATIONS AND POWER OF FLIGHT OF THE MOTH The moth has great power of wing, 23—It flies many hundreds of miles north of cotton belt, 23—Appearance of the worm on cotton first planted great distances from any other cotton thus accounted for, 23—Migrating habit developed only in later months when they become numerous, 24.	21
Different opinions and beliefs, 24—Difficulty surrounding the question, 24—Erroneous conclusions easily drawn, 25—Fallacy of the belief that the chrysalis winters underground, 25—26—Ability of the moth to survive the winter, 26—Other moths easily mistaken for it, 27—Theory of annual introduction of the species from some exotic country, 28—Arguments for and against the theory, 29, 30—Hibernation of the moth in the southern portion of the belt substantiated, 30—But a very small proportion of those which abound in the fall live to perpetuate their kind the next spring, 31—Localities where the moth hibernates, 31.	24
Vertebrates	31 33
Introduction of English sparrow, 33. Invertebrates	33 33
34—Beetles which prey upon the worm and others which presumably do so, 35—Enemies among the half-winged bugs or Heteroptera, 36—Among two-winged flies or Diptera, 36—Among nerve-winged insects or Neuroptera, 37.	
Parasites Previous belief that none preyed on Aletia, 38—Infesting the egg, 38—Infesting and issuing from the worm, 39—Flesh flies and Tachina flies, differences between them, 40-42—Issuing from the chrysalis, 42—A parasite known but not named in 1852—Ten distinct parasites infesting Aletia, and two other undetermined species, 47.	38
12. Remedies: Means of coping with the insect	47
Prevention	48
Mechanical means of killing the worms	49
Simple methods of brushing and dragging, 50. Machines for brushing off the worms	50

12. Remedies: Means of Coping with the Insect—Continued. The Ewing brushing machine, 50—The Helm brushing machine, 51—The Wood Smith brushing machine, 52—The Iske brushing machine, 53—Use of kerosene and coal oil with shallow pans or cloth frames, 54.	,
Poisoning the worms. Impetus given to invention by the use of mineral poisons, 54— Value of concerted action, 54—General principles to be observed in poisoning, 55—Dry vs. wet applications, 55—Advantage in being prepared, 55.	54
Arsenieal compounds	55
London purple	60
Pyrethrum powder Its origin and nature, 62—Its remarkable insecticide qualities, 62—Most effectual on the cotton-worm, 63—Perfectly harmless to man, 63—The extract of the plant has the same virtue, 63—Growth of the plant in California, 63—Experiments conducted, 64—The alcoholic extract very potent, 64—Probable cost of using it, whether as powder or extract, 65—Highly encouraging prospects from its use, 65.	62
Kerosene oil	65
Cotton-seed oil	66
Carbolic acid	67
Sulphur Extracts or decoctions from various plants Yeast ferment; fungus infection Proposition by Dr. Hagen, 68—Dr. Bail's experiments, 68—Not in accord with the leading mycologists, 68—Experiments with beer-mash, 69—With beer-mash and yeast ferment, 70.	67 68
Machines and contrivances for powdering	74

12, Remedies: Means of Coping with the Insect—Continued. and sprinkler, 78—The Davis duster, 78—The Levy duster, 79—The Taylor duster, 80—The Allen duster, 81—The Young	
duster, 81—The Goodheart duster, 81. Machines and contrivances for sprinkling and spraying	81
Machines for applying vapors	97
The Steinmann vaporizer, 97—The Pearl vapor generator, 99. Destruction of the moth	100
Results so far not encouraging, 100—Action must be taken early in the season, 100.	100
Lamps for attracting the moth	100
Usually resorted to when too late to do practical good, i00-	
Fires, 100-Trouble of keeping them up throughout the night,	
101—Beneficial insects killed by lamps and fires, 101—Ad-	
vantage of lamps, 101—The Garrett lamp, 101—The Binkley	
lamp, 102—The McQueen lamp, 103—The Rigel lamp, 103—	
The Walker lamp, 103—Simple form of lamp, 104—The Pugh	
lamp, 104—Lamps in motion, 105—Le Blanc's cotton-moth	
destroyer, 105—The Fordtran cotton-moth destroyer, 105. Poisoned sweets and fluids as means of destroying the meth	106
Sweets attractive, 106—Ripe fruits and melons often suffer, 106—	100
Baits that allure and kill, 106—Experiments with them, 107—	
Moths less attracted during summer than in autumn, 107—Lit-	
tle inducement to use baits in autumn, 108—They are most	
beneficial in spring, 108-White rags, 108-Theory of Dr. J. L.	
Lupton, 108.	
APPENDIX	109
Answers to circular from West Feliciana Parish, Louisiana, 109-	
From Evergreen, Ala., 111- From Henderson, Tex., 113-From	
Hempstead, Tex., 115-Different views as to hibernation and	
mode of perpetuation of the insect, 116—Answers to circular	
from Livingston, Ala., 117-From Parish of Concordia, La.,	
119—From Natchitoches, La., 122—From Jackson, Miss., 126—	
From Larissa, Tex., 127—From Perry County, Ala., 128—	
From Walterborough, S. C., 128—From Jamestown, Fla., 130—	
From San Antonio, Tex., 131—From Kirkwood, Miss., 133—	
From Woodville, Miss., 135.	D-1-

1737. [RILEY, C. V.] The 17-year Cicada in Iowa. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, pp. 25-26.

Review of C. E. Bessey on the 17-year Cicada in Iowa; limits of the broods of 1854-1871, 1861-1878, and 1862-1879 of Cicada [= Tibicen] septendecem in Iowa; occurrence of the last brood in Missouri; comparison of the distribution of these broods with the distribution of timber trees.

- 1738. [RILEY, C. V.] Vegetal-feeding ground-beetles. < Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26.
 - Notes from various sources upon the phytophagous habits of Carabidæ.
- 1739. [RILEY, C. V.] The pear-leaf blister. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26. Notice: <Ibid., March, 1880 [v. 3], n. s., v. 1, p. 74.
 - Review of T. J. Burrill's "The pear-leaf blister"; ravages of Typhlodromus pyri on leaves of pear-tree; characters of this mite; noxious habits of Acarina, especially of Phytoptus.
- 1740. [RILEY, C. V.] Fuller's rose-beetle in California. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26.

 Occurrence and ravages of Aramigus fulleri in California.
- 1741. [RILEY, C. V.] Lepidium vs. bed-bugs. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26.

 Notice of S. M.'s "Lepidium, the bed-bug destroyer;" Acanthia lectularia said to be attracted to and killed by Lepidium sp.
- 1742. [RILEY, C. V.] Aniseed and grain weevils. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 26.

 Notice of reports that Calandra granaria were attracted from grain to aniseed and killed by it.
- 1743. RILEY, C. V. Two valuable insecticides. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, pp. 41-45.

 From Bull. No. 3, U. S. Ent. Commission, pp. 60-65. See No. 1736, for synopsis of contents.
- 1744. [RILEY, C. V.] Use of buckwheat to destroy insects. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 48.

 Extract from "The Farmer" with remarks on the planting and plowing in of buckwheat as a means against injurious insects.
- 1715. [RILEY, C. V.] Tipula eggs in stomach of cat-bird. <Amer. Ent., .
 February, 1880 [v. 3], u. s., v. 1, p. 50.

 Answer to communication from S. A. Forbes; two kinds of eggs found in stomach of cat-bird; these birds eat gravid females of Tipula.
- 1716. [RILEY, C. V.] Beetles supposed to be feeding on wheat. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 50.</p>
 Answer to letter from P. H. M.; larva of Cryptorhopalum sp. supposed to have injured grains of wheat, was probably feeding on the remains of true grain insects; food-habits of larvæ of Dermestidæ.
- 1747. RILEY, C. V. The apple-twig borer. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, pp. 50-51, figs. 11-12.

 Answer to letter from H. G. Wolcott; habits, food-plants, and means against Amphicerus bicaudatus: figure of male and female image and of bored apple-twigs; habits and food-plants of Sinoxylos basillars and of Bostrichus.
- 1748. [RILEY, C. V.] Stinging caterpillars. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 51.

 Answer to letters from T. Pollard and G.W. S.; food-plants, general appearance, vernacular name, urticating properties and description of image of

Lagoa opercularis.

- 1749. RILEY, C. V. A new genus of *Proctrotrupidæ*. <Amer. Ent., February, 1880 [v. 3], n. s., v. 1, p. 52, figs. 13-14. Review: <1bid., December, 1880 [v. 3], n. s., v. 1, p. 293.
 - Description and figures of Didictyum [=Hexaplasta] zigsag n. g. et n. sp. reared from chrysalis of Aletia argillacea [=zylina].
- 1750. RILEY, C. V. Agricultural advancement in the United States.
 <Farmer's Review, 1880, v. 4: 4 March, p. 158; 11 March, p. 174.</p>
 S.-b. No. 23, pp. 153-158. Notice: <Ibid., 4 March, 1880, p.—.
 S.-b. No. 23, p. 150. Reprint: <Journ. Amer. Agric. Assoc., April, 1881, v. 1, pp. 47-54. Separate of reprint: <August, 1881, pp. 47-54.
 - Suggestions for the organization of an agricultural association; comparison of the relations of government to the advancement of agriculture in England, Germany, France, and the United States; necessary changes in the United States Department of Agriculture.
- 1751. RILEY, C. V. Parasites of the plum Curculio. <Farmer's Review, 4 March, 1880, v. 4, p. —, 2 figs. S.-b. No. 23, pp. 169–170.
 - Figures of larva, pupa, cocoon, and images of Sigalphus curculionis; habits, variations, usefulness, and dissemination of the same; habits, colors, and seasons of Porison [= Thersilochus] conotracheli.
- 1752. [RILEY, O. V.] Trapping the carpet-beetle. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, pp. 53-55, fig. 15.

 Figures larva, pupa, and image of Anthronus scrophularia; food-plants and habits of image and ravages of larva; importation, distribution, vernacular names of and means against the same.
- 1753. [RILEY, C. V.] Silk-worm eggs: Silk culture. Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 55.

 Method of obtaining eggs of Serioaria mori and information on silk culture.
- 1754. [RILEY, C. V.] [Hickory Scolytus.] < Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 58.

 Occurrence of Scolytus quadrispinosus in Washington Territory.
- 1755. [RILEY, C. V.] [Tenacity of life.] <Amer. Ent., March, 1880 [v.3], n. s., v. 1, p. 68.

 Resistance of Cleonus sp. to the influence of various insecticides.
- 1756. [RILEY, C. V.] [Danais archippus.] < Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 73.

 Occurrence of a colorational variety on the island of Antigua.
- 1757. [RILEY, C. V.] Reports of the U. S. Entomological Commission. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 73.

 Method of obtaining the first report of the U. S. Entomological Commission.
- 1758. [RILEY, C. V.] [Habits of the cotton-moth.] < Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 74.
 - Request for information on the habits of the imago and the food-plants of the larva of Aletia argillacea [=xylina] during March and April.
- 1759. [RILEY, C. V.] Typhlodromus pyri. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 74.

 Reference to early account of the pear-leaf blister-mite.

- 1760. [RILEY, C. V.] Food-habits of ground-beetles. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 75.
 - Note on request of S. A. Forbes for Carabidæ found in situations suggesting herbivorous habits.
- 1761. [RILEY, O. V.] Moths and butterflies caught by the tongue. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 75.
 - Notice of several articles on the capture of Noctuida and Sphingida by the flowers of Physianihus albens; capture of Sphingida by the flowers of Nerium oleander and Enothera grandiflora, and of Syrphus sp. by the flowers of Bidens chrysanthemoides.
- 1762. [RILEY, C. V.] [Queen bees in the mails.] Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 75.
 - Circumstances under which queen bees may be sent by mail; exclusion of pinned insects from the same.
- 1763. [RILEY, O. V.] Common tiger-beetle. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, pp. 77-78.

Answer to inquiry of J. L. Seney; habitat, distribution, and food-habits of Cioindela repanda.

- 1764. [RILEY, C. V.] Gall on *Pelargonium*. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 78.
 - Answer to letter from Mary Treat; gall at base of *Pelargonium* probably made by a mite; the *Podura* found in it were undoubtedly feeding on diseased gall-tissue.
- 1765. [RILEY, C. V.] Catalogues and monographs of insects. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 78.
 - Answer to letter of P. S. B.; mention of monographic works and catalogues of North American insects.
- 1766. [RILEY, O. V.] Leaf-hoppers injuring wheat fields. <Amer. Eut., March, 1880 [v. 3], n. s., v. 1, p. 78.
 - Answer to letter of R. L. B.; habits and ravages of Cicadula exittosa, Diedrocephala flaviorps, and Jassus sp. in the southern United States, and of Jassus sexuotatus in Europe; means against the same.
- 1767. RILEY, C. V. A new leaf-hopper injurious to small grain. <Amer. Ent., March, 1880 [v. 3], n. s., v. 1, p. 78.
 - Description of *Diedrocophala flavicops* n. sp. injurious to wheat and cats in Texas in 1876.
- 1768. RILEY, C. V. The bird question dispassionately considered. <Farmer's Review, 1 April, 1880, v. 4, p. 211. S.-b. No. 23, pp. 147-148.
 - Extract from E. Perris's "Birds vs. Insects," with introductory; conclusions as to the value of birds in agriculture as destroyers of noxious insects.
- 1769. RILEY, C. V. The cotton-worm in the United States. <Amer. Ent., April, 1880 [v.3], n. s., v.1, pp. 93-95. Reprint, with slight changes: <Proc, Amer. Assoc. Adv. Sci. for 1879, 23 February, 1881, v. 28, pp. 464-466. Separate of reprint: <Salem, Mass., August, 1880, 3 pp.
 - Date and manner of first appearance of larvæ of Aletia argillacea [=xylina]; number of annual generations and the existence of parasites upon it; his-

- 1769 RILEY, C. V.—Continued.
 - tory and refutation of the annual immigration theory; division of the cotton belt into regions wherein, respectively, A. argillaces [=xylina] is permanent and temporary; food-habits and enemies of the same.
- 1770. [RILEY, C. V.] The migrations of butterflies. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, pp. 100-102, figs. 34-35.
 - Migratory flights of Lepidoptera supposed to be due to excessive multiplication of individuals and to a true migratory instinct; directions, times, and extent of migrations of Danais archippus; explanation of the migratory instinct; distribution of D. archippus and Pyrameis cardsi; figures the former and its manner of clustering.
- 1771. [RILEY, C. V.] Gouty gall on blackberry and raspberry canes. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 107.
 - Answer to letters of P. S., T. A. C., and J. W.; seasons, ravages of and means against *Agrilus ruficollis*.
- 1772. [RILEY, C. V.] Sowing cotton seeds in hot-beds and transplanting as a means of preventing injury from the cotton-worm. < Amer. Ent., April, 1883 [v. 3], n. s., v. 1, p. 107.
 - Impracticability of W. J. W.'s suggestion as indicated in title.
- 1773. [RILEY, C. V.] Not the cotton-moth. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 107.
 - Answer to inquiry of D. B. Woodbury; food-plants of Telype velleda.
- 1774. [RILEY, C. V.] Moths caught in Alabama: Muscle-shaped barklouse on apple-trees South Amer. Ent., April, 1880 [v. 3], n. s., v. 1, pp. 107–108.
 - Mentious several insects received from J. F. Bailey, of Marion, Ala.; larva of Papilio philenor feeds on Aristolochia; Bombus virginious 2 robs hives of Apis mellifica; Mytilaspis pomicorticis [=pomorum] not before received from so far south; means against the same.
- 1775. [RILEY, C. V.] Chrysalides supposed to be those of Aletia. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 108.
 - Answer to letter of W. J. Jones; pupse of Agnetic incrmis [=saucia] and Prodonia lineatella from cotton field.
- 1776. [RILEY, C. V.] Apple-twig borer. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 108.
 - Answer to inquiry of T. V. M.; occurrence and habits of Amphicerus bicandatus.
- 1777. [RILEY, C. V.] Clover-weevil. <Amer. Ent., April, 1880 [v. 3], n. s., v. 1, p. 108.
 - Answer to inquiry of W. W. F.; several "weevils" infest clover-seeds; Hylesinus trifolii infests the roots and lower part of the stem.
- 1778. [RILEY, C. V.] Effects of cold applied to the chrysalides of butterflies. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 110-111.

 Notice of the experiments of W. H. Edwards; effects of hibernation in the chrysalis and image states.
- 1779. [RILEY, C. V.] Moth issuing from a larva. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 114.
 - Notice of J. J. Weir on the issuance of *Orgyis* sp. from the larva without passing through the pupal stage.

- 1780. [RILEY, C. V.] The rose-slug. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 115-116, figs. 42-43.
 - Description of eggs, larvæ, and imago, history and means against Selandria [= Monostegia] rosæ; figures, egg, larvæ, imago, with details of structure, and injured leaf.
- 1781. [RILEY, C. V.] Dr. Asa Fitch. <Amer. Ent., May, 1880 [v. 3], n. s, v. 1, pp. 121-123.
 - Biographical sketch of Asa Fitch, with notice of his collection of insects and his writings.
- 1782. [RILEY, C. V.] Flea-beetle on young tobacco plants. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 123.
 - Leaves of Nicotiana tabacum eaten by Epitrix hirtipennis [= Crepidodera parvula] and probably by E. [= C.] cucumeris; leaves of Solanum eaten by the first named and by E. [= C.] brevis; distribution of these and other species of Epitrix [= Crepidodera].
- 1783. [RILEY, C. V.] The proboscis of the common house-fly. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 125. Review of paper by G. Macloskie.
- 1784. [RILEY, C. V.] Notes on South American Lepidoptera. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 125-126.

 Report of meeting of the Entomological Society of London; length of proboscis of and presence of scent organ in Sphingida; perception of colors in
 - scis of and presence of scent organ in Sphingidæ; perception of colors in Rhopalocera; secondary sexual character in Callidryas and other genera; according to R. Meldola the proboscis of Macrosila cluentius is 23.5cm (94 inches) long.
- 1785. [RILEY, C. V.] Floating apiaries. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 126-127.

 Notice of an unsuccessful attempt to increase the product of an apiary by floating it southward late in the season and back to the north in the spring, on the Mississippi River.
- 1786. [RILEY, C. V.] American Staphylinida wanted. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 127.
 - Notice of A. Fauvel's works on Staphylinida and his request for additional material.
- 1787. [RILEY, C. V.] Probable parthenogenesis in the Hessian fly. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 127.
 - Review of paper by H. A. Hagen; Cecidomyia destructor less injurious than formerly in the older wheat-growing regions, its area of abundance having moved westward with the westward extension of wheat culture.
- 1788. [RILEY, C. V.] Raspberries destroyed by weevils. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 127.

 Ravages of Otiorhynchus picipes in England; warning against its introduction
 - Ravages of *Ottorhynchus picipes* in England; warning against its introduction into the United States.
- 1788a. [RILEY, C. V.] Cotton culture and the insects affecting the plant at Bahia, Brazil. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 128-129.
 - Letter of R. A. Edes, with remarks on specimens sent.

- 1789. [RILEY, C. V.] Hemispherical larva at bottom of ant hill. < Amer. Ent., May, 1880 | v. 3], n. s., v. 1, p. 129.

 Answer to inquiry of D. S. Sheldon; habitat of larva of Microdon globosus.
- 1790. [RILEY, C. V.] Chrysalides dug up in cotton field, mistaken for those of the cotton-worm. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 129.

 Answer to letter of R. Worrel.
- 1791. [RILEY, C. V.] Aleurodes on Oxalis. < Amer. Ent., May, 1880
 [v. 3], n. s., v. 1, pp. 129-130.

Answer to letter of S. A. Conrad; description of all stages of Aleurodes sp. from leaves of Oxalis sp.

- 1792. [RILEY, C. V.] Larvæ in stomach of black-bass. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 130.

 Answer to inquiry of S. A. Forbes; two coleopterous larvæ from the stomach of *Mioropterus salmoides*; one a dytiscid, the other perhaps a dascyllid.
- 1793. [RILEY, C. V.] Insects from stomach of rock-bass. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 130.

Answer to letter of S. A. Forbes; coleopterous larva from the stomach of common sun-fish (*Lepiopomus pallidus*) and rock-bass (*Bocous lineatus*).

- 1794. [RILEY, C. V.] New enemy to sugar-cane. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 130, fig. 48.
 - Answer to letter of D. Th.; ravages of Ligyrus rugiceps upon sugar-cane, maize, and grasses; recommends use of lamp and kerosene pan, and figures one form of such apparatus.
- 1795. [RILEY, C. V.] Parasites of the plum Curculio. <Amer. Ent., May, 1880 [v. 3], n. s., v. 1, pp. 131-132, figs. 49-50.

 Answer to letter of W. S. Barnard; quotes from 3d Ann. Rept. State Ent. Mo., pp. 24-26; habits of Sigalphus curculionis; habits, varieties, and figures of all stages of the same; food-habits of Semasia [= Grapholitha] prunivers on which the Sigalphus is parasitic.
- 1796. [RILEY, C. V.] Blister-beetles from New Mexico. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 132.

 Answer to inquiry of J. M.; Macrobasis albida common in the Southwest and valuable as a vesicant; Diplotaxis not known to be a vesicant.
- 1797. [RILEY, C. V.] Abnormal cocoon. Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 132.

 Answer to letter of W. S. Barnard; occurrence of Callosamia [= Attacus]

 promethea and Orgyia antiqua at Ithaca, N. Y.; description of abnormal cocoon of the latter found on sugar-maple.
- 1798. [RILEY, C. V.] Insects found about orange-trees. < Amer. Ent., May, 1880 [v. 3], n. s., v. 1, p. 132.
 - Answer to inquiry of J. S. Barnwell; list of insects from Darien, Ga.; most of them have no relation to orange trees; notes on Aphis sp., Chilocorus birulnerus, Syrphus sp., Neoclytus erythrocephalus, Drasterius amabilis, Platynus punctiformis, Forficula sp., Gryllotalpa borealis, and Psoous venosus.

- 1799. [RILEY, C. V.] Luperus brunneus, Crotch. < Amer. Ent., May, 1880, [v. 3], n. s., v. 1, p. 132.
 - Luperus noxius Riley MSS., is a pale form of L. brunneus; description of the typical and other forms of the same; distinctive characters of L. brunneus and L. morulus from other North American species and from each other; characters of the genus Calomicrus.
- 1800. [RILEY, C. V.] Professor Riley on army-worms. <N. Y. Weekly Sun, 20 June, 1880. S.-b. No. 26, p. 89.

 Interview with reporter; summary of present knowledge of Lencania unipuncts.
- 1801. RILEY, C. V. Cotton-caterpillars. <Selma [Ala.] Times, 25 June, 1880.
- 1802. RILEY, C. V. Notes on our commoner insects. <Amer. Eut., June, 1880 [v. 3], n. s., v. 1, pp. 133, 134, figs. 51-52.

 Description and figures of larva, pupa, cocoon, and image of Arctia [=
 Pyrrharctia] isabella; habits, seasons, food-plants, parasites, vernacular names and variations of the same; figure of Ophion macrurum; descriptions of Ichneumon carulous, I. signatipes, and Trogus obsidianator; description of larva of Ecpantheria scribonia.
- 1803. BILEY, C. V. The white-grub fungus. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, pp. 137-140, figs. 53-55.</p>
 Description and figures of Torrubia ravenelii and of its fructification; forms in which it occurs on larvæ of Lachnosterna quercina [= fusca]; list of some
- articles upon it; its synonymy and distribution; list of related species and their distribution; occurrence of related species on other insects.

 1804. RILEY, C. V. The true and the bogus Yucca moth, with remarks
- on the pollination of Yucca. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, pp. 141–145.
 - History of observations on *Prodoxus decipiens*; habits, parasite, and foodplants of the same; evolution of *Prodoxus* and *Pronuba*; criticism of errors
 resulting from mistaking *Prodoxus decipiens* for *Pronuba yuccasella*; failure
 of *Yucca angustifolia* to become fertilized in the absence of *Pronuba yuccasella*; *Hyponomeuta quinquepunctella* Chambers is the same as *Prodoxus decipiens*.
- 1805. [RILEY, C.V.] Intermittance of phosphorescence in fire-flies.

 Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 146.

 Criticism of a discussion before the London Entomological Society; the winged images of Lampyrida can intermit their light; the larva and wingless
- 1806. [RILEY, C. V.] Grain Aphis vs. rust. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 147.

images can suppress their light.

- Ravages of Aphis avenæ [Nectarophora granaria] and of rust on wheat and oats in Georgia; the work of aphides promotes the growth of some kinds of rust.
- 1807. [RILEY, C. V.] Mold and Phylloxera. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 147.
 - Criticism of statements by Rommier; development of mycelium on phylloxerized roots does not result in the destruction of the *Phylloxera*.

- 1808. [RILEY, C.V.] Infecting Phylloxera with fungus disease. < Amér. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.
 - Abstract of discussion at meeting of Académie des Sciences de Paris on the infection of *Phylloxera vastatrix* by parasitic fungi; particular fungi infect only particular insects; method of experimentation to determine which fungi to use and how to apply them; improbability of success in these experiments.
- 1809. [RILEY, C. V.] Fungus in Cicada. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.
 - Notice of C. H. Peck's description and J. Leidy's note on Massospora cicadina; Cicada [= Tibicen] septendecim and C. [= T.] tredecim subject to the attacks of the same.
- 1810. [RILEY, C. V.] On the nature of the phosphorescence of the glowworm. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.

 Conclusions drawn by Jousset de Bellesme; probability that the phosphor-
- escent substance is a gaseous product; nature of phosphorescence.

 1811. [RILEY, C. V.] Death of mules caused by insects. < Amer. Ent.,
 June, 1880 [v. 3], n. s., v. 1, p. 148.

About 6,000 mules reported killed in the Ouachita Valley, Louisiana, by the attacks of Simulium sp.

- 1812. [RILEY, C. V.] [May beetles swarming in Alabama.] < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 148.
 - Abundance and ravages of Lachnosterna queroina [=fusca] on oak-trees around Mobile, Ala., in May, 1880.
- 1813. [RILEY, C. V.] Fungus diseases of beneficial insects. < Amer. Eut., June, 1880 [v. 3], n. s., v. 1, p. 149.
 - Abstract of communication by C. Brongniart and Max Cornu on an epizootic among Syrphus mellinus caused by a species of Entomophthora; almost all insecticides liable to the objection that they kill useful as well as noxious insects.
- 1814. [RILEY, C. V.] Early appearance of cotton-worm. <Amer. Ent.,
 June, 1880 [v. 3], n. s., v. 1, p. 149.

 Extracts from letter of L. M. Rell, and from the Golied Free 1 Green and

Extracts from letter of J. M. Bell and from the Goliad [Tex.] Guard and from G. Witting on the date of appearance of Aletia argillacea [= xylina] in Texas.

- 1815. [RILEY, C. V.] The butterfly tongue. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 149.
 - Notice of and extract from paper by E. Burgess; conclusions as to the manner in which butterflies imbibe their liquid food.
- 1816. [RILEY, C. V.] Cottony maple scale. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 149.

Notice of J. D. Putnam on Pulvinaria innumerabilis.

- 1817. [RILEY, C. V.] Necrological. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 150.
 - Obituary notices of E. A. H. v. Kiesenwetter, S. C. Snellen van Vollenhoven, and F. L. de Laporte.
- 1818. [RILEY, C. V.] Effects of severe cold on insects. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 150.

A steady even if severe winter not prejudicial to insect life.

- 1819. [RILEY, C. V.] Revision of the *Lampyridæ*. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 150.
 - Notice of H. S. Gorham's work on the Lampyrida and his request for additional material.
- 1820. [RILEY, C. V.] Apple-twig borer. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 151.
 - Answer to inquiry of A. S. H.; occurrence of Amphicerus bicaudatus on appletrees in Virginia.
- 1821. [RILEY, C. V.] Grape-vine flea beetle. <.\text{Mer. Ent., June, 1880}
 [v. 3], n. s., v. 1, pp. 152-153.

 Auswer to letter of J. Nilis; ravages of Graptodera [= Haltica] chalybea on
- grape-vines in Pennsylvania.

 1899 (Proper C. W.) Carried cell on each trian.
- 1822. [RILEY, C. V.] Cynipid gall on oak twigs. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 153, fig. 56.
 - Answer to letter of J. A. Warder; figure of gall of Cynips (n. sp.†) from twig of Quercus prinus var. acuminata; comparison of this gall with that of C. [=Andrious] q.-punctata; the flies bred from these galls issue in spring and are all females, but probably have a bisexual form producing a different gall.
- 1823. [RILEY, C. V.] White-grub fungus. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 153.

 Answer to letter of H. S.; occurrence of Torrubia ravenelii at Iola, Kans.
- 1824. [RILEY, C. V.] Galerita janus. < Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 153, fig. 57.
 - Answer to inquiry of H. D. M. Fair; distribution, habitat, transformations, and seasons of Galerita janus; figure of larva of G. lecontei.
- 1825. [RILEY, C. V.] Not Fuller's rose-beetle. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, pp. 153-154, fig. 58.

 Answer to inquiry of J. Stewart; habits and synonymy of Agonoderus comma [=pallipes]; figure of the same.
- 1826. [RILEY, O. V.] Not Aletia chrysalides. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 154.
 - Answer to letter of G. P. White; pups of Agrotic saucia from cotton fields; food-habits of larva; ravages of Aletia argillacea [=xylina] at Brown Station, Ala., in August, 1879.
- 1827. [RILEY, C. V.] Mud-wasp and parasite. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 154, figs. 59-60.

 Answer to inquiries of Mary Treat: nidification of several species of Odyngrus.
 - Answer to inquiries of Mary Treat; nidification of several species of Odynerus; food-habits of O. birenimaculatus; figure of O. flavipes and of a nest of Odynerus sp.; figure of Cryptus juncus.
- 1828. [RILEY, C. V.] Monographs again. <Amer. Ent., June, 1880, [v. 3], n. s., v. 1, pp. 154-155.

 Answer to letter of C. D. Marsh; references to works. Elaterida, Curculio-

nide and Coccide.

- 1829. [RILEY, C. V.] Rearing wood-borers. <Amer. Ent., June, 1880 [v. 3], n. s., v. 1, p. 155.
 - Answer to letter of W. H. Harrington; directions for rearing larvæ which bore in wood or under bark.

- 1830. BILEY, C. V. On a new tineid genus allied to *Pronuba*, Biley. <Amer. Ent., June, 1880, [v. 3], n. s., v. 1, pp. 155-156.
 - Description of *Prodoxus* n. g. [p. 155] of *Tinoins* and of larva, pupa, and imagos of *P. decipiens* n. sp. [p. 155].
- 1831. RILEY, C. V. A parasite on *Prodoxus decipiens*. <Amer. Ent., June, 1880, [v. 3], n. s., v. 1, p. 156.

 Description of larva and image of *Exothecus prodoxi* n. sp.; habits of larva.
- 1832. RILEY, C. V. How to manage the cotton-worm: Suggestions to cotton planters. <Farmer's Review, 8 July, 1880. S.-b. No. 24, p. 68.</p>
 Means against Aletia xylina.
- 1833. [RILEY, C. V.] The cotton-worm investigation. <Selma [Ala.]

 Morning Times, 21 July, 1880, v. 55, No. 190, p. 3. Reprint:

 <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 197.

 List of persons engaged in the cotton-worm investigation; statement of work
- to be done by each.

 1834. [RILEY, C. V.] A foe to cottonwood. < Amer. Ent., July, 1880
- [v. 3], n. s., v. 1, pp. 159-161, figs. 61-64. Extract: <Suppl. to Amer. Ent., July, 1880, p. 1.

 Description of egg, larva, and image and figures of all stages of Melasoma
 - Description of egg, larva, and image and figures of all stages of Metasoma [=Lina] scripta; description and figures of variations of the image; habits, ravages, seasons, food-plants, and means against the same; figure of M. [=L.] lapponica and of larva of M. [=L.] populi; description of these larvæ and of that of M. [=L.] tremulæ; food-plants and distribution of these species; acquisition of new habits by insects.
- 1835. [RILEY, C. V.] The northern army-worm. Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 170-171, figs. 72-75.

 Ravages of Leucania unipuncta in 1880; natural history, means against, description, and figures of all stages of the same.
- 1836. [RILEY, C. V.] The periodical Cicada. Amer. Ent.. July, 1880
 [v. 3], n. s., v. 1, pp. 172-173, fig. 76.

 Broods of Cicada [= Tibicen] septendecim and C. [= T.] tredecim which appear in 1880.
- 1837. [RILEY, U. V.] Use of guano for grape Phylloxera. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 173. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 2.
 - Application of sulphide of carbon with infusorial earth or guano.
- 1838. [RILEY, C. V.] Fertilizers of alpine flowers. < Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 175. Reprint: < Suppl. to Amer. Ent., July, 1880, p. 2.
 - Relative frequency, according to H. Müller, of the visits of insects to flowers in high alpine regions as compared with such visits at lower levels.
- 1839. [RILEY, C. V.] Carnivorous habits of caddis-worms. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 176.

 Notice of paper by G. C. Goody.

- 1840. [RILEY, C. V.] Development of the eyes and luminosity in the fire-flies. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 176.

 Abstract of H. S. Gorham's observations.
- 1841. [RILEY, C. V.] Grape Phylloxera not at the Cape. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 176. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 1.
 Examination of roots of unhealthy vines by R. McLachlan and R. Trimen show that the disease of the vines is not caused by Phylloxera vastatrix.
- 1842. [RILEY, C. V.] [Bill providing for the extermination of insects.]

 Amer.Ent., July, 1880 [v. 3], n. s., v. 1, pp. 176–177.

 Text of and comments on a bill introduced into the California Assembly, providing for the extermination of insects.
- 1843. [RILEY, C. V.] [Appropriation for the U.S. Entomological Commission.] <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 177.

 Notice of the appropriation by Congress of \$25,000 for completing the work of the U.S. Entomological Commission.
- 1844. [RILEY, C. V.] Pronuba vs. Prodoxus. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 177-178.

 Comments on letter of V. T. Chambers.
- 1845. [RILEY, C. V.] Interesting cotton-worm notes from Vera Cruz, Mexico. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 179. Reprint with slight omission: <Suppl. to Amer. Ent., July, 1880, p. 3.
 - Letters from S. T. Trowbridge and R. de Zayas Enriquez, with notes on the irregular occurrence of Aletia argillacea [=xylina] in Vera Cruz.
- 1846. [RILEY, C. V.] Clover-root borer. <Amer. Ent., July, 1880
 [v. 3], n. s., v. 1, pp. 179-180, fig. 81. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 4. See: <Prairie Farmer, 31 July, 1880. S.-b. No. 45, p. 15.
 - Answer to letter of W. A. Henry; description and figures of larva, pupa, and imago of Hylesinus trifolii; ravages, habits, and means against the same.
- 1847. [RILEY, U. V.] Mud-wasp and spider egg-nest. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 180, figs. 82-84.

 Answer to letter of Mary Treat; figures of cells and of image of Eumenes fraterna, which stores its cells with larvæ of Paleacrita vernata; figure and descriptions of egg-nests of Epeira sp.
- 1848. [RILEY, C. V.] Worm in joints of wheat. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, pp. 180-181, fig. 85. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 4.
 - Answer to letter of J. K. P. Wallace; ravages of an unknown, apparently hymenopterous, larva in wheat stalks; figure of larva and pupa of *Meromyza americana* and of stalks injured by the same.
- 1849. [RILEY, C. V.] Linden and ash destroyers. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 181. Reprint: <Suppl. to Amer. Ent., July, 1880, p. 4.

- 1849. RILEY, C. V.—Continued.
 - Answer to letter of Shelby Reed; season of appearance of Odontota rubra on Tilia; larvæ of Hyphantria textor [= cunea] feeding on the same; Saperda calcarata boring in poplar, and an unknown larva (Neoclytus capræa?) boring in black-ash.
- 1850. [RILEY, C. V.] Larva boring along the axis of apple-twigs. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 181.

 Answer to letter of T. V. Munson; twigs of apple-trees bored by larvæ of

Answer to letter of T. V. Munson; twigs of apple-trees bored by larvæ of Oberea sp.?

- 1851. [RILEY, C. V.] Aquatic larvæ. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 181.
 - Answer to letter of S. A. Forbes; larve of Anax junius and Palingenia [= Hexagenia] bilineata named; the latter common in the stomach of fishes.
- 1852. [RILEY, C. V.] First appearance of cotton-worm in prairie belt.

 Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 181. Reprint:

 <Suppl. to Amer. Ent., July, 1880, pp. 3-4.
 - Answer to letter of J. F. Bailey; larvæ of Aletia argillacea [=xylina] appear first on lands where the cotton is luxuriant.
- 1853. [RILEY, C. V.] Gyrinus larva; terrestrial insects in stomach of shad. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 182.
 - Answer to letter of S. A. Forbes; character of larva of Gyrinus; a mass of terrestrial insects including Typhlocyba vitis? a muscid, a Eurytomid, Jassus sp., Triphleps insidiosus, and other species from the stomach of Ohio shad.
- 1854. RILEY, C. V. Further remarks on the differences between *Pronuba* and *Prodoxus*. <Amer. Ent., July, 1880 [v. 3], n. s., v. 1, p. 182.
 - Differences between Pronuba yuccasella and Prodoxus decipiens in the form, sculpture, and color of the terminal joint, and of the ovipositor.
- 1855. [RILEY, C. V.] The grape-vine flea-beetle, Graptodera chalybea Illig. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 183-184, fig. 86.
 - Habits, seasons, ravages of, and means against Grapiodera [= Haltica] chalybea; figures, injured leaf, and larvæ, cocoon, and image of the same.
- 1856. RILEY, C. V. Further notes and observations on the armyworm. <Amer. Ent., 1880 [v. 3], n. s., v. 1: August, pp. 184, 185; September, pp. 214, 215. Reprint, with changes: <Sci. Amer., 4 September [v. 57], n. s., v. 43, p. 152. S.-b. No. 23, pp. 161-162.
 - Progress of knowledge upon the number of annual generations of Leucania unipunots; stages of growth at which it hibernates; the destructive generation probably not the first of the season; explanation of the partial efficacy of the burning of fields in winter as a means against this insect; connection of wet and dry seasons with its increase; its natural habits; errors of A. Fitch.
- 1857. [RILEY, C. V.] Sprinklers and atomizers. <Amer. Ent., 1880 [v. 3], n. s., v. 1: August, pp. 185-189, figs. 87-98; September, pp. 211-214, figs. 111-117.
 - Extracts from pp. 56-57 and reprint of pp. 85-94 of Bull. No. 3, U.S. Entomological Commission. See No. 1736 for synopsis of contents.

- 1858. RILEY, C. V. The use of pyrethrum. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 193-195.
 - Record of experiments by W.A. Henry upon the effect of Pyrethrum powder on Halticidæ, Meloidæ, Pieris rapæ, Coreus [= Anasa] tristis, and Blattidæ; and of the fumes of burning pyrethrum on various insects.
- 1859. [RILEY, C. V.] [Colorado potato-beetle in New Hampshire.] <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 195.
 Ravages of Doryphera 10-lineata in Coos County, N. H., in 1879.
- 1860. [RILEY, C.V.] Retarded development in a blister-beetle. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 196. Final transformations of individuals from a single batch of eggs of *Epicauta vittata* occurring at the first, second, and third year after hatching.
- 1861. [RILEY, C. V.] Ox-eye daisy as an insecticide. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 196.

 Experiments by W. S. Barnard upon the effect of alcoholic extracts of the flowers and stems of the ox-eye daisy on insects; "no evidence that they will prove of any practical value."
- 1862. [RILEY, C.V.] Directions for raising pyrethrum. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 197.

 Directions for sowing the seeds and cultivating the plants of pyrethrum.
- 1863. [RILEY, C. V.] State Entomologist for New York. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 197, 198.

 Notice of the appointment of J. A. Lintner as State Entomologist of New York.
- 1864. [RILEY, C. V.] Economic investigations in the South and West. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 198.

 Meeting of the U. S. Entomological Commission; plan of work to be done in 1880-1881; partial list of persons engaged in the work.
- 1865. [RILEY, C. V.] [Number of entomologists in Europe.] <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 198.

 Tabulation of the number of entomologists of the several countries of Europe.
- 1866. [RILEY, C. V.] [Catalogus coleopterorum by Gemminger and Harold.] <Amer. Ent., August, 1880 [v. 3], ii. s., v. 1, p. 199. Commendation of the above work and of the supplementary list of *Elaterida* by E. Candèze; need of co-operation for the completion of the whole catalogue.
- 1867. [RILEY, C.V.] Carnivorous propensity of plant-feeders. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 200.

 Larvæ of Plusia brassicæ devour those of Pieris rapæ and Pionea rimosalis in default of more natural food.
- 1868. [RILEY, C. V.] Beetles injuring cabbages and fuchsias. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 200, fig. 106.

 Answer to letter of G. T.; ravages, distribution, and figures of Epicarus imbricatus; description of eggs, ravages, food-plants, and means against Graptodera [= Haltica] carinata.
- 1869. [RILEY, C. V.] Spider and nest. < Amer. Ent., August, 1880, [v. 3], n. s., v. 1, p. 200.

Answer to letter of Mrs. J. B. Harrison; distribution and distinctive characters of Acrosoma stellatum.

- 1870. [RILEY, C. V.] Damage to wheat: Worm boring in the stalk. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 201.
 - Answer to letter of A. R. Frost; food-plants and ravages of larva and figure of larva and imago of Gortyna nitela.
- 1871. [RILEY, C. V.] Larvæ from stomach of blue-bird. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 201.
 - Answer to letter of S. A. Forbes; larva of Meracantha contracta and of Callimorpha sp. from stomach of blue-bird; habits and characters of the same.
- 1872. [RILEY, C. V.] Butterfly larva injurious to cotton squares. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 201.

 Answer to letter of B. F. Cooke; larva of Theela (pages) feeding on leaves
- and bolls of cotton-plant; a species of *Microgaster* parasitic in the larva.

 1873. [RILEY, C. V.] Cut-worms from stomach of robin. < Amer. Ent.,
- August, 1880 [v. 3], n. s., v. 1, p. 201.

 Answer to letter of S. A. Forbes; larvæ of Agrotis (messoria?) from stomach of robin; Agrotis cochrani = A. messoria.
- 1874. [RILEY, C. V.] Large phosphorescent larva. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 201-202, fig. 108.

 Answer to letter of S. F. Clarke; occurrence in Maryland and Missouri of a luminous larva (Melanacies sp. ?) [= Phengodes sp.]; figures of the larva, of its head and leg enlarged and of its probable parent; occurrence in the more northern States of a similar larva, probably that of Asaphes memnonius.
- 1875. RILEY, C. V. Worms injuring wheat. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 202.

 Answer to letter of I. Monoghan: rayages of large of Mercange americans.
 - Answer to letter of J. Monaghan; ravages of larva of Meromyza americana.
- 1876. RILEY, C.V. Ash-root borer: Supposed eggs of Odontota. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 202-203.

 Answer to letter of Shelby Reed; habits and ravages of Parandra brunnes; Pimpla sp. parasitic upon its larva; probable manner in which the eggs of Odontota rubra are laid.
- 1877. [RILEY, C. V.] Army-worm notes and inquiries: Its work on clover. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 203.

 Answer to letter of L. T. Derousse; Leucania unipuncta frequently leaves clover-plants untouched while eating the grasses growing with the clover, but under some circumstances destroys young clover-plants.
- 1878. [RILEY, C. V.] Ichneumon from stomach of bluebird. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 203.

 Answer to letter of S. A. Forbes; Lampronota sp. eaten in quantity by bluebirds.
- 1879. [RILEY, C. V.] Leaf miner on white oak. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 203.

 Answer to letter of E. W. Claypole; description, habits, and ravages of Lithecolletis cincinnaticlls.
- 1880. [RILEY, C. V.] Screw-worm: Its parentage in doubt. < Amer. Eut., August, 1880 [v. 3], n. s., v. 1, p. 203.

 Answer to letter of A. R. Kilpatrick; Lucilia macellaria supposed to be the parent of the "screw-worm"; food-habits of certain Muscides.

- 1881. [RILEY, C. V.] Silk culture: How to dispose of cocoons. < Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 203-204.
 - Answer to letter of C. F. Durksen; need of silk-filatures in the United States; present means of selling cocoons.
- 1882. [RILEY, C. V.] Best cotton-worm destroyer. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 204.
 - Answer to letter of J. G. Dauterive; reference to the most reliable poison and the best machines and methods for applying poison to kill Aletia argillacea [= xylina].
- 1883. [RILEY, C. V.] Twice-stabbed lady-bird. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 204, fig. 109.
 - Answer to letter of H. N. Patterson; Coccinellida beneficial by feeding on Aphidida and Coccida; figure of Chilocorus bivulnerus; significance of the presence of this beetle in large numbers on trees.
- 1884. [RILEY, C. V.] Pseudo-scorpion. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, p. 204.
 - Answer to inquiry of C. H. S. Davis; habitats and food-habits of Chelifer camoroides and other Chernetidæ.
- 1885. [RILEY, C. V.] Bluebirds feeding on parasitic and predaceous insects. <Amer. Ent., August, 1880 [v. 3], n. s., v. 1, pp. 204–205.
 - Answer to letter of S. A. Forbes; bluebirds more destructive to parasitic and predaceous insects than the thrushes; percentage of such food found in the stomachs of the bluebird; larvæ of Leucania unipuncta, Telephorus bilineatus, and Nephelodes violans eaten by the same; probable hibernation of the first named; habits, hibernation, distribution, and popular name of the Nephelodes.
- 1886. RILEY, C. V. The cotton destroyers. <New Orleans Democrat, 21
 September, 1880, v. 5, No. 276, p. 8. S.-b. No. 23, pp. 170-173.
 Reprint: <Selma [Ala.] Times, 29 September, 1880. <Southern Enterprise [Atlanta, Ga.], December, 1880, v. 5, pp. 77-82.
 S.-b. No. 23, pp. 184-189; No. 61, pp. 3-5; No. 63, pp. 57-59;
 60-62. Reprint, with slight changes: <Proc. Amer. Assoc.
 Adv. Sci. for 1880, 14 October, 1881, v. 29, pp. 642-649. Separate: <Salem, July, 1881, pp. 26-33. Extract: <Sci. Amer., 16
 October, 1880 [v. 57], n. s., v. 43, p. 241. S.-b. No. 23, pp. 159160. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, pp. 245-247.
 <Prairie Farmer, 30 October, 1880, v. 51, No. 44, p. 2. S.-b. No.
 23, pp. 153. Abstract: <Farmer's Review, 7 October, 1880. S.-b.
 No. 23, pp. 162-163. <Journ. Appl. Sci., November, 1880, v.
 11, pp. 170-171.
 - Facts and principles established by the U. S. Entomological Commission, applicable to the whole cotton belt, regarding the times and manner of first appearance of Aletia argillacea [=xylina] and Heliothis armigera upon the cotton-plant; habits of the larvæ and images and the best means of destroying the same; relative efficacy of several insecticides; methods of preparing and applying the same; importance of early poisoning; main object of the cotton-worm inquiry accomplished.

- 1887. [RILEY, O. V.] The use of poisons to destroy insects. **Farmer's**Review, 23 September, 1880, v. 5, p. 200. S.-b. No. 23, pp. 148-149; 165-166. Reprint: **Amer. Ent.**, October, 1880 [v. 3], n. s., v. 1, p. 244.
 - Advocacy of the use of Paris green and London purple as means against certain insects; danger of the use of the same against many insects; criticism of A. J. Cook's recommendation for their use against Paris aterrima and Carpocapea pemonella; limitations within which they may be used.
- 1888. [RILEY, C. V.] Supplementary instructions to agents of the United States Entomological Commission. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 218.
 - Call for experiments to ascertain the minimum quantity of poisons which can be used effectually against Aletia argillacea [=xykisa]; effect of poisons on the several stages of this insect; food-plants, parasites, and enemies of the same; effect of yeast ferment upon it.
- 1889. RILEY, C. V. Dimorphism in locusts (Acrididæ). < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, pp. 219-220.
 - Review of paper by S. H. Scudder; certain forms described as species of Pesotettix are dimorphic forms of Caloptenus; extract from 8th Ann. Rept. State Ent. Mo., 1876, p. 115.
- 1890. [RILEY, C. V.] A scale insect on maple hitherto unobserved by American entomologists. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, pp. 220-221.
 - Notice of paper by Miss E. A. Smith, with emended reprint of a portion of the same; habits, colors, and behavior of young larvæ and transformations of the male of *Pseudococcus aceris*.
- 1891. [RILEY, C. V.] How flight in insects is directed. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 221.
 - Abstract of paper by Jousset de Bellesme; direction of flight not determined by the motion of the wings, but principally by the displacement of the center of gravity, resulting from the changes of position of other parts of the body.
- 1392. [RILEY, C. V.] Entomological work at the Department of Agriculture. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 222.

 Notice of the continuation of the appropriations by Congress for field-work and experiments in the entomological division of the U. S. Department of Agriculture.
- 1893. [RILEY, C. V.] Entomological papers read before the A. A. A. S. Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 223. Titles of twenty-eight [28] papers read at the Boston meeting.
- 1894. [RILEY, C. V.] Entomologists at Boston. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 223.
 - Notice of the 29th meeting of the A. A. S. at Boston, August-September, 1880.
- 1895. RILEY, C. V. Winged Phylloxera in California. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, pp. 224-225.
 - Letter from J. S. Hyde, with remarks; extracts from papers by E. W. Hilgard; winged fertile females of *Phylloxera vastatrix* found in California; local evidence of the spread of this insect; means of eradicating the same.

- 1896. [RILEY, C. V.] Worm infesting meal sacks. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 229.
 - Answer to letter of J. Greenwood, jr.; description of larva and image of Ephestia zew [=interpunctella]; food-habits of the larva.
- 1897. [RILEY, C. V.] Hesperid larva feeding on Canna. < Amer. Eut., September, 1880 [v. 3], n. s., v. 1, p. 229.
 - Answer to letter of N. S. Reed; occurrence of Pamphila ethlius in Illinois and South Carolina; its larva injurious to Canna flacoida.
- 1898. [RILEY, C. V.] Grape-vine apple-gall. < Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 229, fig. 119.
 - Answer to letter of A. R. McCutchen; occurrence of Cecidomyia ? vilis-pomum and of Calosoma scrutator in Georgia; figures gall of the first named; food-habits of the Calosoma.
- 1899. [RILEY, C. V.] Apple-tree plant-lice in Oregon. <Amer. Ent., September, 1880 [v. 3], n. s., v. 1, pp. 229-230.

 Answer to letter of H. B. May: occurrence and rayages of Ankis mali in
 - Answer to letter of H. B. May; occurrence and ravages of Aphie malit in Oregon; means against the same.
- 1900. [RILEY, C. V.] Phylloxera work. Wood-lice on grape-vine roots. Amer. Ent., September, 1880 [v. 3], n. s., v. 1, p. 230.
 - Answer to letter of H. B. Trimble; occurrence of *Phylloxera vastatrix* at West Chester, Pa.; roots of grape-vines destroyed from unknown cause, perhaps by young of *Phylosoia* or *Porcellio*.
- 1901. RILEY, C. V. New hickory galls made by Phylloxera. <Amer. Ent., September, 18:0 [v. 3], n. s., v. 1, p. 230.
 - Description of Phylloxera carya-scissan sp. and P. carya-avellana n. sp. from Florida; larva of Diplosis found in the galls.
- 1902. RILEY, C. V. Food-habits of the longicorn beetles or wood-borers. <Amer. Ent., 1880 [v. 3], n. s., v. 1; October, pp. 237-239; November, pp. 270-271.
 - Food-plants and food-habits of the Prionida and Cerambycida of the United States.
- 1903. [RILEY, C. V.] Additional experiments with Pyrethrum. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 242.
 - Details of experiments made by H. G. Hubbard on the effect of very minute quantities of Pyrethrum powder on very young larvæ of Aleka argillacea [=xylina].
- 1904. [RILEY, C. V.] A new enemy to the strawberry. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, pp. 242-243, fig. 121.
 - Review of paper by A. J. Cook; extracts from the same; description and habits of larva and imago of *Paria aterrima*; means against the imago; habits of larva of *Colaspis flavida*; figure of the same; number of segments and spiracles in larvæ of Coleoptera.
- 1905. [RILEY, C. V.] A new enemy to corn: The long-horned Diabrotica. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 247.
 - Notice of recent articles on the ravages of the larva of Diabrotica longicornis; food-habits and means against the same; food-habits of Drasterius amabilis.
- 1906. [RILEY, C. V.] Phylloxera congress in Spain. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 247.
 - Notice of a congress to be held in Spain to consider all topics connected with the ravages of *Phylloxera vastatrix*.

- 1907. [RILEY, C. V.] The grape Phylloxera not permanently destructive. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 248.
 - Recovery of grape-vines formerly injured by *Phylloxera vastatrix* in California and Missouri; belief that in Europe grape-vines will be grown again on the lands of late years ravaged.
- 1908. [RILEY, C. V.] Sale of silk-worm eggs. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 248.
 - \$6,000,000 worth of silk-worm eggs sent from Japan to France via San Francisco in each of the four years 1874-1877.
- 1909. [RILEY, C. V.] [Death of S. S. Haldeman.] < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 248.

 Obituary notice.
- 1910. [RILEY, C. V.] [Retirement of Mr. Fuller.] < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 248.
 - Announcement of the retirement of A. S. Fuller as assistant editor of the American Entomologist.
- 1911. [RILEY, C. V.] Insect enemies of growing rice. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 253.
 - Answer to letters of J. L. Leconte and J. Screven; ravages of and means against the "grub" [= Chalepus trackypygus] and the "maggot" [= Lissorhoptrus simplex] in rice fields in Georgia.
- 1912. [RILEY, U. V.] Blind-eyed Smerinthus. < Amer. Ent., October 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to inquiry of Mrs. J. B. Harrison; occurrence at lamp-light in N. H. of Smerinthus excecatus; food-plant of the larva of the same.
- 1913. [RILEY, C. V.] White waxy secretion on stems of bitter-sweet. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to inquiry of J. A. Lintner; method of oviposition and characters of egg-mass of *Enchophyllum* [= *Enchonopa*] binotata; seasons, habits, and food-plants and description of the eggs of the same.
- 1914. [RILEY, C. V.] Prickly-ash larva: Tachinid eggs. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to inquiry of R. W. Jones; larva of Papilio cresphontes on prickly ash; eggs of Tachina sp. on larva of Aletia argillacea [=xylina].
- 1915. [RILEY, C. V.] Worms on cabbage: Boll-worm feeding on leaf. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to inquiry of R. W. Jones; food-habits of larve of Heliothis armigers and of Piones rimosalis.
- 1916. [RILEY, C. V.] Buggy peas. <Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254.
 - Answer to letter of A. Berger; abundance of Bruchus pisi in Wisconsin; means against it.
- 1917. RILEY, C. V. Glow-worm. < Amer. Ent., October, 1880 [v. 3], n. s., v. 1, p. 254, fig. 123.
 - Answer to letter of J. J. Dean; occurrence of larva of *Photuris pennsylvanics* at Chatham, N. J.; figures larva and imago of the same; larva and imago of *Photinus pyralis*, also luminous; males and females of these species winged; female of *Lampyris noctiluca* wingless and more luminous than the male.

- 1918. [RILEY, C. V.] The use of fungus growths to destroy insects. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, pp. 269-270.
 - Abstract of paper by A. N. Prentiss; the constant presence of spores of fungi in the air vitiates the value of most experiments on the application of the fungi to insects; insects rarely affected by the spores which are in the air while the plants are more so affected; yeast more injurious mechanically than infectiously.
- 1919. [RILEY, C. V.] New species of scale insects. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, pp. 275-276.

 Review of paper by W. H. Ashmead on the red scale.
- 1920. [RILEY, C. V.] Remedy for cabbage worms. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 276.</p>
 Pyrethrum powder the most satisfactory means against all larvæ affecting the leaves of cabbage.
- 1921. [RILEY, C. V.] Pyrethrum for the screw-worm. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 276.

 Notice of paper by A. R. Kirkpatrick; directions for the use of pyrethrum powder against the "screw-worm" [= Lucilia macellaria]; objections to the use of other remedies.
- 1922. [RILEY, C. V.] Oviposition in the *Tortricidæ*. <Amer. Ent. November, 1880 [v. 3], n. s., v. 1, p. 276.

 Notice of and extract from paper by C. H. Fernald.
- 1923. [RILEY, C. V.] About *Phora* being merely a scavenger and not a true parasite. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 277.
 - Letter from C. R. Osten Sacken, with remarks; Phora aletiæ not a parasite; oviposition and habits of larva of the same.
- . 1924. [RILEY, C. V.] Gall on Solidago leaves. <Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 278.

 Answer to letter of H. Barnes; occurrence of galls of Cecidomyia carbonifera on leaves of Solidago nemoralis? at Mulberry Corners, Ohio.
 - 1925. [RILEY, C. V.] Oak gall: Cynips q.-decidua Bass. < Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 278.

 Answer to letter of J. Schenck; galls of Cynips q. decidua? found on leaves of Querous muhlenbergii at Mount Carmel, Ill.
 - 1926. [RILEY, C. V.] Insects from stomach of lark, robin, and sunfish.

 Amer.Ent., November, 1880 [v. 3], n. s., v. 1, p. 278.

 Answer to letter of S. A. Forbes: Diplotarie sordida from stomach of meadow-

Answer to letter of S. A. Forbes; Diplotaxis sordida from stomach of meadow-lark; egg of a reduviid from that of a robin; larva of an ephemerid (Polymitarcys alba*) from the stomach of a sunfish.

- 1927. [RILEY, C. V.] Supposed hibernating Aletia chrysalis. < Amer. Ent., November, 1880 [v. 3], n. s., v. 1, p. 278.

 Answer to letter of J. W. Davidson; pupa resembling that of Achatodes zeas found in stalk of maize at Uniontown, Ala.
- 1928. RILEY, C. V. On the natural history of certain bee-flies (Bomby-liidæ). <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, pp.

- 1928. RILEY, C. V.—Continued.
 - 279-283, figs. 147-151. Review: <Ent. Mo. Mag., February, 1881, v. 17, pp. 206-207.
 - Advance print of pp. 262-267 and a general abstract of pp. 267-269 of the section entitled "Bee-fly larvs, family Bombylidae," in 2d Rept. U. S. Entomological Commission. See No. 1969 for synopsis of contents.
- 1929. RILEY, C. V. On a new pyralid infesting the seed-pods of the trumpet-vine. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, pp. 286-288, figs. 152-158.
 - Description, with figures, of Clydonopteron n. g. [p. 287] and of larva, pupa, and image of C. tecomes n. sp. [p. 288], and of the abode of this insect in pods of the trumpet-vine; habits of the insect.
- 1930. [RILEY, C. V.] Experiments with yeast-ferment on various insects. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, pp. 289, 290.
 - Reports by J. E. Willet and A. J. Cook upon their experiments in the application of beer and yeast to various insects; no infection communicated to the insects by these applications.
- 1931. RILEY, C. V. Notes on the imported elm leaf-beetle. < Amer. Ent., December, 1880 [v. 3], n. s., v. 1, pp. 291-292.
 - Answer to letter of J. L. Leconte; seasons, transformations, enemies, and ravages of and means against Galeruca zamthomelæma; hibernation of Chrysomelidæ.
- 1932. [BILEY, C. V.] Synonyms of parasites' mistakes corrected. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 293. Review: <Ca. Ent., February, 1881, v. 13, pp. 31-33, fig. 3.
 - Review of paper by L. O. Heward; comparison of the alleged characters of Antigaster and Eupelmus to show grounds for the founding of the former; Didictyum synonymous with Hexaplasta; H. sigzag not a parasite of Aletis argillacea [= xylina], but of Phora aletis; habits of H. zigzag.
- 1934. [RILEY, C. V.] Mandible of Lithocolletis guttifinitella. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 294, fig. 138. Figure, with explanation, of the mandible of Lithocolletis guttifinitella.
- 1935. [RILEY, C. V.] Excessive injury by a beetle in Russia. < Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 294.</p>
 Ravages of Anisoplia austriaca in southern Russia; description of the same; its habits and vernacular name.
- 1937. [RILEY, C. V.] Fungus foes. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p.297.
 - Answer to inquiry of J. J. Brown; Cie fuscipes and all Cieids infest fungi growing on old trees and logs.

- 1938. [RILEY, C. V.] The twig-girdler. <Amer. Ent., December, 1889 [v 3], n. s., v. 1, p. 297, figs. 155-156.
 - Answer to letter of W. R. Maxwell; figures of larva, pupa, and image of Oncideres cingulata and of twig injured by the image; food-plants, habits, and transformations of this insect.
- 1939. [RILEY, C. V.] The bedeguar of the rose. < Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 298, fig. 154.

Answer to letter of L. C. Bryan; account of the gall of Bhodites resæ; figure of the same.

- 1940. [RILEY, C. V.] Minute borers in cherry, peach, and plum-trees. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 298.
 - Answer to letters of J. L. Bennett and M. H. Boye; ravages, food-plants, and distribution of Scolytus rugulosus; food-habits of the species of Scolytus.
- 1941. [RILEY, C. V.] Smilax injured by cut-worms. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 298.

Answer to letter of G. Thommen; habits, ravages, and food-plants of Agretis saucia.

- 1942. [RILEY, C. V.] Honey-producing oak-gall. <Amer. Ent., December, 1880 [v. 3], n. s., v. 1, p. 298.
 - Answer to letter of H. C. McCook; secretion of saccharine matter by galls growing on Quercus undulata; description of these galls, Cynips quercus-mellaria n. sp.; Myrmeocoystus horius-deorum [= melliger] collects the saccharine matter from these galls.
- 1943. [RILEY, C. V.] Dr. Hagen's mystery. <Ca. Ent., December, 1880, v. 12, pp. 263-264.
 - Review of paper by H. A. Hagen; Prodoxus decipiens confounded with Pronuba guecasella.
- 1944. RILEY, C. V. Acorn-gall. <Trans. Acad. Sci. St. Louis, 1880, v. 4, p. 1 Proc.
 - Gall mistaken for abortive acorn, by G. B. Emerson, in his "Trees and shrubs of Massachusetts."
- 1945. RILEY, C. V. Silk-culture in the United States. Condensed account of the silk-worm and how to inaugurate a new source of wealth. <Western Farmer's Almanac for 1881, 1880, pp. 35-39, 4 figs. S.-b. No. 23, p. —.
 - Practicability and desirability of the culture of *Seriosria mori* in the United States; requisites to its success; superiority of *S. mori* over all other insects for silk-culture; brief illustrated natural history of the same; directions for silk-culture.
- 1946. RILEY, C. V. Legislation to control insects injurious to vegetation. <Farmer's Review, 20 January, 1881. S. b. No. 23, p. 148. Reprint: <Amer. Nat., April [25 warch], 1881, v. 15, pp. 322–323. <Indiana Farmer, 16 April, 1881. S. b. No. 23, p. 149. Notice: <Farmer's Review, 20 January, 1881. S. b. No. 23, p. 170.
 - Appointment of C. H. Dwinelle, by the California State Horticultural Society, on a committee to consider what legislation is desirable to check the spread of noxious insects and to force land-owners to destroy the same; notice of the previous passage of such laws and of anticipated objections to them.

- 1947. RILEY, C. V. Larval habits of bee-flies, Bombyliidæ. <Amer. Nat., February [25 January], 1881, v. 15, pp. 143-145, figs. 1-3. Notice: <Western Stock Journ. and Farm, March, 1881, v. 11, p. 58.
 - Habits of Systechus and Triodites as parasites on eggs of Acridide; notice of paper by T. A. Chapman; figures larva, pupa, and image of Systechus oreas and compares its larva and pupa with those of Bombylius major.
- 1948. RILEY, C. V. Experiments with pyrethrum: Safe remedies for cabbage-worms and potato-beetles. <Amer. Nat., February [25 January], 1881, v. 15, pp. 145-147.
 - Details of experiments made by A. J. Cook and W. R. Hubbert upon the effect of a dusting of pyrethrum powder upon larvæ of *Pieris rapæ*, larvæ and imagos of *Doryphora* 10-lineata, and upon *Eriosoma* [= *Pemphigus*] tessellata, Coreus [= Anasa] tristis, and flies and mosquitoes.
- 1949. RILEY, C. V. Insect enemies of the rice plant. < Amer. Nat., February [25 January], 1881, v. 15, pp. 148-149.
 - Chalepus trackypygus feeds on roots of the rice-plant; conjectures as to other enemies of the rice-plant; Cecidomyia oryzæ injurious to the same in India.
- 1950. [RILEY, C. V.] The "yellow-fever fly." < Amer. Nat., February [25 January], 1881, v. 15, p. 150.
 - Review of paper by H. A. Hagen; food-habits of larvæ of Sciara sp.; occurrence of swarms of images of the same.
- 1951. [RILEY, C. V.] An aquatic Sphinx larva. <Amer. Nat., February [25 January], 1881, v. 15, p. 151.

 Abstract of paper by H. A. Hagen.
- 1952. RILEY, C. V. Notes on the grape Phylloxera and on laws to prevent its introduction. <Amer. Nat., March [24 February], . 1881, v. 15, pp. 238-241. Notice with extracts: <Pacific Rural Press, 23 April, 1881. S.-b. No. 42, p. 17.
 - Remarks upon letter of I. Bush; precautions to be adopted against the introduction of *Phylloxera vastatrix*; summary of the life-history of the same.
- 1953. RILEY, C. V. Hibernation of the cotton-worm moth: Ease with which mistakes are made. <Amer. Nat., March [24 February], 1881, v. 15, pp. 244-245, figs. 1-3.
 - Extract from letter of I. A. Wimbish; Leucania unipuncta mistaken for Aletia argillacea [=xylina]; characters of Aletia; figures ovipositor and eggs of L. unipuncta and images of both species.
- 1954. [RILEY, C. V.] On some interaction of organisms. <Amer. Nat., April [25 March], 1881, v. 15, pp. 323-324.
 - Review of paper by S. A. Forbes, with extracts; relative unimportance of special parasites as compared with predaceous animals of varied tastes in the limitation of the numbers of any species of animals; need of conservative action and exhaustive inquiry in the attempt to interfere with the order of nature.
- 1955. [RILEY, C. V.] Insect locomotion. < Amer. Nat., April [25 March], 1881, v. 15, p. 325.

١.

Results of G. Carlet's studies on the order in which the feet are moved in the walking of Hexapoda and Arachnida. 1956. [RILEY, C: V.] Plant-feeding habits of predaceous beetles. <Amer. Nat., April [25 March], 1881, v. 15, pp. 325-327.</p>

> Citation of evidence from numerous sources proving that certain Carabida and Coccinellide occasionally feed on plants, seeds, and spores.

1957. RILEY, C. V. Notes on Papilio philenor. < Amer. Nat., April, [25 March], 1881, v. 15, pp. 327-329, figs. 1-3.

Description of egg and newly-hatched larva of Papilio philonor; figure of larva, chrysalids, and image of the same; food-plants, distribution, and its occurrence in swarms.

1958. [RILEY, C. V.] Entomological notes. < Amer. Nat., April [25] March], 1881, v. 15, pp. 330-331.

Notice of H. A. Hagen's paper on Simulium pictipes, with additional notes; abstracts and minor notices of other papers and items of news.

1959. [RILEY, C. V.] Second report of the United States Entomological Commission for the years 1878 and 1879, relating to the Rocky Mountain locust and the Western cricket, and treating of the best means of subduing the locust in its permanent breeding grounds, with a view of preventing its migrations into the more fertile portions of the trans-Mississippi country in pursuance of appropriations made by Congress for this purpose, with maps and illustrations. < Washington: 1880 [4 April, 1881], pp. 18+322+80, 10 figs., 17 pl., 9 maps.

TABLE OF CONTENTS.

LETTER OF TRANSMITTAL PREFACE	IX VIII
CHAPTER I: By A. S. Packard and C. V. Riley.	
Additions to the chronology of locust ravages	1
Utah, 12-13; in Montana, 13-14.	
CHAPTER II: By C. Thomas.	
The relation of the locust and its ravages to agriculture and the settlement of the Territories	14

CHAPTER III: By C. Thomas.

Facts concerning and laws governing the migrations of locusts in all countries

Most species of Acridida not migratory, 31-Migratory disposition not caused by anatomical differences, 31-32-Writers on locust flights, 32-33—Earliest accounts of locust flights in Europe, 33-34—Locust flights in Europe in the middle ages, 34-37; in the seventeenth century, 37-38-The locust invasion of 1693 in Europe, 38-39-Locust flights in Europe in the eighteenth and nineteenth centuries, 40-41-Locusts in Asia and adjacent islands, 41; in Cyprus, 42; in Arabia and Syria, 43-44; in Persia, 44; in the Steppes, 45; in China, 46; in Manila, 46; in India, 46-47; in Australia, 47; in New Zealand, 47-Locust flights in Africa, 48; in northern Africa, 48-49; in Egypt, 49in Algiers, 50-51; in Abyssinia, 51; in central Africa, 51; in eastern Africa and the Canary Islands, 52-53-Locusts in South America, 53-54-Prevalence of locusts in deserts and dry regions, 54-55-Locust flights not governed by laws of periodicity, 55-56-Permanent breeding grounds of locusts, 56-57-Their geographical distribution, 58-Caloptenus spretus not in Mexico, 58-Migratory locusts in Yucatan and the West Indies, 58-59-In Nicaragua, 59-Geographical limits of Caloptenus spretus, 59; of Pachytylus migratorius, 59-60-Difficulty of determining species, 60-61-Definition of permanent home of migratory locusts, 62-Range of Pachytylus migratorius, 62-63; of Acridium peregrinum, 63-64-Comparison with the laws of geographical zoology, 65-66-Meeting point of three sub-regions, 67-Each species has its permanent breeding ground, 68-General character of the permanent breeding grounds, 68-69-Breeding ground of Calentonus italicus, 69; of Pachytylus migratorius, 69-70; of Acridium peregrinum, 70; of Caloptenus spretus, 70-71.

CHAPTER IV: By C. Thomas.

Habits and characteristics of locusts in all countries within their areas of permanent distribution, so far as these relate to their movements

Caloptenus epretus always migratory, 72-73-Flights of C. epretus, 73-74—Invading swarms come from northwest, 74-75—Other opinions eliminated, 75-76—Swarms in 1875 and 1876, 76-77—Direction of returning swarms of C. spretus, 78; of other migratory locusts, 78-79-Direction of local flights, 79-81-Returning swarms do little injury, 81-82-Distance to which swarms may migrate, 82-84—Length of a single flight, 84-85—Swarms able to cross large bodies of water, 85-86-Swarms carried far by winds, 86-87-The mode of flight, 87-Position of the locust while flying, 87-88-Locusts flying with the wind, 88-89-Method of falling of swarms, 89-Formation of swarms, 89-91-Movements of swarms of C. spretus, as reported by Dr. Child, 91-93-Other reports, 94-Movements of swarms in other countries, 95-96-Swarms usually alight at night, 96-97-The height at which swarms move, 98-100-Different directions of swarms at the same time and place, 100-Fall of locusts into the sea, 100-102-Great flying power of locusts, 102-General causes of migration, 103-Excessive numbers, 103-104-Want of food, 104-Excessive heat,

31

·	
1969. [RILEY, C. V.]—Continued.	
CHAPTER IV—Continued.	
105-Dry condition of the atmosphere, 105-106-Arid condition	
of the mountain region, 196-107—Immediate causes of migration,	
107–108.	
CHAPTER V: By C. Thamas.	
Influence of meteerological conditions on the development and mi-	
grations of locusts	109
Influence of heat and dryness, 109-110-Packard's table of locust	
years, 111–112—Table of rain-falls for 1860 to 1866, 112—Influence	
of the rainfall, especially in 1864, 112-113—Average monthly tem-	
perature for 1864 and 1866, 113-115—Rain-fall of the years 1863 to	
1867, 116—Temperature of the years 1863 to 1867, 117—Extreme	
variations of monthly mean temperature from 1863 to 1867,	
118-119-Monthly and annual mean temperature from 1872 to	
1876, 119-121—Monthly and annual rain-full from 1872 to 1878, 122-	
124—Extreme fluctuations between monthly mean temperature,	
125-126—Greatest variations between monthly mean tempera-	
tures, 126-Variations of temperature between months of con-	
secutive years, 127-128—Daily temperature at Fort Sully from	
1872 to 1875, 129-134—Relation of temperature to the develop-	
ment of the eggs, 135-136—Professor Riley's experiments, 137—	
Professor Abbe's theory on the subject discussed, 137-140—Table	
• showing temperature and direction and velocity of wind at	
Western stations, 141-154—Relation of temperature and winds to locust flights, 155.	
CHAPTER VI: By A. S. Packard.	
The senthern limits of the distribution of the Recky Mountain locust.	156
Few locusts permanently breeding in New Mexico and none in Ari-	100
zona, 156—Caloptenus spretus probably not in southern New Mex-	
ico, central and southern Arizona, etc., 156—History of locust	
invasions in New Mexico from 1864 to 1879, 157-159—Southern	
range of the lecust in Mew Mexico, 159—Changes in the map in	
the Report I, showing the distribution of the locust, 159-160.	
CHAPTER VII: By A. S. Packard.	
Summary of locust flights from 1877 to 1879	160
General remarks on maps Nos. 2 to 4, 160-161-Flights of locusts	
in 1878, 161-162-Flights in 1879, 162-163.	
CHAPTER VIII: By A. S. Packard.	
The western cricket	163
Difference in habits between the western cricket and the locust,	
163-Occurrence of Anabrus purpurascens, 163-164; of A. simplex,	
164—Mode of egg-laying, 164—Movements of an army of the west-	
ern cricket, 164-165—Its ravages, 165-166—Its food, 166—Its ene-	
mics and parasites, 166—Its breeding habits, 166–167—Remedies,	
167-168-Geographical distribution of the species of Anabrus,	
168-169—Synopsis of the species of Anabrus and its allies, 169-	
170—External anatomy of Anabrus, 170: the head, 170-172; the	
thorax, 179-174; the abdomen, 174-175—Internal anatomy of	
Anabrus purpurascens, 175; the digestive system, 175 176; the	
nervous system, 176-177; the breathing apparatus, 177-178.	
CHAPTER IX: By A. S. Packard.	190
The air-sacs of locusts with reference to their powers of flight	178

sects, 178-179—The air-sacs of the Acridii, 179—Use of the air-sacs in flight, 179-182—Their origin, 182-183.

CHAPTER X: By C. S. Minot.

Histology of the locust (Caloptenus) and the cricket (Anabrus)......

Study of histology of insects, 183-184—Explanations of Figs. 1 and 2, Plate II, 185-186—The cuticula, 186-188—The epidermis, 188-189—Sense organs, 189—Nervous system, 190—Trachese, 191-195—Air-sacs and spiracles, 195—Muscles, 195-196—Organs of circulation, 196—Connective tissue, 196-198—Ovary, 198-201—Anterior execum of the oviduct, 201—Uterus of locusta, 201-203—Male organs, 203—Testis, 203-204—Development of the spermatozoa, 204, 207—Vasa deferentia, 207—Ductus ejaculatorius, 207—Vesicula seminales, 207-208—Digestive canal, 208-210—Crop, 210-211—Provertriculus, 211-212—Stomach, 212-214—Diverticula, 214-215—Gastro-ileal folds, 215-217—Heum, 217—Colon, 217-218—Rectum, 218-219—Summary on the digestive canal, 219-221—Malpighian vessels, 222.

CHAPTER XI: By A. S. Packard.

The nervous system in general, 223-224—The brain of insects compared with that of vertebrates, 224-226-The brain of the adult locust, 226—Histological elements of the brain, 226-228—The sections of the brain, 228-230-Internal topography of the brain, 230—The central body, 230, 231—The mushroom bodies, 231-234-The optic lobes, 234—The optic ganglion, 234—The antennal or olfactory lobes, 235—The commissural lobes, 235—The brain of locusts compared with that of other insects, 235-236-Structure of the brain in the embryo locust, 236—The brain of the embryo locust compared with the first thoracic ganglion, 238—The brain in the second embryonic stage, 238-239-Structure of the subcesophageal ganglion, 239—The brain of the freshly-hatched larva of Caloptenus spretus, 239; of the third larval stage of C. bivittatus, 239; of the second or last pupal stage of C. spretus, 240-241-Bibliography of the internal structure of the brain of crustacea and insects, 241-242.

CHAPTER XII: By C. V. Riley and C. Thomas.

marks on Cratypedes Putnami, 259.

Locust ravages in California

Position of the genus Camnula in the Œdipodini, 242-243—Description of the genus Camnula, 243; of Œdipoda pellucida, 243; of Œdipoda atrox, 243-244; of Camnula pellucida, 244-246—The locust in California in 1878, 246; in Sierra Valley, 247; in the vicinity of Loyalton, 247-248; of Sierraville, 249—History of locust devastation in California, 250—Enemies of the Californian locust, 251—Protective measures, 251-252—Life history of Camnula atrox, 252-253—The red or locust mite, 253—Damage done by the locust in Sierra Valley, 254-255—The locust in California in 1879, 255-257—Description of Œdipoda obliterata, 257-259—Re-

CHAPTER XIII: By C. V. Riley.

183

223

242

250

CHAPTER XIII-Continued.

eggs of Camnula pellucida, 263—Habits of bee-fly larvæ, 263–264—Observations on the larval habits of Bombylii, 264–265—Larval habits of Argyramæba, 265–266—Abundance of bee-flies and blister-beetles in the Western country, connected with the abundance of locusts, 266—Life history of Systæchus oreas, 266–267—Description of its larva, 267; of the pupa, 267–268; of the imago, 268—Description of Systæchus oreas, 268—Larva of Triodites mus, 268–269—Imago of Triodites mus, 269—Dr. T. A. Chapman's observations on Bombylius major, 269—Hair-worms and red-mites abundant upon locusts in California, 270—Chalcid-fly parasitic on locusts, 270—Synonymy of the locust-egg parasite, 270—Diggerwasps killing locusts, 270.

CHAPTER XIV: By C. V. Riley.

Courses that may be adopted by the General Government to lessen locust injury.....

271

Importance of destroying the locusts in their native breedinggrounds, 271-Recapitulation of means suggested in first report, 271-Importation of English rooks, 271-Importance of burning over the permanent breeding-grounds, 272-Breeding-grounds occupy comparatively small areas in the permanent region, 273— Protection from invading swarms, 273—Locust warnings through the Signal Bureau, 273—Diverting swarms by means of smoke, 274—Co-operation of governments and governmental institutions. 274—Apathy apt to result from periods of immunity from locust invasions, 274-How the Government can aid, 275-Surface characteristics of the permanent region and the proportion of burnable land, 275-Number of square miles in the permanent region, 276—The plains area east of the mountains, 276—Its vegetation, 276-Burnable land practically indentical with grazing-land, 277-The plains area in the British possessions, 278-In the United States, 279—The mountain area, 280—Timber-lands in the mountain area dependent upon latitude and altitude, 280-Northern section of the mountain area, 280, 281-River valleys in Montana, 281-Valley of the Yellowstone and its tributaries, 281-Of the Madison River, 282-Of the Jefferson and Missouri Rivers, 283—The Wind River and the Bighorn Mountains, 283— The Green River Basin, 283-Valley of the Snake River and its tributaries, 286, 289-Southern section of the mountain area, 289-In southern Wyoming, 290-291-In Colorado, 291-The San Luis Valley, 292-Mountain area in New Mexico, 292-The plateau area, 293-Its extent, 293-The Roan Plateau, 293-The Uinta Valley, 293—The Grand River Valley, 294—The Great Sage Plain, 294—The San Juan River Valley, 295—The Colorado River Plateau, 205-The Great Basin area and the Wasatch Mountains, 296-Valleys of the Bear River and its tributaries, 296-The Wasatch Range and its valleys, 298-Mountain Ranges in Nevada, 299-300-The Mojave Desert, 300-Preventive measures in the plains region, 300—Caloptenus spretus breeds, especially in British America and Montana, 300-Fertile grass-land in the Territories, 300-Its extent, 302-Eucouragement to settlement, 302-Modification of climate by settlement and cultivation, 302-Professor Thomas' communication to Governor Pillsbury, 303—The settlement of Dakota will benefit Minnesota, 304 - Forest planting

CHAPTER XIV-Continued.

on the coteau of the prairies, 305-Preserving of lakes, ponds, and swamps in Minnesota, 306—Encouragements to railroads, 307— Irrigation, 307—Extent of the arid region, 307-308—Only a small portion of the land irrigable, 308-Agriculture in the arid West and in the Mississippi Delta require assistance of the National Government, 308—Prevention of floods by means of reservoirs, 309—Cost of irrigation, 310—Influence of irrigation on the locust question, 310-Mr. H. Gannett's report on irrigation in the arid region, 310-Waste of water in irrigating, 311-Amount of irrigable land, 311-Amount of water necessary for irrigation, 312-313—Irrigable land in Colorado, Utah, California, &c., 313— Arable land in the Territories, 314—Amount of land in actual cultivation, 314-Irrigation by means of artesian wells, 315-Methods of irrigation, 315-316-Uselessness of reservoirs and great irrigating canals in the arid region, 316-How irrigation should be regulated, 316—Preservation of forests; tree-planting, 317— Judicious burning in spring, 317-Permanent establishment of locust signals and warnings, 318-Co-operation with the Dominion Government, 319-Preventive measures in the mountain and plateau areas, 319-Breeding-grounds of the locust in the mountain area, 319-Influence of settlement, 320-Replanting of forests, 320-Importance of locust signals to the farmer in Colorado and Utah, 321-Very destructive locust invasions not likely to recur in the future, 321—The locust has ceased to be an object of dread, 322.

APPENDIX I:

Miscellaneous data and replies to Circular No. 1.... Circular No. 1, [3]—Nebraska data, [4]—Data for Minnesota, [17]— Data for Iowa, [20]—Data for Dakota, [21].

APPENDIX II: By S. H. Scudder.

List of the Orthoptems collected by Dr. A. S. Packard, jr., in the western United States in the summer of 1877..... Bradynotes opimus, Scudd., n. sp., [24] - Pezotettix pacificus, Scudd., n. sp., [24]—Gomphocerus shastanus, Scudd., n. sp., [25]— Circotettix maculatus, Scudd., n. sp., [26]—Trimerotropis latifasciata, Scudd., n. sp., [26]-T. similis, Scudd., n. sp., [27]-T. coruleipes, Soudd., n. sp., [27]—Psinidia wallula, Soudd., n. sp., [27].

APPENDIX III:

Report of John Marten..... Observations of John Marten, special agent, on locusts in Iowa, [22]—In Minnesota, [30]—In Dakota, [30]—In Nebraska, [31].

APPENDIX IV: By B. P. Mann and C. Thomas.

Bibliography of some of the literature concerning destructive locusts.....

Introductory and explanatory remarks, [33]—Bibliography on the destructive locusts of Europe, Asia, and Africa, [33]-Supplementary list to the bibliography on locusts in foreign countries, [50]—Additions to bibliography on the locusts in America, [55].

APPENDIX V:

Notes furnished by Gen. A. J. Myer, U. S. A., Chief Signal Officer, on locusts in Texas, chiefly in the year 1877, [57]-Locust data from Indian Territory in 1877, [61].

[3]

[29]

[33]

1 959.	[RILEY, C. V.]—Continued.	•
	APPENDIX VI:	
	On the flight of locusts.	[63]
	Translation, by F. P. Spofford, of a memoir by Gaetano de Lucre-	
	tus, [63]—Translation of d'Azara's paper on locust invasions in	
	Spain from 1754 to 1757, by F. P. Spofford, [66].	
	APPENDIX VII:	
	Notes of a journey made to Utah and Idaho in the summer of 1878, by A. S. Packard, jr	[69]
	Observations on locusts in Nebraska and Colorado, [69]—In Wy-	[]
	oming, [69]—In Utah, [69]—In Idaho, [70]—Observations on	
•	the journey returning East, [71].	
	APPENDIX VIII:	
	Yersin's researches on the functions of the nervous system of the articulate animals	[73]
	Translation, by Dr. A. S. Packard, jr., of Saussure's abstract from	
	Yersin's Recherches sur les fonctions du système nerveux dans	
	les animaux articulés, [73].	

- 1960. RILEY, C. V. Notes on North American Microgasters, with descriptions of new species. <Trans. Acad. Sci. St. Louis, 16 April 1881, v. 4, pp. 295-315, 9 figs. Separate: <[St. Louis, Mo.], 6 April, 1881, 20 pp., 9 figs.</p>
 - Characteristics and habits of the Microgasters; description of larva of Apanteles aletiæ; the cocoons and their construction; effect of the parasites upon the longevity of the host; secondary parasites; habits and descriptions of several new species and varieties, for a list of which see the Systematic Index; figures A. aletiæ, healthy and parasitized larvæ of Chærocampa pampinatrix [= Ampelophaga myron]; also mass of Microgaster cocoons, and the formation of the cocoon by the larva.
- 1961. [RILEY, C. V.] Exuviation in flight. < Amer. Nat., May [16 April], 1881, v. 15, p. 395.</p>
 Criticism of R. McLachlan's observations; process of casting subimaginal skin

in Polymitarcys alba.

- 1962. RILEY, C. V. The rascal leaf-crumpler in Georgia. <Amer. Nat., May [16 April], 1881, v. 15, p. 400.</p>
 Occurrence and ravages of Acrobasis nebulo [=indiginella] in Bryan County, Georgia.
- 1963. [RILEY, C. V.] Vertical vs. horizontal insect boxes. < Amer. Nat., May [16 April], 1881, v. 15, p. 401.

Review of pamphlet by A. Prendhomme de Borre; reply of G. H. Horn to Preudhomme de Borre.

- 1964. [RILEY, C. V.] Insects affecting the China tree. Amer. Nat.,
 May [16 April], 1881, v. 15, pp. 401-402.

 General immunity of the China tree from the attacks of insects and its consequent value as a shade tree; occurrence of a Lecanium and Coroplastes n. sp. upon it in Alabama; injuries to its leaves by Atta fervens in Texas.
- 1965. [RILEY, C. V.] Galls on Eucolyptus. < Amer. Nat., May [16 April], 1881, v. 15, p. 402.</p>
 Critical review of paper by R. McLachlan; the supposed dipterous galls prob-

ably cecidomyidous; the supposed lepidopterous galls probably not lepidopterous, but inhabited by a lepidopterous inquiline,

- 1966. [RILEY, C. V.] North American Anthonyide. < Amer. Nat. May [16 April], 1881, v. 15, p. 402.
 - Notice and summary of paper by H. A. Hagen.
- 1967. [RILEY, C. V.] Galls and gall-insects. < Amer. Nat., May [16] April, 1881, v. 15, pp. 402-403.
 - Review of paper by H. F. Bassett, with additional notes on the gall of Cymins [= Andricus] querous-californica found on Querous douglasti and infested by Ozognathus cornutus; the Cynips produced from this gall all females; carlier accounts of the gall and of the habits of O. cornutus.
- 1968. RILEY, C. V. Scale insect on raspberry. <Farmer's Review. 21 April, 1881, v. 6, p. 243. S.-b. No. 23, p. 150. Reprint: <Amer. Nat., June [19 May], 1881, v. 15, p. 487.
 - Remarks on communication from R. B. Fulton; occurrence and ravages of a scale (Diaspis harrisii 1) [= Chionaspis furfurus] on black-cap raspberry in Mississippi; mode of dissemination and means of extermination of Coccide.
- 1969. RILEY, C. V. Descriptions of some new Tortricidæ, leaf-rollers. <Trans. Acad. Sci. St. Louis, May, 1882, v. 4, pp. 316–324.</p> Separate: <[St. Louis, Mo.], 28 April, 1881, 9 pp.
 - Description of two (2) new genera and thirteen (13) new species of Tortricida; see the Systematic Index for a list of the same; mention of the food-plants of some of the species.
- 1970. RILEY, C. V. Larval habits of bee-flies. < Amer. Nat., June [19] May], 1881, v. 15, pp. 438-447, pl. 6. Separate: <pp. 438-447. Adapted from the 2d Rept. U. S. Ent. Commission, pp. 262-269; see No. 1959 for synopsis of contents.
- 1971. [RILEY, C. V.] The periodical Cicada alias "seventeen-year locust." <Amer. Nat., June [19 May], 1881, v. 15, pp. 479-482, Correction: < Ibid., July [22 June], 1881, p. 578.
 - Extract from 1st Rept. State Ent. Mo., with additional notes; figures eggs, pupes, and image of Cicada [= Tibicen] septendecim, and of twig punctured by the image for the deposition of her eggs.
- .372. BILEY, C. V. A new species of oak cocoid mistaken for a gall. <Amer. Nat., June [19 May], 1881, v. 15, p. 482.
 - Description of Kermes galliformis n. sp. [p. 482] occurring on Querous palustris in the southern and central United States; the coccid infested by Euclemensia bassettella.
- 1973. [RILEY, C. V.] The "water-weevil" of the rice-plant. < Amer. Nat., June [19 May], 1881, v. 15, pp. 482-483. Extract: <Ann. Rept. [U.S.] Commissioner Agric. for 1881 and 1882, pp. 130-133. Note on the injury done to rice plants by the larve of Chalepus trackypyque and by other larvæ, probably Lissorhoptrus simplex; letter from J. Screven; seasons, habits, and ravages of the last-mentioned species.
- 1974. [RILEY, C. V.] The impregnated egg of Phylloxera vastatrix. <Amer. Nat., June [19 May], 1881, v. 15, pp. 483, 484. Reprint: <Amer. Wine and Grape Grower, 1 July, 1881, v. 3, p. 104.</p> S.-b. No. 42, pp. 12-13.
 - Verification by P. Graell of author's conclusion that the impregnated egg of Phyllowera vastatric would generally be found to hatch in the same season in which it was laid.

- 1975. RILEY, C. V. Works on North American Microlepidoptera. Amer. Nat., June [19 May], 1881, v. 15, pp. 484-486.
 - Review of works of T. de Grey [Lord Walsingham]; discussion of new genera; list of Tortrioidæ of economic interest common to Europe and North America; additions to the synonomy of Tortrioidæ.
- 1976. [RILEY, C. V.] Moths mistaken for *Aletia*. <Amer. Nat., June [19 May], 1881, v. 15, pp. 486-487.

Platyhypena [= Hypena] scabra and Phoberia atomaris mistaken for Alstia ergillacea [= xylina]; hibernating habits of the first species.

- 1977. [RILEY, C. V.] Specific value of Apatura alicia Edw. < Amer.
 Nat., June [19 May], 1881, v. 15, p. 487.
 Critical review of paper by W. H. Edwards.
- 1978. BILEY, C. V. Antigaster vs. Eupelmus. <Ca. Ent., May, 1881, v. 13, p. 114.

Validity of L. O. Howard's reasons for considering Antigaster a synonym of Eupelmus.

- 1979. RILEY, C.V. The periodical Cicada alias "seventeen-year locust." <Farmer's Review, 16 June, 1881, v. 6, p. 370. S.-b. No. 42, pp. 18-20.
 - Extract from 1st Rept. State Ent. Mo., with additional notes and request for further information in regard to the chronology and geographical distribution of the broads of *Cicada* [= Tibicen] septendecim and C. [= T.] tredecim which appear in 1881.
- 1980. [RILEY, C. V.] Dimorphism in *Oynipidæ*. < Amer. Nat., July [22 June], 1881, v. 15, p. 566.
 - Claim of author to have made the first record at least in North America of proof of dimorphism in Cymipidæ; notice of writings of B. D. Walsh, H. F. Bassett, and H. Adler on the subject; list of species of Cymipidæ in which the occurrence of dimorphic forms has been proven, and of closely allied species in which no alternate generation seems to occur.
- 1981. [RILEY, C. V.] Blepharocerides. <Amer. Nat., July [22 June], 1881, v. 15, pp. 567-568.

 Assume of various investigations into the natural history of Blenharocerides.
 - Account of various investigations into the natural history of Blepharocerida; description of larve and pupe of these flies.
- 1982. [RILEY, O. V.] Braula cæca not particularly injurious to the honey-bee. Amer Nat., July [22 June], 1881, v. 15, p. 568.

 Notice of paper of J. Fedarb; ravages of Braula cæca in hives generally overestimated at present and formerly unknown.
- 1983. [RILEY, C. V.] Economic entomology in England. < Amer. Nat., July [22 June], 1881, v. 15, p. 568.

 'Notice of Miss E. A. Ormerod's report for 1880.
- 1984. [BILEY, C. V.] The cultivation of pyrethrum and manufacture of the powder. <Amer. Nat., 1881, v. 15. July [22 June], pp. 569-572; September [23 August], pp. 744-746; October [23 September], pp. 817-819. See: <Amer. Wine and Grape Grower, December, 1881, v. 4, p. 22. S.-b. No. 51, p. 149.

٠١.

Partial history of the use of pyrethrum powder as an insecticide; directions for the cultivation of the plants and for the production and use of the powder

- 1985. [RILEY, C. V.] Hudson Bay Lepidoptera. < Amer. Nat., July [22 June], 1881, v. 15, pp. 572, 573.
 - Review of paper of J. J. Weir; reprint of the list of Lepidopters and of the remarks on the explanation of the relations of the fauna of Hudson's Bay to that of Europe; probability that the species identical with those of Europe are recent immigrants.
- 1986. [RILEY, C. V.] Trade in insects. < Amer. Nat., July [22 June], 1881, v. 15, p. 573.
 - General decline in prices of insects within sixty years or more; statement of some prices obtained for Coleoptera recently.
- 1987. [RILEY, C. V.] Ants injurious in Arizona. <Amer. Nat., July [22 June], 1881, v. 15, pp. 573, 574.
 - Statement by H. H. Rusby of ravages of Formicidæ, which occur in vast colonics.
- 1988. [RILEY, C. V.] Covering of egg-puncture mistaken for *Dorthesia*. <Amer. Nat., July [22 June], 1881, v. 15, p. 574.
 - Waxy material covering the egg-punctures of Enchophyllum [= Enchenops] binotata, labeled Dorthesia viburni and D. celastri in collection of A. Fitch.
- 1989. [RILEY, C. V.] [Dolerus unicolor.] <Amer. Nat., July [22 June], 1881, v. 15, p. 574.
 - Doubts the statement of H. Keenan that the images of *Dolerus unicolor* injure the fruit buds of pear-trees.
- 1990. [RILEY, C. V.] Supposed army-worm in New York and other Eastern States. <Amer. Nat., July [22 June], 1881, v. 15, pp. 574-577.
 - Ravages of Nephelodes violans and Crambus vulgivagellus in New Jersey, Long Island, and northern New York; natural history, vernacular names, and description of the larva of the former species; previous accounts of this larva
- 1991. [RILEY, C. V.] Migration of butterflies. <Amer. Nat., July [22 June], 1881, v. 15, p. 577.
 - Report by J. H. Mellichamp of the eastward flight of thousands of *Pieris me-*susts in small groups of two, three, or more individuals, on 1 and 2 June,
 1881, over Bluffton, S. C.; description of larva and pupa; food-plants of
 larva.
- 1992. [RILEY, C. V.] Classification of the mites. < Amer. Nat., July [22 June], 1881, v. 15, pp. 577-578.
 - Abstract of letter of G. Haller; observations on the appendages and systematic position of the Acarina.
- 1993. [RILEY, C. V.] Carrying out the law. < Amer. Nat., July [22 June], 1881, v. 15, p. 578.
 - A fine of £5 imposed upon a man in England for importing living Doryphors decembineata into that country.
- 1994. RILEY, C. V. Locusts and locusts. <N. Y. Tribune, 22 June, 1881. S.-b. No. 32, pp. 55-56; No. 42, pp. 7-11.
 - Cicada [= Tibicen] septendecim and C. [T.] tredecim compared with Caloptenus spretus to explain and correct the confusion of the species by those who call them all "locusts"; differences in the habits and characters of these

- 1994. RILEY, C. V.—Continued.
 - insects, their periodicity and distribution; no fear that *C. spretus* will commit great ravages this year; recommendation of a system of observations and warnings by the United States Signal Service to guard against unexpected inroads of *C. spretus*.
- 1995. RILEY, C. V. The caterpillar nuisance. <Evening Star [Washington, D. C.], 24 June, 1881, v. 57, No. 8802, p. 3. S.-b. No. 23, p. 152; No. 42, p. 13. Reprint: <Amer. Nat., September [23 August], 1881, v. 15, pp. 747-748, 1 fig.
 - Natural history of *Hyphantria textor* [= cunea]; means against it; methods of applying poisoned liquids to trees.
- 1996. RILEY, C. V. Directions for cultivating pyrethrum for insect powder. <Gardener's Mo. and Hortic., June, 1881, v. 23, pp. 172-173. S.-b. No. 23, p. 153.
 - Directions for raising Pyrethrum cineraria folium from seed; climate not favorable to the growth of P roseum; preparation of these plants for the destruction of insects.
- 1997. RILEY, C. V. Cotton-worms and Cicadas. Prof. Stelle's logic! <Selma [Ala.] Times, 19 July, 1881. S.-b. No. 24, pp. 44-45; No. 39, p. 123.
 - Criticism of statements of J. P. Stelle; impossibility of exterminating Alstia xylina; waves of destructiveness; Cicada [= Tibicen] tredecim abundant in sections of Alabama, as predicted.
- 1998. RILEY, C. V. The Rocky Mountain locust alias Western grass-hopper. <Amer. Agric., July, 1881, v. 40, pp. 283-284, 6 figs. S.-b. No. 23, p. 142.
 - Summary of facts concerning the egg-laying development, habits, food-plants, destructive powers, migration, flight, and enemies of and means against Caloptenus spretus; figures of the eggs of the insect in all stages, of the manner of oviposition, and of pans for the destruction of the locusts; prospects of future injury.
- 1999. RILEY, C. V. Lepidopterological notes. <Papilio, July, 1881, v. 1, pp. 106-110.
 - Advance print of extracts from Bull. No. 6, U. S. Entomological Commission, pp. 56-58, 78, 82-83. See No. 2026 for synopsis of contents.
- 2000. RILEY, C. V. Further notes on the pollination of Yucca and on *Pronuba* and *Prodoxus*. <Proc. Amer. Assoc. Adv. Sci. for 1880, 14 October, 1881, v. 29, pp. 617-639, 16 figs. Separate: <[Salem, Mass, July, 1881], 23 pp., 16 figs.
 - Recapitulation of published observations on Pronuba yuccasella and Prodoxus decipiens, with additional observations and references to literature; habits and functions of the two species; descriptions and figures of the generic characters of Pronuba, Prodoxus, and Hyponomeuta, and of the specific characters of Pronuba yuccasella, P. maculata n. sp., Prodoxus decipiens, P. intermedius n. sp., P. marginatus n. sp., P. cinercus n. sp., P. conescens n. sp., Hyponomeuta malinella, and H. multipunctella, especially the structure of the ovipositor and the male characters, and the venation of the wings; discussion of the structure of the ovipositor in Lepidoptera characterization of the new family Prodoxida to comprise Pronuba and Prodoxus; generic and specific synonymy; reply to criticism of T. Meehan; list of insects frequenting Yucca; synonymical list of the described Prodoxia.

- 2001. RILEY, C. V. Additional notes on the army worm, Lewomic unipuncta. <Proc. Amer. Assoc. Adv. Sci. for 1880 [14 October], 1881, v. 29, pp. 640-642. Separate: <Salem, Mass., July, 1881, pp. 24-26. Abstract: <Boston Daily Advertiser, 2 September, 1880, v. 136, p. 4.</p>
 - Number of annual generations of Loucania unipuncta; stages at which hibernation takes place; occasional immigration of moths into uninfested districts; preferred breeding-places; connection of wet and dry seasons with the abundance of these insects.
- 2002. RILEY, C. V. The hitherto unknown life-habits of two genera of bee-flies, Bombyliidæ. <Proc. Amer. Assoc. Adv. Sci. for 1880 [14 October], 1881, v. 29, p. 649. Separate: <Salem, Mass., July, 1881, p. 33.
 - Food-animals and food-habits of Systechus oreas and Triodites mus; parallelism in the life-history of Bombyliidæ and Meloidæ; connection of the abundance of insects of these families in the western United States with the prevalence of Acrididæ there; retardation of development in the early stages of these parasites explained as a beneficial characteristic.
- 2003. RILEY, C. V. A remarkable case of retarded development. <Sci. Amer., 20 August, 1881 [v. 59], n. s., v. 45, p. 116. S.-b. No. 23, p. —.
 - Report of a case in which eggs of Caloptenus spretus remained undeveloped for a period of four and one-half years after being laid; remarks upon the problems involved in the explanation of retarded development.
- 2004. [RILEY, C. V.] Blepharoceridæ. <Amer. Nat., September [23 August], 1881, v. 15, p. 748.
 - Discovery by J. Q. Adams of pupe and images of *Blepharocerida* at Watertown, N. Y.
- 2005. [RILEY, C. V.] Remarkable case of retarded development. Amer.">Nat., September [23 August], 1881, v. 15, pp. 748-749.
 - Eggs of Caloptenus spretus, buried about 25 cm. (10 inches) under ground, remained unhatched and alive for four and one-half years, and hatched upon being exhumed.
- 2006. [RILEY, C. V.] Promotion of silk-culture in California. < Amer. Nat., September [23 August], 1881, v. 15, p. 749.
 - Answer to letter of Mrs. T. H. Hittell; popularization of silk-culture in California; preparation of the fiber for the market.
- 2007. [RILEY, C. V.] Locust flights in Dakota. >Amer. Nat., September [23 August], 1881, v. 15, pp. 749-750.
 - Flights of Caloptenus spretus in Dakota on 7th and 16th July, 1881; large numbers of Diplax rubicundula accompanying the locusts.
- 2008. [RILEY, C. V.] The Hessian fly. <Amer. Nat., September [23 August], 1881, v. 15, p. 750.
 - Extensive damages done by Cecidomyia destructor in Illinois and Missouri; its abundance in the western prairie States in 1881.

- **2909.** [RILEY, C. V.] The genuine army-worm in the West. < Amer. Nat., September [23 August], 1881, v. 15, p. 750.
 - Occurrence and ravages of Loucania unipuncia in Illinois and Indiana in 1881; these larves either from eggs of hibernated moths or of a second brood; other insects mistaken for this in other States; periods of the Loucania and influence of the seasons upon it.
- 2010. [RILEY, C. V.] A new imported enemy to clover. <Amer. Nat.,

 September [23 August], 1881, v. 15, pp. 750-751.

 Clover injured by Phytonomus punctatus at Barrington, N. Y., in July, 1881.
- 2011. [RILEY, C. V.] Another enemy of the rice-plant. <Amer. Nat., September [23 August], 1881, v. 15, p. 751.</p>
 Rice-plants injured by the larva of Laphygma frugiperds in Georgia in the summer of 1861.
- 2012. [RILEY, C. V.] Canker-worms. < Amer. Nat., September [23 August], 1881, v. 15, p. 751.
- gust, 1881, v. 19, p. 791.

 Excessive ravages of Paleacrita vernata in Tazewell County, Illinois, in 1881.
- 2013. [RILEY, O. V.] Lepidopterological notes. < Amer. Nat., September [23 August], 1881, v. 15, pp. 751-752.
 Advance extract from Bull. No. 6, U. S. Entomological Commission, pp. 55-56. See No. 2026 for synopsis of contents.
- 2014. RILEY, C. V. The periodical Cicada. < Amer. Agric., August, 1881, v. 40, p. 132, 5 figs. S.-b. No. 23, p. 141.
 - Brief sketch of the natural history of Cicada [= Tibicen] septendecim and C. [= T.] tredecim, with figures of all stages and of punctured twigs; the reputed stinging by these insects probably done by Stisns grandis [= Sphecius speciosus]; figure of the Stisns; recommends the substitution of the name "periodical Cicada" in place of the name "locust."
- 2015. RILEY, C. V. Entomological notes. <Farmer's Review, 22 September, 1881. S.-b. No. 42, pp. 54-55. Reprint: <Amer. Nat., [3] December, 1881, v. 15, pp. 1012-1013. See: <Sci. Amer., 11 February, 1882. S.-b. No. 38, p. 1½.
 - Abstract and critical review of paper by M. Saint-André; attempted explanation of the immunity of grape-vines in sandy soil from the ravages of *Phyl-loxera vasiatrix*.
- 2016. RILEY, C. V. Notes on *Hydrophilus triangularis*. <Amer. Nat., October [23 September], 1881, v. 15, pp. 814-817, figs. 1-2.

 Description and figures of egg-case and figures of eggs and of male imago,
 - with structural details of Hydrophilus triangularis; characters and life-history of the larva and pupa; figures of eggs, egg-case, larva, and pupa of H. piceus of Europe.
- 2017. [RILEY, C. V.] Migration of plant-lice from one plant to another.
 <Amer. Nat., October [23 September], 1881, v. 15, pp. 819-820.</p>
 Exposition of J. Lichtenstein's theory that most Aphidida, especially gall-making Pemphigini, live upon two different plants in passing through their cycle of development; cites instances in which the host-plants are of different families.
- 2018. [RILEY, C. V.] The chinch-bug. <Amer. Nat., October [23 September], 1881, v. 15, pp. 820-821.
 - The abundance of Blissus loucopterus is connected with the moisture of the season; extraordinary noxiousness of this insect in the dry season of 1881 in various parts of the United States.

- 2019. [RILEY, C. V.] Phylloxera laws. < Amer. Nat., October [23 September], 1881, v. 15, p. 821.
 - Summary of the existing laws regulating the traffic in plants, with a view of preventing the introduction of Phyllozera vastatrix into different countries.
- 2020. [RILEY, C. V.] One half the vine area of France affected by Phylloxera. <Amer. Nat., October [23 September], 1881, v. 15, p. 821.
 - Statement that nearly one-fourth of the area of vineyards in France is destroyed by *Phylloxera vastatrix* and as much more attacked; submersion the only sure remedy, and the use of American resisting stocks the only available preventive.
- 2021. [RILEY, C. V.] London purple and Paris green. <Amer. Nat., October [23 September], 1881, v. 15, p. 821.</p>
 London purple more than twice as efficacious and less expensive than Paris green as an insecticide; it needs to be well ground; composition of London purple.
- 2022. [RILEY, C. V.] Entomologist for the Pacific coast. < Amer. Nat.,
 October [23 September], 1881, v. 15, pp. 821–822.
 Notice of attempts made to have a State entomologist appointed in California.
- 2023. [RILEY, C. V.] Dilar in North America. <Amer. Nat., October [23 September], 1881, v. 15, p. 822.</p>
 Notice of paper by R. McLachlan.
- 2024. [RILEY, C. V.] Locusts in Nevada. < Amer. Nat., October [23 September], 1881, v. 15, p. 822.
 - Statement from Reno [Nev.] Journal that the countless locusts which hatched in western Nevada in the spring of 1881 flew toward the Sierra Nevada without doing any injury.
- 2025. [RILEY, C. V.] Odor in butterflies. <Amer. Nat., October [23 September], 1881, v. 15, p. 822.

 Notice of paper by Miss M. E. Murtfeldt.
 - 2026. RILEY, C. V. General index and supplement to the nine reports on the insects of Missouri. <Bull. No. 6, U.S. Ent. Commission [24 March], 1881, 178 pp.
 - Order of matter: (1) Introduction. (2) Table of contents. (3) Corrections. (4) Notes and additions. (5) Descriptions of new species and varieties. (6) List of descriptions of adolescent states. (7) List of descriptions, mostly amplified, of species not new. (8) List of illustrations by reports. (9) Classified list of illustrations. (10) General index. (11) Index to foodplants.
 - 2027. [RILEY, C. V.] The permanent subsection of entomology at the recent meeting of the A. A. A. S. < Amer. Nat., 1881, v. 15, November [28 October], pp. 909-912; [3] December, pp. 1008-1011.
 - Notice of the meeting at Cincinnati, Ohio, August, 1881, with abstracts and notices of papers read.
 - 2028. RILEY, C. V. The new imported clover enemy. <Amer. Nat., November [28 October], 1881, v. 15, pp. 912-914.
 - Occurrence and habits of *Phytonomus punctatus* in New York; description of the egg and larva of the same; food-plants of the European species of *Phytonomus*.

- 2029. RILEY, C. V. Crambus vulgivagellus. <Amer. Nat., November [28 October], 1881, v. 15, pp. 914-915.
 - Excessive abundance of this species in the eastern United States in 1881; description of the egg; method of oviposition.
- 2030. RILEY, C. V. Larval habits of Sphenophori that attack corn. Amer. Nat., November [28 October], 1881, v. 15, pp. 915-916. Several species of Sphenophorus injurious to maize-plants in different parts of the United States; habits and ravages of S. robustus.
- 2031. [RILEY, C. V.] Effect of drought on the Hessian fly. <Amer. Nat., November [28 October], 1881, v. 15, p. 916. See: <N. E. Farmer, 10 December, 1881. S.-b. No. 45, p. 39. <Farmer's Review, 15 December, 1881. S.-b. No. 45, p. 99.
 - Hot and dry weather dries up and kills Cecidomyia destructor and its parasites.
- 2032. [RILEY, C. V.] Simulium from Lake Superior. <Amer. Nat.,

 November [28 October], 1881, v. 15, p. 916.

 Notice of paper by H. A. Hagen: lawys and pupp from Lake Superior similar.
 - Notice of paper by H. A. Hagen; larvæ and pupæ from Lake Superior similar to those of S. pictipes, but the images from the same locality are different.
- 2033. [RILEY, C. V.] Coleopterous cave fauna of Kentucky. < Amer. Nat., November [28 October], 1881, v. 15, pp. 916, 917.</p>
 Notice of H. G. Hubbard's investigations; Adelops contains but one species, which seems to be equally common in all caves in the State; Anaphthalmus contains several species, and more may be expected to be found.
- 2034. [RILEY, C. V.] Hemipterological studies. < Amer. Nat., November [28 October], 1881, v. 15, p. 917.</p>
 Notice of paper of V. Signoret.
- 2035. [RILEY, C. V.] Entomology in Buffalo, N. Y. < Amer. Nat., November [28 October], 1881, v. 15, p. 917.
 Notice of Vol. 4, No. 1, of the Bulletin of the Buffalo Society of Natural Science; list of entomological papers therein.
- 2036. [RILEY, C. V.] Lampyridæ. <Amer. Nat., November [28 October], 1881, v. 15, p. 917.
 Notice of paper by J. L. Leconte.
- 2037. [RILEY, C. V.] Severe cold and hibernating apple-worms. Amer.Nat., November [28 October], 1881, v. 15, p. 917.

 According to A. J. Cook larves of Carpocapsa pomonella were killed in their cocoons where exposed to the severe cold of the winter of 1880–1881.
- 2038. RILEY, C. V. [Address delivered 4 November, 1881, at the cotton convention held in Atlanta, Ga., 2-4 November, 1881.] <[U. S. Department of Agriculture.] Address of Hon. George B. Loring... and other proceedings of the cotton convention, Washington, 1881, pp. 19-35. Reprint: <Atlanta Constitution, 5 November, 1881. S.-b. No. 24, p. 34.</p>
 - Beneficial and injurious influence of insects; methods of counteracting the same; ravages and natural history of and search for means against Alstia argillacea [=xylina]; improved methods and contrivances for the application of poisons to plants.

- 2039. RILEY, C. V. The chinch bug. <Amer. Agric., November, 1881, v. 40, p. 476, figs. 1-3. S. b. No. 42, pp. 21-23.

 Rayages food plants and natural history of Rlissus legenterus: descriptions.
 - Ravages, food-plants, and natural history of Blissus leucopterus; descriptions and figures of its several stages and of dimorphic form of the imago; effect of wet weather upon it.
- 2040. [RILEY, C. V.] Retarded development in insects. < Amer. Nat., [3] December, 1881, v. 15, pp. 1007-1008. Reprint: < Proc. Amer. Assoc. Adv. Sci. for 1881, [13 October], 1882, v. 30, pp. 270, 271. Notice: < Gardener's Chronicle, 27 May, 1882, v. 17, pp. 708-709. S.-b. No. 42, p. 31.
 - Eggs of Caloptenus spretus retained their vitality four and one-half years under abnormal environment and then hatched on exposure to normal conditions; speculations on the cause of the phenomena of retardation of development.
- 2041. [RILEY, C. V.] Preparation of Diptera. Amer. Nat., [3] December, 1881, v. 15, p. 1008.

 Notice of paper of J. Mik.
- 2042. [RILEY, C. V.] Another herbivorous ground-beetle. Amer.
 Nat., [3] December, 1881, v. 15, p. 1011.

 Anisodactylus confusus injuring strawberry-plants in California.
- 2043. [RILEY, C. V.] A disastrous sheep parasite. <Amer. Nat., [3]

 December, 1881, v. 15, p. 1011.

 A parasite (Trickodectes ovis?) doing great injury to sheep in Illinois.
- 2044. [RILEY, C. V.] Locusts in the West. < Amer. Nat., [3] December, 1881, v. 15, p. 1013.
 - Caloptenus spretus scarce in Colorado and Kansas; Camnula pellucida abundant on the Pacific coast; destructiveness of locusts in many parts of South America and of Pachytylus migratorius in Turkey; means adopted against the latter.
- 2045. [RILEY, O. V.] Structure of the claw in *Psocina*. <Amer. Nat., [3] December, 1881, v. 15, pp. 1013-1014.

 Notice of paper of H. A. Hagen.
- 2046. [RILEY, C. V.] Insect collection for sale. <Amer. Nat., [3] December, 1881, v. 15, p. 1014.</p>
 Notice of the collection of Coleoptera left by C. Trabrandt.
- 2047. RILEY, C. V. Peach-tree bark-borer. Important note from Prof. C. V. Riley. <Rural New Yorker, 24 December, 1881, v. 40, p. 866. S. b. No. 42, p. 13.
 - Habits and ravages of Phlaotribus liminaris; means against bark-borers.
- 2048. RILEY, C. V. The chinch-bug. <Amer. Agric., December, 1881, v. 40, p. 515, figs. 1-4. S.-b. No. 42, pp. 23-26.
 - Figures Anthocoris [= Triphleps] insidiosus and Harpactor [= Milyas] cinctus as enemies of Blissus leucopterus and Nysius destructor [= angustatus] and Piesma cinerea as likely to be confounded with it; mentions other enemies of the chinch-bug; means against it; importance of irrigation.
- 2049. RILEY, C. V. On the oviposition of Prodoxus decipiens. <Amer. Nat., January, 1882 [30 December, 1881], v. 16, pp. 62-63. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1881 13 October],</p>

- 2049. RILEY, C. V.—Continued. 1882, v. 30, p. 272. Separate: <[Salem, Mass., February, 1882], p. 3.
 - Time and manner of oviposition of Prodoxus decipiens in stem of Yucca filamentosa.
- 2050. [RILEY, C. V.] Clover insects. Amer. Nat., January, 1882 [30 December, 1881], v. 16, p. 63.

 Notice of paper of J. A. Lintner.
- 2051. RILEY, C. V. Horn's classification of the *Carabidæ*. <Amer. Nat., January, 1882 [30 December, 1881], v. 16, pp. 63-64. Notice of paper of G. H. Horn.
- 2052. [RILEY, C. V.] The butterfly-trees of Monterey again. <Amer Nat., January, 1882 [30 December, 1881], v. 16, p. 64.

 Swarming and migrating habits of Danais archippus; hibernation of the same.
- 2053. [RILEY, C. V.] Interest felt in economic entomology in California. <Amer. Nat., January, 1882 [30 December, 1881], v. 16, p. 65.
 - Notice of a call issued by the Board of State Horticultural Commissioners of California for a State convention to consider horticultural subjects, including means against insects.
- 2054. [RILEY, C. V.] Obituary. < Amer. Nat., January, 1882 [30 December, 1881], v. 16, p. 65.</p>
 Biographical notice of J. D. Putnam; announcement of the death of G. V. Muiszoch.
- 2055. RILEY, C. V. New insects injurious to agriculture. <Amer. Nat., February [25 January], 1882, v. 16, pp. 151-152. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1881, [13 October], 1882, v. 30, pp. 272-273.
 - The occurrence of previously unnoticed noxious insects due to, 1, the introduction of species from abroad; 2, previously existing unnoticed species; and 3, the acquisition of new habits by previously known innoxious species; in the last case the acquisition of new habits may be accompanied by the acquisition of new descriptional character, thus constituting a new species; new species thus become developed within brief periods of time.
- 2056. [RILEY, C. V.] New entomological periodicals. <Amer. Nat., February [25 January], 1882, v. 16, pp. 152–153. Notice of several prospective periodicals.
- 2057. [RILEY, C. V.] Locust probabilities for 1882. < Amer. Nat., February [25 January], 1882, v. 16, p. 153. Favorable prospects of immunity from ravages of Caloptenus spretus in 1882.
- 2058. [RILEY, C. V.] Entomological notes. < Amer. Nat., February [25 January], 1882, v. 16, p. 153.

 Notice of J. J. Weir's paper on the "Lepidoptera of the Outer Hebrides," etc.
- 2059. [RILEY, C. V.] Bibliography of gall literature. <Amer. Nat., March [24 February], 1882, v. 16, pp. 246-247. Notice of the record of F. Thomas,

- 2060. RILEY, C. V. A new depredator infesting wheat-stalks. < Amer. Nat., March [24 February], 1882, v. 16, pp. 247-248, fig. 1.
 - Isosoma allynii French is a species of Eupelmus probably parasitic on some of the wheat-stalk feeders and on some species of Chlorops; habits and figures of larva and description of female image of Isosoma tritici [p. 247]; comparisons between I. tritici and I. hordei; alleged habits of I. lineare of Europe; relative conspicuousness of the "humeral spot" in European, American, and Australian species of Isosoma. See No. 2063.
- 2061. RILEY, C. V. Further notes on the imported clover-leaf weevil (*Phytonomus punctatus*). <Amer. Nat., March [24 February], 1882, v. 16, pp. 248, 249.
 - Identity of *Phytonomus opimus* Lec. with *P. punctatus* Fab.; instances in which injurious insects have been overlooked for a long time; probability that plant-feeding Coleoptera imported from Europe will not spread far from the Atlantic coast; some species, injurious in Europe, are thus far innoxious in America; hibernation of and means against *P. punctatus*.
- 2063. RILEY, C. V. The wheat Isosoma. A new depredator infesting wheat-stalks. <Rural New Yorker, 4 March, 1882. S.-b. No. 42, pp. 55-56. See: <Sci. Amer. Suppl., 17 June, 1882. S.-b. No. 37, p. 57.</p>
 - Description of female image of Isosoma tritici; habits, ravages, and distribution of and means against the same; comparison with related species; a probable parasite of it, described as Isosoma allynii by G. H. French, belongs to the genus Eupelmus. See No. 2060.
- 2064. RILEY, C. V. Possible food-plants for the cotton-worm. < Amer.
 Nat., April [22 March, 1882], v. 16, pp. 327-329. Reprint:
 <Rept. [U. S.] Com. Agric. for 1881 and 1882, 1882 [January, 1883], pp. 164-166. Separate: < pp. 164-166.
 Food-plants of Alctia xylina. See No. 2119 for synopsis of contents.
- 2065. [RILEY, C.V.] Mode of feeding on the larva of *Dytiscus*. < Amer. Nat., April [22 March], 1882, v. 16, p. 330.</p>
 Notice of paper of E. Burgess; structure of the mouth of the larva of *Dytiscus*; manner in which this larva feeds.
- 2066. [RILEY, C. V.] Entomological notes. < Amer. Nat., April [22 March], 1882, v. 16, p. 330.

 Brief notices of several articles recently published with items of news.
- 2067. [RILEY, C. V.] Riley's researches. Orange insects and the cotton-worm; how the orange insect operates and the cure for its ravages; the hibernation of the cotton-worm; settlement of a mooted question. <Florida Daily Times, 29 March, 1882. S.-b. No. 42, pp. 28-29.
 - Report of interview with C. V. Riley; discovery by W. H. Ashmead that "orange rust" is caused by a mite; habits of and means against the same;

- 2067. RILEY, C. V.—Continued.
 - successful application by H. G. Hubbard of an emulsion of kerosene and milk as a means against this mite and against scale-insects; proof of the hibernation of Alstia argillacea [=xylina] in the extreme south of the United States; means against the Alstia.
- 2068. RILEY, C. V. Prof. C. V. Riley and the Yucca moth. < Gardener's Mo. and Hortic., March, 1882, v. 24, p. 92. S.-b. No. 23, pp. 198-199.
 - Critical review and reply to editorial of T. Meehan.
- 2069. RILBY, C. V. The *Noctuidæ* in the Missouri entomological reports. <Papilio, March, 1882, v. 2, pp. 41-44. Separate: <N. Y., 1882, 4 pp.
 - Critical review of paper of A. R. Grote; discussion of the synonymy of the Noctuidæ in the Missouri entomological reports.
- 2070. RILEY, C. V. Silk-culture in the United States. <Sci. Amer., 1
 April, 1882 [v. 60], n. s., v. 46, p. 193. S.-b. No. 42, pp. 93-96.
 Statement of the dangers to be avoided and the obstacles to be overcome for the successful establishment of silk-culture in the United States.
- 2071. RILEY, C. V. Little known facts about well known animals. A lecture delivered in the National Museum, Washington, D. C., April 8, 1882. < Washington, April, 1882, 32 pp., figs. See: < National Farmer, 13 July, 1882. S.-b. No. 42, p. 15. < Prairie Farmer, 22 July, 1882. S.-b. No. 46, p. 99.
 - The entomological subjects include brief and popular illustrated accounts of the life-histories and transformations of the house-fly (Musca domestica) and mosquito (Culex pipiens); the development of Trombidium from Astoma; parasitism; truth of the theory of evolution; reasons for the prevalent ignorance of natural history; interest and value of a knowledge of natural history.
- 2072. RILEY, C. V. Lichtenstein's theory as to dimorphic asexual females. <Amer. Nat., May [24 April], 1882, v. 16, p. 409. Criticism of J. Lichtenstein's theory that winged female Aphididæ and asexual female Cynipidæ are larvæ, and their eggs pupæ; errors in the comparison
- 2073. [RILEY, C. V.] Naphthaline cones for the protection of insect collections. <Amer. Nat., May [24 April], 1882, v. 16, pp. 409-410.

of these forms with the hypermetamorphotic stages of Meloida.

- Use of and objections to the use of naphthaline cones; formula of a preferable protective mixture.
- 2074. [RILEY, C. V.] Injurious insects in California. <Amer. Nat., May [24 April], 1882, v. 16, p. 410.
 Notice of treatise of Matthew Cooke.
- 2075. [RILEY, C. V.] Sarcophaga lineata destructive to locusts in the Dardanelles. <Amer. Nat., May [24 April], 1882, v. 16, pp. 410-411.
 - Notice of communications, by F. Calvert, regarding the parasitic habits of Saroophaga lineata, which preys upon Œdipoda cruciata; egg-pods of the same locust destroyed by larvæ of Callostoma fascipennis.

- 2076. [RILEY, C. V.] Parasitic Diptera. < Amer. Nat., May [24 April], 1882, v. 16, p. 411.
 - Dilophus parasitic in larvæ of Chetoptria hypericans; the Bibionida have hitherto been known only as vegetable feeders in the larval state.
- 2077. RILEY, C. V. The cotton worm. <Times Democrat [New Orleans], 7 May 1882, v. —, p. 4. S. b. No. 42, pp. 29-31; 112-113; No. 63, pp. 47-48. Reprint: <Rept. [U. 8.] Com. Agric. for 1881 and 1882, 1882 [January, 1883], pp. 157-159. Separate: <pp. 97-99.

See No. 2119 for synopsis of contents.

- 2078. RILEY, C. V. The house-fly. <Prairie Farmer, 13 May, 1882. S.-b. No. 39, p. 122. See: <Colman's Rural World, 15 June, 1882. S.-b. No. 68, p. 194. <Times-Democrat [New Orleans], 24 June, 1882. S.-b. No. 42, p. 31. <N. E. Farmer, 5 August, 1882.
 - Wide distribution of Musca domestica; identity of M. harpyia with the species; description of eggs; habits, transformations, and hibernation.
- 2079. RILEY, C. V. Habits of Cybocephalus. < Amer. Nat., June [20 May], 1882, v. 16, p. 514.

 Cybocephalus pigritulus feeds on scales of Chionasnia vinifolii on Pinus elliottii in
 - Cybocephalus nigritulus feeds on scales of Chionaspis pinifolii on Pinus elliottii in South Carolina, and C. californicus on a scale on apple-trees in California.
- 2080. [RILEY, C. V.] One effect of the Mississippi floods. <Amer. Nat., June [20 May], 1882, v. 16, pp. 514, 515.
 - Probability that *Ligyrus rugiceps* will have been drowned out of the region submerged for months by the Mississippi River; liability of the importation of new foes with new seed-plants.
- 2081. [RILEY, C. V.] Doryphora decemlineata in England. < Amer. Nat., June [20 May], 1882, v. 16, p. 515.
 - A living specimen carried from North America to London, England, in a barrel of potatoes.
- 2082. [RILEY, C. V.] Dr. Dimmock's inaugural dissertation. < Amer. Nat., June [20 May], 1882, v. 16, p. 515.
 - Review of G. Dimmock's paper on "The anatomy of the mouth-parts and of the sucking apparatus of some Diptera."
- 2083. [RILEY, C. V.] The triungulin of *Meloidæ*. <Amer. Nat., June [20 May], 1882, v. 16, p. 515.
 - The triungulin of Meloe proscarabæus was described by J. L. Frisch in 1727.
- 2084. [RILEY, C. V.] Fossil tineids. <Amer. Nat., June [20 May], 1882, v. 16, p. 515.
 - Notice of paper of V. T. Chambers; occurrence of the supposed burrow of a tineid larva in a fossil leaf of Acer sp.
- 2085. [RILEY, C. V.] Classification of North American Coleoptera. Amer. Nat., June [20 May], 1882, v. 16, pp. 515-516.
 - Notice of the approaching completion of the classification of the Coleoptera of North America by J. L. Leconte and G. H. Horn.

- 2086. RILEY, C. V. Hibernation of the army-worm. < Amer. Nat., June [20 May], 1882, v. 16, p. 516.
 - Confirmation of views in regard to the hibernation of the larvæ of Leucania unipuncta; places of oviposition; prospects of extensive injury by these insects in the more northern States in 1882.
- 2087. RILEY, C. V. Chinch-bug and army-worm prospects. <Rural New Yorker, 27 May, 1882. S.-b. No. 61, p. 27.
 - Blissus loucoptorus abundant in dry weather; Leucania unipuncta in wet weather; the latter hibernates principally as a larva; its eggs secreted in old grass and stubble; means against L. unipuncta.
- 2088. RILEY, C. V. Successful management of the insects most destructive to the orange. <Sci. Amer., 27 May, 1882 [v. 60], n. s., v. 46, pp. 335-336, 5 figs.
 - Treats of Cocoidæ injurious to the orange; means against the same; use of kerosene emulsion; figures stages of Mytilaspis pomicorticis [=pomorum], M. gloveri, and M. citricola.
- 2089. RILEY, O. V. The utilization of ants in horticulture. <Nature, 8 June, 1882, v. 26, p. 126. Reprint: <Gardener's Chronicle, 17 June, 1882, v. 17, p. 805. S. b. No. 42, p. 32.
 - Abstract of paper of C. J. Macgowan; capture and sale of two species of ants which build nests in trees; colonization of these ants in orange orchards to destroy injurious insects.
- 2090. RILEY, C. V. The army-worm vs. the clover hay-worm. Remedies for the army-worm. <Rural New Yorker, 10 June, 1882.
 S.-b. No. 47, pp. 158-159; No. 61, pp. 32-33. Reprint: <Nat. Farmer, 22 June, 1882.
 S.-b. No. 37, p. 123. <Lancaster Farmer, July, 1882.
 S.-b. No. 46, p. 104. <Home and Farm, 1 July, 1882.
 S.-b. No. 37, p. 122.
 - Abundance of Leucania unipuncta and Asopia costalis in Alabama and Tennessee; confusion of the army-worm with the clover hay-worm; prevalence of enemies of and means against the former.
- 2091. RILEY, C. V. Repelling insects by malodorants. < Amer. Nat., July [22 June], 1882, v. 16, p. 596.
 - Critical review of paper of J. A. Lintner; odorous substances repel insects more by their toxic properties than by their odor; failure of attempts by the use of strongly smelling substances to prevent oviposition; sight, touch, and taste generally more important in insect economy than smell.
- 2092. [RILEY, C. V.] Habits of Bittacus apterus. <Amer. Nat., July [22 June], 1882, v. 16, pp. 596-597.
 - Notice of paper of C. R. Osten Sacken; locomotion and food-habits of Bittacus apterus in California.
- 2093. RILEY, C. V. Habits of Coscinoptera dominicana. <Amer. Nat. July [22 June], 1882, v. 16, p. 598.
 - Discovery by F. H. King that Coscinoptera dominicana is inquilinous in ants' nests in its earlier stages; similar habits of related species.
- 2094. [RILEY, C. V.] Sun-spots and insect life. <Amer. Nat., July [22 June], 1882, v. 16, pp. 598-599.
 - Review of paper of A. H. Swinton; dependence of certain insect phenomena on the periods of sun-spots, through the determination by the latter of meteorological conditions.

- 2095. RILEY, C. V. The silk-worm. No. 1-3. < National Farmer, 13, 20, 27 July, 1882. S.-b. No. 35, p. 15; No. 48, pp. 32, 49.
 - Nature of the silk-worm; stages, enemies, diseases, races; wintering and hatching of eggs; feeding and rearing of larves; the coccess; spinning and gathering, choking chrysalids; egg-laying, reproduction, reeling, foodplants.
- 2096. RILEY, C. V. Change of habit; two new enemies of the eggplant. <Amer. Nat., August [28 July], 1882, v. 16, pp. 678-679.
 - Sudden acquisition by Doryphora juncta and by Cassida texana of the habits of feeding on Solanum melengena, these species having been found previously on S. carolinense and S. elwagnifolium respectively; occurrence of C. texana on S. carolinense; distribution of the two insects.
- 2097. BILEY, C. V. Notes on Microgasters. < Amer. Nat., August [28 July], 1882, v. 16, pp. 679-680.</p>
 Critical review of paper of A. S. Packard; synomymical notes on Packard's species.
- 2098. [RILEY, C. V.] Are honey-bees carnivorous? < Amer. Nat., August [28 July], 1882, v. 16, p. 681.

 Observations of Fritz Müller on the carnivorous habits of Brazilian Apide.
- *2099. [RILEY, C. V.] The "overflow bugs" in California. <Amer. Nat., August [28 July], 1882, v. 16, pp. 681-682.

 Extract from letter of Mrs. A. E. Bush, with introductory remarks; cosurrence of Platynus maculicollis in such abundance as to become a nuisance.
- 2100. [BILEY, C. V.] Insects and drouth. <Amer. Nat., September [24 August], 1882, v. 16, p. 745.
 - Occurrence of excessive drouth in the spring and early summer of 1880 in New England; statement by S. Lockwood of the exceptional abundance of several insects in New Jersey in that year.
- 2101. RILEY, C. V. Probable sound organs in sphingid pupes. < Amer. Nat., September [24 August], 1882, v. 16, pp. 745-746.

 Occurrence of a peculiar structure on the abdominal joints of the pupe of
 - ccurrence of a peculiar structure on the abdominal joints of the pupe of certain Sphingidæ; genera in which this structure is observed; probable connection of this structure with the function of producing sound as observed in Sphinz atropos.
- 2102. RILEY, C. V. Is Cyrtoneura a parasite or a scavenger? Amer. Nat., September [24 August], 1882, v. 16, pp. 746-747.

 Cyrtoneura stabulans bred from pups of Alctia ergillacea [= zylina] usual food of this species; doubt whether the species is a parasite or is only a scavenger in decayed pups; Phora alctia merely a scavenger; great liability of pups of Alctia to decay
- 2103. [RILEY, C. V.] Habits of Polycaon confertus Lec. < Amer. Nat., September [24 August], 1882, v. 16, p. 747.
 - Polycaon confertus bores in twigs of apple- and pear-trees and grape-vines; the larvæ probably live in the dead and dry wood of forest-trees.
- 2104. RILEY, C. V. Dinoderus pusillus as a museum pest. < Amer. Nat., September [24 August], 1882, v. 16, p. 747.
 - Dinoderus pusillus feeds on cork and paper lining in an insect box; they occur usually in drugs and other stored and dry vegetal products.

- 2105. RILEY, C. V. Myrmecophilous Coleoptera. <Amer. Nat., September [24 August], 1882, v. 16, pp. 747-748.
 - Larvæ and images of Euphoria hirtipes live in hills of Formica rufa; pupæ of Hymenorus rufpes in nests of Formica fusca and of H. obscurus in nests of another species of ant; known myrmecophilous habits of Cetonia, Cremastochilus, Euparia castanea (in nests of Solenopsis xyloni [= geminatu]), Tenebrionidæ, and Anthicus.
- 2106. [RILEY, C. V.] Discontinuance of publication. <Amer. Nat., September [24 August], 1882, v. 16, p. 748.

 Announcement of the discontinuance of publication of "Revue Coléoptérologique;" critical review of the same.
- 2107. RILEY, O. V. Buffalo tree-hopper injurious to potatoes. < Amer. Nat., October [28 September], 1882, v. 15, p. 823. Habits, food-plants, and ravages of Ceresa bubalus.
- 2108. [RILEY, C. V.] Moths attracted by falling water. < Amer.
 Nat., October [28 September], 1882, v. 16, p. 826.
 Notice of paper of J. S. Gardner; gleaming water-falls in Iceland as attractive to moths as artificial light would be.
- 2109. [RILEY, C. V.] A new museum pest. <Amer. Nat., October [28 September], 1882, v. 16, p. 826. Reprint: <Psyche, September-October, 1882 [1 March, 1884], v. 3, p. 408.

 Perimegatoma variegatum as a pest in collections of insects.
- 2110. [RILEY, C. V.] Fleas feeding on lepidopterous larvæ. <Amer. Nat., October [28 September], 1882, v. 16, p. 826.</p>
 Notice of paper of C. J. Boden; abundance of fleas (Pulex sp.) where few or no warm-blooded animals occur may be explained by the feeding of the fleas on insects.
- 2111. [RILEY, C. V.] [Gasteracantha cancer.] < Gonzales [Tex.] Inquirer, 30 September, 1882. S.-b. No. 42, p. 15.

 Habits of Gasteracantha cancer; the male unknown.
- 2112. RILEY, C. V. Oicada septendecim. < Gardener's Mo. and Hortic., September, 1882, v. 24, pp. 274-275. S. b. No. 38, p. 42; No. 39, p. 6; No. 42, pp. 2-4.
 - Orthography of the names C. [= Tibicen] septendecim and C. [= T.] tredecim; natural relations of the same; indistinguishability of the species of certain genera; dimorphic forms more numerous than usually recognized; Massospora cicadina parasitic on Cicada.
- 2113. RILEY, C. V. Remarkable felting caused by a beetle. <Rural New-Yorker, 14 October, 1882, v. 41, pp. 699-700. S.-b. No. 42, p. 16. Reprint, with changes: <Amer. Nat. [2] December, 1882, v. 16, pp. 1018-1019.

 Perception of the felting of the interior of a pillow-ticking with fragments.
 - Description of the felting of the interior of a pillow-ticking with fragments of feathers formed by the ravages of Attagenus megatoma within a feather pillow.
- 2114. RILEY, C. V. The buckeye-leaf stem-borer. <Amer. Nat., November [28 October], 1882, v. 16, pp. 913-914. See: <Sci. Amer. Suppl., 16 December, 1882. S. b. No. 47, p. 145.
 - Sericoris instrutana †Claypole = S. [= Steganoptycha] claypoleana n. sp.; habits and food-plants of this species and of Protectoras asculana; distinctions between these two species.

- 2115. [RILEY, C. V.] Efficacy of chalcid egg-parasites. < Amer. Nat., November [28 October], 1882, v. 16, pp. 914-915. See: < Prairie Farmer, 2 December, 1882. S.-b. No. 60, p. 123.
 - Beneficial influence and occasional vast abundance of egg-parasitic Chalcidida and especially of Trichogramma pretiosa and of Telenomus sp.
- 2116. [RILEY, C. V.] On the biology of Gonatopus pilosus Thoms. <Amer. Nat., November [28 October], 1882, v. 16, p. 915.
 - Notice of paper of J. Mik; parasitism of Gonatopus pedestris on Athysanus maritima, and of G. pilosus on Deltocephalus xanthonourus; transformations and habits of G. pilosus; probable parasitism of a Gonatopus (G. contortulus?) on Amphiscepa bivittata.
- 2117. RILEY, C. V. Species of *Otiorhynchidæ* injurious to cultivated plants. <Amer. Nat., November [28 October], 1882, v. 16, pp. 915-916. See: <Sci. Amer. Suppl., 16 December, 1882. S.-b. No. 47, p. 145.
 - Aramigus fulleri the only species of North American Otiorhynchides whose development and earlier stages are known; food-plants of several species; discovery by G.P. Peffer of the injuries of Anametic grises upon roots of apple- and pear-trees.
- 2118. [RILEY, C. V.] Bombyliid larvæ destroying locust eggs in Asia Minor. <Amer. Nat., November [28 October], 1882, v. 16, pp. 916-917.
 - Notice of and extract from communications of F. Calvert; parasitic habits of Callostoma fascipennis; similarity of habits of Bombyliida in Asia and in North America; probability that the larvae of Cantharis vesicatoria and of other Meloidae will be found to feed on eggs of Acrididae.

1

2119. RILEY, C. V. Report of the Entomologist. <Ann. Rept. [U. 8.]

Commissioner Agric. for 1881 and 1882, 1882 [January, 1883],

pp. 61-214, 20 pl. Separate: <Washington, 1 December, 1882,

pp. 8+167, 20 pl.

CONTENTS.

Plan of the report, 1—Insufficiency of former appropriations to the Entomological Division, 1—Reorganization of the Division, 1—Great increase in the correspondence of the Division, 2—Preparation of special bulletins, 2—Periodical bulletins desirable, 2—Special reports in preparation, 2—Work of the United States Entomological Commission, 3—The United States National Museum in connection with the Entomological Division, 3—Assistants, agents, and observers of the Division, 3—The illustrations to the

EXTRACTS FROM CORRESPONDENCE

Miscellaneous correspondence on various insects injurious to vegetation in 1881-'82, 4.

Distribution of eggs by the Department, 7—Mr. L. S. Crozier's opinion as to the value of American silk products, 7—Mr. E. Fasnach on the importance of a home market for silk products, 8—Associations, 8—Ladies' Association at Spring Hill, Ala., 8—Women's Silk Culture Association of California, 8—Women's Silk Culture Association of the United States, 8—American Silk Exchange in New York, 8—Sales of eggs and coopons in the

16

27

29

46

2119. RILEY, C. V.—Continued.

SILK CULTURE-Continued.

United States, 9—Various reports from correspondents of the division, 9—Experience in 1882 at the Department in raising worms fed on osage orange or mulberry, 11—New mulberry tree from southern Russia, 13—Business ventures in the United States in connection with silk-producing industry, 13—Sale of eggs, 13—Second edition of the manual of silk culture, 14—Preface thereof, 14—Reasons why the growing interest in silk culture in the United States should be checked at the present time, 15—Import duty on raw silk demanded, 16.

Washington, 24—Miss M. E. Murtfeldt's experiments with powder on various insects, 25.

CHINCH-BUG NOTES, Blissus leucopterus

Professor Thomas' predictions for 1881, 27—1881 a chinch-bug year, 27—Injury in spring of 1882 in the western States, 27—Irriga-

tion as remedy, 28-Use of kerosene emulsions, 29.

in fumes, 20—Alcoholic extract, 20—The powder stirred up in water, 20—Tea or decoction, 21—Reports from persons to whom Pyrethrum seed was sent, 21—Success in raising the plant at

THE ARMY WORM, Leucania unipuncta..... Habits and natural history, 30-History of investigation in America, 30-Where the eggs are laid, 30-When the eggs are laid, 31-Mode of oviposition, 31-Prolificacy of the moth, 32-Duration of the egg state, 32-Habits of the young worms, 32-Guenée's characterization of Leucania larvæ, 32-Marching of the worms abnormal, 33-The marching worm not a distinct race, 33-Duration of worm life, 34-Remedies, 34-Burning old grass, 34-Predictions of army-worm years unreliable, 35-Meteorological influences on the species, 35-Ditching and fencing as remedies, 36-Use of arsenical poisons, 36-Kerosene and coal tar, 36-Heavy rolling not to be recommended, 37-"Drawing the rope" of little avail against the worm, 37-Mr. L. O. Howard's report on army-worm injury, 1881, in Illinois and Indiana, 37-Reports from correspondents, spring of 1882, 39-Rev. Samuel Lockwood's account of the invasion of 1880, in New Jersey, 44.

2119. RILEY, C. V.—Continued.

SCALE INSECTS OF THE ORANGE-Continued. enemies of scale insects, 49—Hymenopterous parasites the most efficient destroyers, 49—Sudden outbursts of the pest, 50—Destruction of the scales by their own undue increase, 51-Injury inflicted on young trees, and faulty methods of destroying the scales, 51—Recapitulation, 51—Remedies, 52—Danger in applying undiluted volatile oils, 52-Kerosene, when properly applied, not injurious to the tree, 52 - Process of preparing kerosene butter, 52-Nature of kerosene butter, 53-How it should be applied, 54-The kerosene butter a sure remedy, 54-How it affects the scales and the eggs, 54 - The aquapult force-pump very convenient for applying kerosene emulsions, 54-Effect of kerosene emulsion on the tree, 55—Kerosene emulsions experimented with, 55-Whale oil soap as remedy, 56-Oil of creosote, its effect on scale insects and on the trees, 56-Saponaceous compounds of creosote, 57-Creosote inferior to kerosene as insecticide, 57-Various other substances as remedies, 57-Tables of experiments, 60-Note on more recent discoveries in producing kerosene emulsions, by C. V. Riley, 67.

> 67 68

70

73

76

78

78 78

INSECTS AFFECTING THE RICE-PLANT

Rice production in the United States in 1879, 67.
The rice-grub, Chalepus trachypygus
Mr. L. O. Howard's observations on the occurrence and habits of
the insect, 68—It can easily be kept in check, 68—Structural char-
acters and geographical distribution of the genus Chalepus, 69—
Description of imago and larva, 69.
The water-weevil, Lissorhoptrus simplex
Colonel Screven on the specific identity of "the maggot" and "the
water-weevil," 70-Mr. L. O. Howard's report on the occurrence
of larva and beetle in rice-fields, 70—Habits of the beetle, 71—
Remarkable shape of the larva, 71—Characters of the genus Lis-
sorhoptrus, 72—Description of the image, 72—Description of the
larva, 72.
The rice-stalk borer, Chilo oryzæellus
Mr. Howard's account of the work of the larva and the damage
caused by it, 73—Enemies, 74—Preventive measures, 74—Difficul-
ties in the specific determination of the insect, 74—Characteris-
tics of the genus Chilo, 75—Description of the moth, 75—Descrip-
tion of larva and pupa, 75.
White blast
Colonel Screven's account of the disease, 76—Mr. Howard's observa-
tions on the insects found on the diseased plants, 77—The blast
possibly caused by insect work, 77.
Other insects injurious to growing rice
The "grass-worm" and its injury, 78—The "lubber grasshopper,"
78—Acridium obscurum and various Heteroptera, 78.
INSECTS AFFECTING CORN OR MAIZE
The corn-bill bug, Sphenophorus robustus
Species of Sphenophorous injurious to agriculture, 78-Nature of
damage done by the beetles, 79—Mr. Howard's report on the work
of the larva and its development, 79—Preventive measures, 80—
Burning the stubble in winter time to be recommended, 80—Gen-
eric characters of Sphenophorus and its natural groups, 80—Diag-
nosis of the imago, 81—Description of the larva, 81—Larva of

2119.	RILEY, C. V.—Continued.	
	INSECTS AFFECTING CORN OR MAIZE—Continued.	
	Rhodobænus 13-punctatus and Rhynchophorus simmermanni, 82— Description of the pupa, 82.	
	The smaller corn-stalk borer, Pempelia lignosella	82
	First appearance of the pest, 82—Geographical distribution in the	
	United States, 83—Habits and natural history, 83—Work of the	
	larva, 83—Transformation, 83—Habits of the moth, 83—Prevent	
	ive measures, 84—Color variations of the moth, 84—Mouth parts	
	of the moth, 84—Description of the larva, 84—Description of the	
	pupa, 85. The hell worm elies corn worm. Heliethic sumissing	85
	The boll-worm, alias corn-worm, Heliothis armigera	00
	worm and boll-worm, 85—Number of broods in the northern	
	States, 56—Nature of damage done to corn, 86—The worm also	
	feeding on hard corn, 87—Corn-feeding broods in the southern	
	States, 87-Damage done to tomatoes, 88-Food-plants of the	
	families Solanacem and Leguminosm, 88—Cucurbitaceous and	
	malvaceous food-plants, 99—Various other food-plants, 89—List	
	of food-plants probably still incomplete, 89—Carnivorous habit	
	of the worm, 89—Report upon Heliothie armigera, by Judge Law-	
	rence Johnson, 90.	00
•	THE COTTON-WORM, Aletia xylina	92
	Past work on the practical side of the cotton-worm question, 93—	
	Obstacles and difficulties of the investigation, 93—Natural his-	
	tory, 94—The insect in its different stages, 94—Time of develop-	
	ment and first appearance of the worms, 94—Poisons best applied	
	from beneath, 95-Improvements in machinery for applying poi-	
	sons, 95-Nozzles of various forms, 96-Machinery for poisoning	
	from below, 96.	
	Protection from injury in regions overflowed by the Mississippi	97
	Poisons should be applied to the under surface of the leaves, 98—	
	Paris green and London purple, 96—Sprinkling and sifting, 98—	
	Device for mixing poisons and diluents, 98—Arsenic and arseniate	
	of soda, 98—Early poisoning, 99. Poisoning devices—machine for spraying from below	99
	The skid, 99—Device for thorough mixing and filling the barrel,	33
	100—Details of barrel, pump, and stirrer, 100—Inserting and ex-	
	tricating the stirrer-bar, 100—Fork of the descending pipe, 100—	•
	The nozzles, 102.	
	Damage in 1881	102
	Loss of cotton by worms by States and counties, 102 - Loss of	
	cotton by worms in 1881, tabulated, 104.	
	Possible food plants of the cotton-worm	105
	No other food-plant than cotton yet found, 104—Eggs of Anomie	
	erosa found on Urena lobata, 104—Eggs and young larvæ of Anomis	
	erosa distinguished from those of Aletia xylina, 105—Plants upon which eggs related to Aletia were found in the Department her-	
	barium, 105—Localities for malvaceous plants, with possible	
	food-plants of Aletia indicated, 105.	
	The question of hibernation settled	106
	Theories of hibernation, 106—Eggs deposited early in March, 106—	
	The early brood of worms, 106—Where the moth hibernates,	
	106-Value of fall and winter preventive work, 106.	

2119.

RILEY, C. V.—Continued.	
MISCELLANEOUS INSECTS	107
The Urena anomis, Anomis crosa	107
Habits and natural history; resemblance to the cotton-worm, 107—Distribution, 107—Its eggs mistaken for those of Aletia, 108—Distinguishing characters, 108—Habits of larvæ, 108—Characters of moth, 109—Descriptive, 109.	
The clover leaf-beetle, Phytonomus punctatus	111
Habits of the genus Phytonomus, 111—Ph. punctatus an European insect, 111—Food-plants of European species, 111—History of Ph. punctatus in N. A., 111—Specific identity of Ph. punctatus and Ph. opimus, 112—When first noticed as destructive, 112—Damage in Yates County, N. Y., in 1881, 112—Mr. Schwarz's observations in	
June, 1882, 113—Life history of the species, 113—The eggs, 113—The newly-hatched larvæ, 113—Method of locomotion of larvæ, 114—The cocoon, 114—Method of spinning, 115—Its method of spinning differs from that alleged of the European Ph. rumicis, 115—Principal damage done in the beetle state, 115—Feeds on	•
white and red clover, 115—Length of life in the different states, 115—Number of annual broods, 116—A second brood exceptional, 116—Remedies, 117—Will it spread?, 117—Natural enemies, 117—Description of earlier states, 118.	
The vagabond Crambus, Crambus vulgivagellus History of its injury and identification, 119—Correspondence with Professor Lintner, 119—Abundance of the moths in the eastern States in 1881, 119—Habits and natural history, 119—The eggs, 120—The larva and its tube, 120—Habits of the moth, 120—Damage done by the worms, 120—Parasites, 121—Similarity of habit in an European species, 121—Remedies, 121—Descriptive, 121— Specimens from Vancouver's Island compared, 121—Descriptions of earlier states, 122—Bibliographical list, 122.	119
The wheat Isosoma, Isosoma tritici	123
Past history and habits, 123—When first studied, 123—Professor Thomas finds it in Illinois and Dr. Packard in Virginia, 123—Its work differs from that of the joint-worm, 123—Its injuries in Missouri, 124—Professor French's observations, 124—Comparisons with the joint-worm and other allied species, 124—Errors corrected, 125—Its relationship to the European I. lineare; habits of latter, 125—Number of broods, 125—Remedies, 125—Parasites, 126—Descriptive, 126—Bibliographical list, 126.	
The sorghum web-worm, Nola sorghiella	127
The catalpa sphinx, Ceratomia catalpa	129
Value of the catalpa tree, 129—Its usual exemption from insect attacks, 129—Past history of Sphinz catalpa, 129—Its larvæ used as fish bait, 130—Its distribution, 130—Extracts from correspondence concerning the larva, 130—Characters and natural history: The eggs, 131—The larvæ, 131—The pupa, 131—Description of the moth, 131—Number of broods, 132—Remedies, 132—Descriptive, 132—Peculiarity of structure of pupa, 133.	
The osage orange sphinx, Coratomia kagemi	133
Value of the osage orange, 133—Its comparative exemption from insect attack, 133—Rarity of the osage orange sphinx, 133—Its generic place, 133—Characters of the species, 134—Its affinities, 134—Description of larva, 134.	

2119.	RILEY, C. V.—Continued.	
	REPORT ON MISCELLANEOUS INSECTS, BY PROF. J. HENRY COMSTOCK.	135
	The apple-maggot, Trypeta pomonella. Its importance compared with the codlin moth, 135—Method of work, 135—Food-plants, 135—Its spread, 136—Extracts from correspondence, 136—Early apples principally infested, 136—Technical descriptions of different states, 137—Remedies, 138—Manner of distinguishing it from codlin-moth larva, 138.	135
	The vine-loving pomace-fly, <i>Drosophila ampelophila</i> Reasons for popular name, 139—Reasons for treating of the species, 139—Where found, 139—Rapidity of multiplication, 139—Destructive to grapes, 139—Technical descriptions of different stages, 140—Remedies, 141.	138
	The pretty pomace-fly, Drosophila amana	141
	Technical descriptions of different stages, 141—Remedies, 142.	
	The occilate leaf-gall of the red-maple, Sciara occilaris	142
	Ladybirds, Coccinellidæ General remarks on habits, 144—The ashy-gray ladybird, 144—Description of larva, 144—Description of pupa, 145—The adult, 145—The blood-red ladybird, 145—Description of pupa, 145—The ladybird of the cactus, 145—Descriptions of larva and pupa, 145—The ambiguous Hippodamia, 146—Descriptions of larva and pupa, 146—Other species observed at Los Angeles, Cal., 146.	144
	Methods of destroying scale insects	146
	Lac insects, Carteria. General remarks, 149—The genus Carteria of Signoret, 49—Carteria lacca (Kerr); general appearance, 149—Preparation for study, 150—Technical description, 150—Remarks on Mr. Carter's description, 151—Carteria larreæ n. sp., 151—Remarks on the crossote plant, 151—Mr. J. M. Stillman's paper, 151—Differs from C. lacca, 151—Technical description of C. larreæ, 152—Carteria mexicana n. sp., 152—Where found, 152—General appearance, 152—Technical description, 152.	149
	A new wax insect, Cerococcus querous	153
	Note on the structure of mealy bugs	154
2120.	[RILEY, C.V.] A new rice stalk-borer: Genus-grinding. <ar Nat., [2] December, 1882, v. 16, pp. 1014-1015.</ar 	
	Extract from Rept. [U. S.] Commis. Agric. for 1881 and 1882, pp. 134-with additional remarks; Chilo oryzæellus — Diphryz prolatella; Dip Grote founded on a mutilated specimen, with mistake of maxillary labial palpi.	kryz

- 2121. [RILEY, C.V.] A butterfly larva injurious to pine-trees. < Amer. Nat., [2] December, 1882, v. 16, pp. 1015-1016.
 - Habits and ravages of *Pieris menapia* on yellow-pine and tamarack in Washington Territory.
- 2122. [RILEY, C. V.] The army-worm in 1882. <Amer. Nat., [2] December, 1882, v. 16, p. 1017.
 - Occurrence of Leucania unipuncia in great abundance and with disastrons effects, especially in southern United States in 1882.
- 2123. [RILEY, C. V.] The wheat-stalk worm on the Pacific coast. <Amer. Nat., [2] December, 1882, v. 16, pp. 1017–1018.
 - Isosoma tritici injuring wheat-stalks in Washington Territory; reference to prior notices of this insect.
- 2124. [RILEY, C. V.] Deserved honor. < Amer. Nat., [2] December, 1882, v. 16, p. 1018.
 - Notice of the appointment of Eleanor, A. Ormerod to be consulting entomologist to the Royal Agricultural Society of Great Britain.
- 2125. [RILEY, C. V.] Important work on *Cynipidæ*. <Amer. Nat., [2] December, 1882, v. 16, p. 1018.

 Notice of G. Mayr's "Die europäischen Arten der gallenbewohnenden Cyni-
- piden."

 2126. RILEY, C. V. Emulsions of petroleum and their value as insecticides. <Rural New Yorker, 9 December, 1882, v. 41, pp. 833, 834. S.-b. No. 42, pp. 17-18.
 - Results of experiments made upon methods of using petroleum as an insecticide without injury to plants; soap and milk emulsions the most available; methods of preparing the same.
- 2127. RILEY, C. V. The bean-weevil. <Rural New Yorker, 9 December, 1882, v. 41, p. 835. S.-b. No. 42, p. 18.

 Answer to inquiry of R. J. B.; habits of and means against Bruchus faba.
- 2128. BILEY, C. V. The "cluster-fly." < Prairie Farmer, 23 December, 1882, v. 54, p. 7. S.-b. No. 42, pp. 16-17. Reprint: < Amer. Nat., [5] January, 1883, v. 17, pp. 82-83.
 - Habits and synonymy of *Pollenia rudis*; notice of other accounts of the swarming of Diptera. See No. 2174.
 - 2129. RILEY, C. V. Darwin's work in entomology. <Proc. Biol. Soc. Wash., 1882, v. 1, pp. 70-80.
 - Analysis of the interest shown by C. R. Darwin in entomology and of his contributions to the same.
 - 2130. RILEY, C. V. The cotton-worm. < Western Farmer's Almanae for 1883, 1882, p. 40. S.-b. No. 42, p. 71.
 - Various theories hitherto held in regard to the hibernation of Aletia argillaces [=xylina]; proof of its hibernation; seasons and conditions of development of the first brood of the year, precautionary measures to be adopted.
 - 2131. RILEY, C. V. Pyretbrum, an important insecticide. < Western Farmer's Almanac for 1883, 1882, pp. 41-42. S.-b. No. 42, p. 49. See: < Prairie Farmer, 27 January, 1883. S.-b. No. 42, pp. 56-58. Condensed account of the history of the use of pyrethrum flowers as an insecticide; cultivation of the plants and preparation of the powder; meth-

ods of its application; experiments in the cultivation of the plants.

- 2132. [BILEY, C. V.] New lists of North American Lepidoptera. <Amer. Nat., [5] January, 1883, v. 17, pp. 80-82.
 - Reviews of lists of Brooklyn Entomological Society, of C. H. Fernald and A. R. Grote.
- 2133. [RILEY, C. V.] Naphthaline cones. <Amer. Nat., [5] January, 1883, v. 17, pp. 83, 84.
 - The cones stain the paper lining of boxes; they seem to destroy mites and *Peoci* very soon, but have little effect on *Dermestida*.
- 2134. RILEY, C. V. Emulsions of petroleum as insecticides. <Sci. Amer., 6 January, 1883 [v. 62], n. s., v. 48, p. 3. S.-b., No. 42, pp. 4-7.
 - Notice of experiments made, under author's direction, in 1882, in the use of emulsions of kerosene oil; report of H. G. Hubbard upon experiments made by him; critical review of S. F. Chapin's "Scale insects on deciduous and ornamental trees;" effect of pure kerosene, of emulsions, and of lye upon trees.
- 2135. RILEY, C. V. Entomological notes. <Rural New Yorker, 13 January, 1883. S.-b. No. 42, p. 78. See: <Amer. Nat., 1883, v. 17, pp. 198-199.
 - 1. A new enemy to wax-beans; extract from letter of G. H. Stone, on the food-plants, habits, and ravages of *Epilachna corrupta*; distribution of the same. 2. Spread of the 12-punctured asparagus beetle; increasing noxiousness of *Crioceris* 12-punctata recorded from near Baltimore, Md., by O. Lugger; comparative description of this species with *C. asparagi*. 3. An internal mite in fowls; presence of *Cytoleichus sarcoptoides* in lungs and other parts of diseased chickens.
- 2136. [RILEY, C. V.] The "lignified snake of Brazil." <Evening Star [Washington, D. C.], 20 January, 1883, v. 61, p. 2. S.-b. No. 42, pp. 59-60. Reprint: <Sci. Amer. Suppl., 17 February, 1883. See: <Science, 23 February, 1883, v. 1, p. 84.
 - Discussion of a specimen of problematical character, supposed to be the burrow of a larva under bark; notice of writings upon the subject; frequency with which the true nature of natural objects is mistaken; letter from J. H. Hutchins accompanying a gall of Cecidomyia vitis-pomum mistaken for a hybrid fruit.
- 2137. RILEY, C. V. Utilization of ants in horticulture. <Sci. Amer., 27 January, 1883 [v. 62], n. s., v. 48, p. 49. S.-b., No. 42, pp. 65-66.
 - Abstracts of papers of C. J. Macgowan and H. C. McCook, with additional notes; the introduction of ants might involve objectionable consequences; probability that they would not be of service against Cocoida.
- 2138. RILEY, C. V. Natural sugaring. <Amer. Nat., February [31 January], 1883, v. 17, pp. 197-198. Reprint: <Country Gentl., 31 May, 1883, v. 48. S. b. No. 42a, p. 297.
 - Lachnus platanicala n. sp. [p. 198], abundant in 1882 on sycamore trees; description of the species; attraction of great numbers of insects to its saccharine exudations, and growth of Funago salicina upon these exudations.
- 2139. [RILEY, C. V.] Trogoderma tarsale as a museum pest. < Amer. Nat., February [31 January], 1883, v. 17, p. 199.
 - Notice of paper of F. H. Snow; remarks on the abundance and ravages of Trogoderma tarsale; habits of its larva.

2140. [RILEY, C. V.] Phylloxera in California. < Amer. Nat., February [31 January], 1883, v. 17, pp. 199-200.

Phyllowera vastatrix in California is most injurious in moist soils.

- 2141. RILEY, C. V. The hibernation of Aletia xylina [Say] in the United States a settled fact. <Sci. Amer., 3 February, 1883, v. 48, p. 68. S.-b. No. 42, pp. 66-67. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1882, 1883, v. 31, pp. 468-469. Separate: <Salem, Mass., July, 1883, pp. 468-469. Abstract: <Amer. Nat., April [15 March], 1883, v. 17, pp. 420-421. <Nature, 28 December, 1882, v. 27, p. 214.
 - Proof of the hibernation of Aletia xylina as a moth and of the perpetual existence of the species in Florida.
- 2142. [RILEY, C. V.] Fostering the study of economic entomology.

 <Rural New-Yorker, 12 February, 1883. S.-b. No. 42, p. 58.

 Reprint: <Amer. Nat., April^o[15 March], 1883, v. 17, p. 420.

 Efforts of the French for the promotion of economic entomology.
- 2143. [RILEY, C. V.] Diseases of the chinch-bug. < Rural New-Yorker, 17 February, 1883. S.-b. No. 42, p. 58.
 - Note concerning paper of S. A. Forbes on two fungus parasites of Blissus leucopterus.
- 2144. R[ILEY], C. V. [Instinct of Cicada [= Tibicen] septendecim.]

 Amer. Nat., March [21 February], 1883, v. 17, p. 322.

 Remarks on note of E. W. Claypole; sense of direction in insects.
- 2145. [RILEY, O. V.] Food-habits of Megilla maculata. <Amer. Nat.,</p>
 . March [21 February], 1883, v. 17, pp. 322-323.
 Summary of S. A. Forbes' observations upon the food of Megilla maculate;
 results of other observations on this subject.
- 2146. [RILEY, C. V.] Clothes-moths observed in the United States. <Amer. Nat., March [21 February], 1883, v. 17, p. 323.

 Notice of paper by C. H. Fernald.
- 2147. RILEY, U. V. Entomological notes. <Rural New Yorker, 27 February, 1883. S.-b. No. 42, p. 58. See: <Amer. Nat., April [15 March], 1883, v. 17, pp. 419-420.
 - Abstract of the rules of the international convention at Berne for the prevention of *Phylloxera* ravages; their adoption by Belgium.
- 2148. RILEY, C. V. Notice of an illustrated essay on the *Noctuidæ* of North America. <Bull. Brooklyn Ent. Soc., February, 1883, v. 5, pp. 77-79. Separate: <[Brooklyn, N. Y., 1883], 4 pp. Critical review of A. R. Grote's essay.
- 2149. RILEY, C. V. Concerning canker-worms. <Indiana Farmer, 3
 March, 1883. S.-b. No. 56, p. 69. <Prairie Farmer, 3 March,
 1883. S.-b. No. 56, p. 69. <Pacific Rural Press, 10 March,
 1883. S.-b. No. 54, p. 15. <Lancaster Farmer, March, 1883.
 S.-b. No. 56, p. 72. <Gardener's Mo. and Hortic., April, 1883,
 v. 25. S.-b.No. 42a, p. 631.
 - Description of Paleacrita vernata and Anisopteryx pometaria; ravages of the former; request for information; method of observation.

· · · · ·

- 2150. RILEY, C. V. Dipterous enemies of the *Phylloxera vastatrix*. <Ca. Ent., February [9 March], 1883, v. 15, p. 39.
 - Critical review of paper of T. W. Fyles; the characters given of Diplots grassator are insufficient to distinguish the species; galls of Phylloxera vastatrix inhabited by Leucopis phylloxera Riley MS.; comparison of the early stages of these two Diptera.
- 2151. [RILEY, C. V.] The food relations of the Carabidæ and Coccinellidæ. <Amer. Nat., April [15 March], 1883, v. 17, pp. 417-419.

 Summary of S. A. Forbes's observations on the food relations of the Carabidæ and Coccinellidæ.
- 2152. [RILEY, C. V.] Relations of the Carabidæ and Coccinellidæ to birds. <Amer. Nat., April [15 March], 1883, v. 17, p. 419.

 Reprint of S. A. Forbes's summary, with note; Coccinellidæ not eaten by birds; Carabidæ eaten in proportion as they have phytophagous habits.
- 2153. [RILEY, C. V.] Viviparity in a moth. <Amer. Nat., April [15 March], 1883, v. 17, p. 420.

 Notes Fritz Müller's discovery of a moth seen to deposit living larvæ.
- 2154. [RILEY, C. V.] Damage to silver-plate by insects. <Amer. Nat., April [15 March], 1883, v. 17, p. 420.

 Holes in silver-plate said to have been made by Niptus hololeucus.
- 2155. RILEY, C. V. Possible food-plants of the cotton-worm. <Amer. Nat., April [15 March], 1883, v. 17, pp. 421-422.</p>
 Notice of paper of J. S. Bailey; occurrence of newly issued images of Aletis xylina at Karner, N. Y., 7 and 8 October, 1882, proving that the larva of this insect feeds upon some genus of plants other than Gossypium.
- 2156. [RILEY, C. V.] Agrotis messoria Harr. vs. Agrotis scandens Riley.

 Amer. Nat., April [15 March], 1883, v. 17, p. 422, 2 figs.
 Critical review of views of A. R. Grote; Agrotis lycarum, A. repentis, and A. cochranii = A. messoria; A. scandens is a distinct species; comparison of the imagos; figures larvæ and imagos of the two species.
- 2157. [RILEY, C. V.] An internal mite in fowls. <Amer. Nat., April [15 March], 1883, v. 17, pp. 422-423.</p>
 Lungs, bronchia, and linings of thoracic and abdominal cavities of a sick chicken covered with Cytoleichus sarcoptoides Mégnin; habitat of the same in fowls in Europe; diseases caused by it.
- 2158. [RILEY, C. V.] Prevalence of the screw-worm in Central America. <Amer. Nat., April [15 March], 18-3, v. 17, p. 423.

 Extract from letter of J. C. Zeledon on the abundance and ravages of Lucilia macellaria and related flies in Costa Rica.
- 2159. R[ILEY], C. V. Dried leaves as food for lepidopterous larvæ. Amer. Nat., April [15 March], 1883, v. 17, pp. 423-424.

 Review of paper of A. H. Mundt; larvæ successfully fed upon fresh leaves transported from a distance under pressure; chopping the leaves would possibly permit of more rapid curing and more convenient packing.
- 2160. [RILEY, C. V.] Lepidopterological notes. < Amer. Nat., April [15 March], 1883, v. 17, p. 424.</p>
 Duplication of descriptions of the early stages of Lepidopters to be avoided;

notice of Mrs. C. H. Fernald's list of Noctuida taken in Orono, Me.

2161. [RILEY, C. V.] Obituary. < Amer. Nat., April [15 March], 1883, v. 17, p. 424.

Obituary notices of G. W. Belfrage and F. W. Mæklin.

2162. R[ILEY], C. V. Mosquitões vs. malaria. Sci. Amer., 14 April, 1883, v. 48, pp. 224-225. S. b. No. 42, pp. 63-64; No. 67, p. 8. Abstract: < Amer. Nat., May [18 April], 1883, v. 17, p. 549.

Criticism of the views of Dr. A. F. A. King in support of the thesis that malarial disease is the result of inoculation of the body with malarial poison by the bites of insects; citation of twenty correspondencies in the conditions affecting the prevalence of mosquitoes [Culicidæ] and malarial disease.

2163. RILEY, C. V. Jumping seeds and galls. <Sci. Amer., 14 April, 1883, v. 48, p. 228, fig. S.-b. No. 42, pp. 61-63; No. 67, p. 2. Figures of larva, pupa, and imago of Carpocapea saltitans, with figures and

description of seeds inhabited by the larva of this moth, and description of the plant bearing these seeds; vernacular names of plant and insect; movements imparted to the seeds by the insects and by Cynips q.-sallatorius to the galls of the same.

2164. RILEY, C. V. Reports of experiments, chiefly with kerosene, upon the insects injuriously affecting the orange-tree and the cotton-plant, made under the direction of the entomologist. <Bull. No. 1, Div. Ent. U.S. Dept. Agric., [17 April], 1883, 62. pp.</p>

CONTENTS.

Letter of submittal	3
Introduction	5
Miscellaneous notes on orange insects. By H. G. Hubbard	9
Experiments upon scale insects affecting the orange. By Jos. Voyle	19
Report of observations and experiments. By J. C. Neal	31
Report of observations and experiments on the cotton-worm (Aletia xylina). By R. W. Jones	47
Report upon the cotton-worm, boll-worm, and other insects. By Law-rence Johnson	53
Index	59

2165. RILEY, C. V. Reports of observations on the Rocky Mountain locust and the chinch-bug, together with extracts from the correspondence of the division on miscellaneous insects. < Bull. No. 2, Div. Ent. U. S. Dept. Agric., [17 April], 1883, 36. pp. Second edition, 16 September, 1883.

CONTENTS.

Letter of submittal	
Introduction	
Report of observations in the Northwest on the Rocky Mountain locust,	
by Lawrence Bruner	
Experiments on chinch-bugs. By S. A. Forbes	:
Extracts from correspondence	:
Index	;

2166. [RILEY, C. V.] Insects as food for man. < Amer. Nat., May [18] April], 1883, v. 17, pp. 546-547.

Summary of Max Buchner's observations on the insects used as food by the Bantus tribe of negroes.

- 2167. RILEY, C. V. Number of molts and length of larval life as influenced by food. <Amer. Nat., May [18 April], 1883, v. 17, pp. 547-548.
 - Variability in habits and characters of insects; periods and number of molts observed in larvæ of *Tenebrio molitor*, *T. obscurus* and *Trogoderma tarsale*; conclusion that insufficient food retards development and occasions frequent molting.
- 2168. [RILEY, C. V.] Entomological notes. <Amer. Nat., May [18 April], 1883, v. 17, pp. 549-550.

 Brief notes on recent publications.
- 2169. RILEY, C. V. Larval stages and habits of the bee-fly *Hirmoneura*. <Science, 27 April, 1883, v. 1, pp. 332-334, flgs. 1-3. S.-b. No. 42, p. 52.
 - Summary of the life-history of *Hirmoneura obscura* condensed from Handlirsch, and from Brauer; figures the several stages; correspondences of the structure and early history of the larva; predictions in reference to the larvæ of *Bombyliidæ*.
- 2170. RILEY, C. V. The capitalizing of specific names. < Papilio, March [April], 1883, v. 3, p. 62.
 - Inquiry as to the use and purpose of capitalizing specific names. See No. 2257.
- 2171. RILEY, C. V. Observations on the fertilization of Yucca and on structural and anatomical peculiarities in Pronuba and Prodoxus. <Gardener's Mo. and Hortic. April, 1883, v. 25, pp. 118–119. S.-b. No. 51, p. 122½. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1882, 1883, v. 31, pp. 467–468. Separate: <Salem. Mass., July, 1883, pp. 467–463. Abstract: <Amer. Nat., February [31 January], 1883, v. 17, p. 197.
 - Description of the manner in which Pronuba guccasella gathers the pollen in flowers of Yucca; the work of this species necessary for the fertilization of the capsular species of Yucca, the irregularity of whose fruit is due to its punctures; description of the egg and of the manner of oviposition of this species.
- 2172. RILEY, C. V. The potato-stalk borer. <Rural New Yorker, 12
 May 1883. S.-b. No. 42a, p. 213.
 - Answer to inquiry; life-history and means against Baridius [= Trichobaris] trinotata.
- 2173. RILEY, C. V. Jumping seeds and galls. <Proc. U. S. Nat. Mus., 12 May, 1883, v. 5, pp. 632-635, fig. Separate: <[Washington, 1884], pp. 632-635, fig.
 - Description of seeds of "arrow-weed"; their motions caused by larve of Carpocapse saltitans; figures of the seeds and of the larva, pupa, and image of the Carpocapse; habits and seasons of the same; description of the plant bearing these seeds; character and motions of the gall of Cymips q.-saltatorius.
- - Habits of *Pollenia rudis*; notice of recorded cases of swarming among the Diptera. See No. 2128.

- 2175. [BILEY, C. V.] The new classification of the Coleoptera of North America. <Amer. Nat., June [17 May], 1883, v. 17, pp. 660-661.
 - Notice of work of J. L. Leconte and G. H. Horn.
- 2176. BILEY, C.V. A pretty and unique gall-making tortricid. < Amer. Nat., June (17 May), 1883, v. 17, p. 661, fig. 1.</p>
 Description and figure of image of Grapholitha minana n. sp. reared from galls on stems of Acadia flicing in Arizona.
- 2177. [RILEY, C. V.] Simulium feeding on other insects. < Amer. Nat., June [17 May], 1883, v. 17, pp. 661-662.

 Comment on note of H. A. Hagen.
- 2178. [RILEY, C. V.] Synopsis of the N. A. Heliothina. < Amer. Nat., June [17 May], 1883, v. 17, pp. 662-663.

 Review of paper of J. B. Smith; nature of the generic characters of Noctuids.
- 2179 [RILEY, C. V.] Death of Professor Zeller. < Amer. Nat., June [17 May], 1883, v. 17, p. 663.
 Obituary notice of P. C. Zeller.
- 2180. [RILEY, C. V.] Protection of insect collections. <Amer. Nat., June [17 May], 1883, v. 17, pp. 663-664.

 Power of the larvæ of Dermestidæ to endure the effects of certain insecticides; prerequisites more important than insecticides; seasons when collections are most endangered.
- 2181. [RILEY, C. V.] Cocoon of Telea polyphemus. < Amer. Nat.,
 June [17 May], 1883, v. 17, p. 664.

 Answer to inquiry of H. Morris; cocoon of Telea polyphemus usually falls to
 - Answer to inquiry of H. Morris; cocoon of *Telea polyphomus* usually falls to the ground with the leaves, yet quite frequently it is attached to twigs and does not fall to the ground.
- 2182. [RILEY, C. V.] The sucking organs of bees, wasps, and flies. Amer. Nat., June [17 May], 1883, v. 17, pp. 664–665.

 Notice of paper of K. Kraepelin.
- 2183. [RILEY, C. V.] The "pine moth of Nantucket." < Amer. Nat.,
 June [17 May], 1883, v. 17, pp. 665-666.
 Critical review of paper of S. H. Scudder.
- 2184. [RILEY, C. V.] Entomological notes. < Amer. Nat., June [17 May], 1883, v. 17, pp. 666-667.

 Brief notes on recent publications.
- 2185. RILEY, C. V. Elephantiasis or Filaria disease. <Science, 18 May, 1883, v. 1, pp. 419-421, fig. S. b. No. 42, p. 51, fig. Criticism of the views of Dr. A. F. A. King; notice of the writings of P. Manson and others on the connection of Cular musquite with the life-history of Filaria sanguinis-hominis and on the production of elephantiasis and related diseases by Filaria.</p>
- 2186. RILEY, C. V. Nemestrinidæ. <Science, 8 June, 1883, v. 1, p. 513.

 Note on papers concerning Nemestrinidæ; supplementary to No. 2169.
- 2187. RILEY, C. V. The corn-root Diabrotica. <Rural New-Yorker, 9
 June, 1883. S.-b. No. 54, p. 42.
 Increasing distribution of Diabrotica longicornis; means against it.

- 2188. RILEY, C. V. Elm-leaf beetle. <Sci. Amer., 16 June, 1883, v. 48. S.-b. No. 42a, p. 265.
 - Reply to inquiries concerning Galeruca xanthomelana.
- 2189. RILEY, C. V. A unique and beautiful noctuid. <Amer. Nat., July [20 June], 1883, v. 17, pp. 788-790, fig.
 - Figure of image of Cirrhophanus triangulifer; description of its generic characters; its affinities, synonyms, and probable habits.
- 2190. [RILEY, C. V.] Insects affecting stored rice. < Amer. Nat., July [20 June], 1883, v. 17, p. 790.
 - Lists of insects, chiefly Coleoptera, found in a lot of damaged rice from the Chinese centennial exhibit; two species are carnivorous.
- 2191. RILEY, C. V. Hypermetamorphoses of the *Meloidæ*. <Amer. Nat., July [20 June], 1883, v. 17, pp. 790-791.
 - Proposal of simpler and more natural terms to designate the stages of development of larvæ of Meloidæ.
- 2192. [RILEY, C. V.] Entomological notes. Amer. Nat., July [20 June], 1883, v. 17, pp. 792–793.

 Comments on recent entomological publications; items of news.
- 2193. RILEY, C. V. The grape Phylloxera in France. <Science, 22 June, 1883, v. 1, pp. 576-578.
 - Review and criticism of the report of the Commission supérieure du Phylloxers.

 M. Riter (1) V. Fig. inspects. Science 29 June 1883 v. 1 p. 500
- 2194. RILEY, C. V. Fig insects. <Science, 29 June, 1883, v. 1, p. 599. Review of S. S. Saunders's views on fig insects.
- 2195. RILEY, C. V. Egg-punctures on raspberry- and grape-vines, etc. <Rural New-Yorker, 30 June, 1883, v. 42, p. —. S. b. No. 42, p. 56.
 - Answer to inquiry of T. H. G.; oviposition of *Ecanthus niveus* in stems of raspberry- and grape-vines, and of *Ceresa bubalus* in twigs of apple-trees; habits and ravages of and means against the former; the latter seldom very injurious.
- 2196. RILEY, C. V. Silk culture in the United States. <Rural New-Yorker, 14 July, 1883, v. 42, p. —. S.-b. No. 42a, p. 531.</p>
 Revival of interest in silk culture in the United States; experience of the year; relation of silk culture to import duties; warning against too high expectations as to profits; notes on manuals of silk culture.
- 2197. RILEY, C. V. Economic entomology of Iowa. <Sci. Amer., 14
 July, 1883, v. 49, p. —. S.-b. No. 53, p. 147.
 Review of work done in Iowa, especially of papers of J. N. Dixon, H. Osborn, and A. B. Walton.
- 2198. [RILEY, C. V.] Insect plagues. <Boston Herald, 22 July, 1883, p.—. S.-b. No. 42, pp. 53-54; No. 67, p. 11. Extract: <Mirror and Farmer, 26 July, 1883, v. 35. S.-b. No. 42, pp. 126-127. Newspaper interview; seasons, habits, ravages of and means against Caloptenus atlanis, Nematus cricheonii, and Orgyia leucostigma.
- 2199. RILEY, C. V. Report by C. V. Riley. < Proc. U. S. Nat. Mus.,
 27 July, 1883, v. 6, pp. 104-105. Reprint: < Sci. Amer.
 Suppl., 13 October, 1883, v. 16, p. 6486.
 - Note to F. Humbert's Lucilia macellaria infesting man; references to other mentions of it; its distribution and means against it.

20 ENT

- 2200. RILEY, C. V. Emulsions of petroleum and their value as insecticides. <Proc. Amer. Assoc. Adv. Sci. for 1882, 1883, v. 31, pp. 469-470. Separate: <Salem, Mass., July, 1883, pp. 469-470. Reprint: <Kansas City Rev. of Science and Industry, November, 1883, v. 7, pp. 447-448. S.-b. No. 42, p. 124. Extract: <Sci. Amer., 19 November, 1883, v. 49, p. 294. S.-b. No. 51, p. 153.</p>
 - Description of modes of making emulsions of petroleum for use against insects.
- 2201. [RILEY, C. V.] The old, old question of species. <Amer. Nat., September [15 August], 1883, v. 17, p. 975.

Comments upon the discussion between H. A. Hagen and W. H. Edwards; views of both parties extreme; views as to the true nature of species.

- 2202. [RILEY, C. V.] Myrmecophila. <Amer. Nat., September [15 August], 1883, v. 17, pp. 975-976.
 - Record of recent captures of Myrmecophila in Oregon and District of Columbia; record of former captures in the United States; habits of the genus.
- 2203. [RILEY, C. V.] Salt-water insects used as food. <Amer. Nat., September [15 August], 1883, v. 17, pp. 976-977.

 Occurrence of Ephydra (hians?) in Lake Tetscoco, Mexico; E. gracilis found in Great Salt Lake, Utah, and E. californica in lakes in California; account by W. H. Brewer of the manner in which the last-named species is collected
- and used for food by the Indians living near Mono Lake.

 2204. [RILEY, C. V.] Food plants of Samia cynthia. <Amer. Nat.,
 September [15 August], 1883, v. 17, p. 977.
 - Review of paper of H. H. Birney; list of plants on which Samia [= Attacus] cynthia has hitherto been found feeding; some of these are the favorite food-plants of Callosamia [= Attacus] promethea.
- 2205. [RILEY, C. V.] Bitten by an aphid? <Amer. Nat., September [15 August], 1883, v. 17, p. 977.
 - Letter of S. Swan, with answer; Siphonophora [= Nectarophora] rudbeckies common on Solidago and Rudbeckia; the biting was probably caused by ants or some other insect that escaped notice at the time.
- 2206. RILEY, C. V. Steganoptycha claypoleana. <Amer. Nat., September [15 August], 1883, v. 17, p. 978. Reprint: <Papilio, September-December, 1883, v. 3, p. 191.
 - Comparative description of Steganoptycha claypoleana with Protectoras asculanum; habits of the former.
- 2207. R[ILEY], C. V. Extermination and restriction of Phylloxera in Switzerland. <Rural New-Yorker, 25 August, 1883, v. 42. S.-b. No. 49, p. 74.

Notice of report of Valery-Mayet.

- 2208. RILEY, C. V. Hackberry psyllid galls. <Ca. Ent., August [5 September], 1883, v. 15, pp. 157-159, figs. 6, 7.
 - Critical review of paper of T. W. Fyles; Phyllocera vastatrix has many parasites and Celtis is attacked by many species of gall-insects; characters of Pachypsylla n. g. [p. 157]; figures of galls of P. celtidis-venusta and P. c.-mamma; derivation and orthography of the generic term Celtis.

- 2209. RILEY, C.V. Some recent discoveries in reference to Phylloxera.

 <Science, 7 September, 1883, v. 2, p. 336. S.-b. No. 42, pp. 68,
 69. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1883, 1884,
 v. 32, p. 320. Separate: <Salem, Mass., July, 1884, p. 320.

 <Amer. Nat., December [28 November], 1883, v. 17, p. 1288.
 - Summary of the cycle of development of the genus *Phylloxera*; character of the gall of *P. spinosa* and location of the impregnated egg of the species.
- 2210. RILEY, C. V. The *Psyllidæ* of the United States. <Science, 7 September, 1883, v. 2, p. 337. S.-b. No. 42, pp. 67-68. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1883, 1884, v. 32, p. 319. Separate: <Salem, Mass., July, 1884, p. 319.</p>
 - Characteristics and economic importance of Psyllidæ; status of the present knowledge of this family in the United States; list of new genera and species; food-plants of the same; characteristics of the eggs and young of Psyllidæ.
- 2211. RILEY, C. V. Improved method of spraying trees for protection against insects. <Science, 14 September, 1883, v. 2, p. 378. S.-b. No. 42, p. 68. Reprint: <Proc. Amer. Asoc. Adv. Sci. for 1883, 1884, v. 32, pp. 466–467. Separate: <Selem, Mass., July, 1884, pp. 466, 467.
 - Description of apparatus, especially of "cyclone nozzle" and adjustable hose; mention of insecticide substances.
 - Note.—Nos. 2209-2211 were issued as a separate, Salem, 1884.
- 2212. [RILEY, C. V.] Entomology at Minneapolis. <Amer. Nat., 1883, v. 17: October [17 September], pp. 1068-1070; November [19 October], pp. 1169-1174.
 - Minutes of the meetings of the entomologists at the session of the American Association for the Advancement of Science; reorganization of the entomological club; election of officers and change in the constitution; abstracts of papers read.
- 2213. RILEY, C. V. Notes on Padisca scudderiana. <Amer. Nat., October [17 September], 1883, v. 17, pp. 1069-1070.
 - Habits and variation of Padisca scudderiana; difference between the gall of this insect and that of Gelechia gallasolidaginis.
- 2214. RILEY, C. V. A myrmecophilous lepidopteron. <Amer. Nat., October [17 September], 1883, v. 17, p. 1070.
 - Larva of Helia americalis found in nests of Formica rufa; this species the only one of the Lepidoptera known to develop in ants' nests.
- 2215. [RILEY, C. V.] Enemies of the egg-plant. <Amer. Nat., October [17 September], 1883, v. 17, p. 1070.
 - Extract from a letter of A. Œmler, proving that the occurrence of Cassida texana and Doryphora juncta on Solanum melongena is not accidental or temporary.
- 2216. [RILEY, C. V.] The periodical Cicada in southeastern Massachusetts. < Amer. Nat., October [17 September], 1883, v. 17, p. 1071.</p>
 Note to paper of C. E. Bessey; Cicada [= Tibicen] septendecim at Martha's Vineyard, Mass., in June, 1883, the precursor to septendecim Brood XXI which will appear in North Carolina and Virginia as well as Martha's Vineyard in 1884; accelerated specimens of the same brood received from

Loudoun County, Va., recently.

- 2217. RILEY, C. V. Habits of *Murmidius*. <Amer. Nat., October [17 September], 1883, y. 17, p. 1071.
 - List of insects found in a lot of damaged rice from South America; occurrence of Murmidius ovalis in vast numbers in this rice; its probable food-habits; description of its cocoon; list of families of beetles some of whose larve spin cocoons; habitat of Mychocerus.
- 2218. [RILEY, C. V.] Obituary. < Amer. Nat., October [17 September], 1883, v. 17, p. 1072.

 Notices of V. T. Chambers and Townend Glover.
- 2219. [RILEY, O. V.] Entomological notes. < Amer. Nat., October [17 September], 1883, v. 17, pp. 1072–1073.
- Notes on recent entomological publications; items of news.
- 2220. [RILEY, C. V.] Economic notes. < Amer. Nat., October [17 September], 1883, v. 17, pp. 1073–1074.</p>
 Comments on the ravages of several jujurious insects during the past season.
- 2221. RILEY, C. V. A parasite of the cabbage-worm. < Rural New-Yorker, 6 October, 1883, v. 42. S.-b. No. 42, p. 58.
 - Answer to letter of J. H. B.; parasitism of Pteromalus puparum in larvæ and pupæ of Pieris rapæ.
- 2222. RILEY, C. V. The handmaid moth. <Rural New Yorker, 13 October, 1883, v. 42. S.-b. No. 42, p. 77.
 - Answer to inquiry of H. B. S.; description of larva, pupa, and image of Datana ministra from hickory- and walnut-trees, and of a phytophagic variety of the larva from apple and other trees; habits of the larvæ; the larvæ unusually abundant in 1883.
- 2223. RILEY, C. V. Remarks on Arzama obliquata. <Amer. Nat., November [19 October], 1883, v. 17, p. 1169.</p>
 Description of the egg-mass of Arsama obliquata; colors and habitat of the larva; variations of the imago; number of annual broods.
- 2224. [RILEY, C. V.] Rare monstrosities. <Amer. Nat., November [19 October], 1883, v. 17, p. 1175.

 Notice of monstrosities recorded in Melanippe montanata and Zygana minos.
- 2225. [RILEY, C. V.] The nervous system of insects. <Amer. Nat., November [19 October], 1883, v. 17, pp. 1175-1176. Summary of the observations of Ed. Brandt.
- 2226. [RILEY, C. V.] Hymenorus rufipes as a myrmecophilous species.

 Amer. Nat., November [19 October], 1883, v. 17, p. 1176.
 Images of Hymenorus rufipes raised from larvæ found in nests of Formics fuses;
 character of the pasts of the Formics: food habits of the Hymenorus un-
 - Imagos of Hymenorus rufipes raised from larvæ found in nests of Formica fuses; character of the nests of the Formica; food-habits of the Hymenorus unknown.
- 2227. [BILEY, C. V.] Recent publications. < Amer. Nat., November [19 October], 1883, v. 17, p. 1177.
 Notice of J. H. Comstock's work on Diaspina and of other recent publications.
- 2228. [RILEY, C. V.] Entomological notes. < Amer. Nat., November [19 October], 1883, v. 17, pp. 1177-1179.
 - Review of "General Index of the Entomological Reports of the Province of Ontario;" recent publications and items of news.

99

101

2229. RILEY, C. V. The potato-stalk borer. < Rural New-Yorker, 20 October, 1883. S.-b. No. 42, pp. 78-79.

Answer to letter of S. C. R.; description, habits of, and means sgainst Gortyna nitela; means against Paria aterrima, Graphops pubescens, and other larvæ of Chrysomelidæ injurious to the roots of strawberry-plants.

2230. [RILEY, C. V.] Recent advances in horticultural entomology.

<Rural New-Yorker, 20 October, 1883, v. 42. S.-b. No. 42, pp. 79-81. Reprint: <Proc. 19th Sess. Amer. Pomol. Soc., 1884, p. 45. <Trans. Wisc. State Hortic. Soc., 1886, v. 17, p. —.

Report of address delivered; discussion of measures recommended for adoption to prevent the ravages of insects injurious to horticulture, especially of Carpocapsa pomonella and Conotrachelus nenuphar; correction of statements in regard to the oviposition of Saperda bivittata [= candida] and † Bembex marginata; advance in knowledge of the life-history of Aphidida and in the development of machinery for the application of poison sprays to plants; relative value of the principal insecticides.

2231. RILEY, C. V. On a gall-making genus of *Apionine*. <Bull. Brooklyn Ent. Soc., October, 1883, v. 6, pp. 61-62. Separate: <Brooklyn, N. Y., 1883, 2 pp.

List of North American gall-making Coleoptera; description of *Podapion* n. g. [p. 62] and of the gall and image of *P. gallicola* n. sp., found on twigs of *Pinus inops*; probable life-habits, inquilines, and parasite of this species.

2232. RILEY, C. V. Report of the Entomologist. <Ann. Rept. [U. 8.]

Commissioner Agric. for 1883, pp. 99–180, 13 pl. Separate:

<Washington, 31 October, 1883, pp. 5+pp. 99–180+pp. 7, 13 pl.

TABLE OF CONTENTS.

Scope and limitations, 99—Cabbage insects, 99—Lesser locust, 99—Protection of fruit and shade trees, 99—Office force, assistants and agents, 99—Observations and report on insects injurious to forest-trees, by Dr. A. S. Packard, jr., 99—Special report on insects affecting the orange, by H. G. Hubbard, 99—Work of agents, 100—Monographs in preparation, 100—Commission sent to Brazil to study certain insects, 100—Publication of third report of the United States Entomological Commission, 100—Publication of special bulletins, 100—Correspondence and work of the division, 101—The illustrations to the report, 101.

Legislation, 101—California State Board of Silk Culture, 101—Premiums offered in California, 101—Home culture, 102—Guaranty of egg supply, 102—Woman's Silk Culture Association of California, 102—California Silk Culture Association, 102—Reports from correspondents of the division, 102—104—Issue of "Silk Culture Directory," 104—Establishment of school in New Jersey, 105—Organization of Southern Silk Industrial Association, 105—A new machine patented, 105—Report on the future of silk culture in the United States, by United States Consul Peixotto, of Lyons, France, 105—Silk culture by the Mennonites, 106—Work at the Department, 106.

2232. RILEY, C. V.—Continued.	
Cabbage worms	. 107
Interest in and importance of the subject, 107—Accurate estimations of loss impossible, 107.	te
Imported cabbage-worm, Pieris rapa	. 108
History of its spread, 108—Ravages, 109—Characters, 110—Habit	
110—Food-plants, 111—Seasons of appearance, 111—Natural enuica, 111.	
Southern cabbage-butterfly, Pieris protedics	. 114
Range, 114—Ravages, 114—Characters, 114—Food-plants, 115- Parasites, 115.	
The potherb-butterfly, Pieris oleracea	. 115
Range, 115—Characters, 116—Habits, 116—Number of broods, 116- Food-plants, 117—Enemies, 117.	
The larger cabbage-butterfly, Pieris monuste	. 117
Rauge, 117—Characters, 117—Food-plants, 118—Parasites, 118—Description of early states, 118.	
The cabbage plusia, Plusia brassica	. 119
Its range, 119—Food-plants, 119—Habits and natural history, 119-	
Close relation to an European species, 120-Natural enemie	3,
120—Botrytis rileyi, a new species of fungus, 121—Remedies, 12	l.
The cabbage mamestra, Mamestra trifolii	. 123
Its habits and natural history, 123—Remedies, 124—Description 124.	1,
The zebra cabbage-worm Mamestra picta	. 124
History, 124—Description, 125—Habits, 125—Broods, 125—Remedies, 125.	
The vabbage pionea Pionea rimosalis	. 196
Its past history, 126—Notes by Prof. Cyrus Thomas, 126—Habit and natural history, 127—Parasites, 127—Remedies, 127—De	
scriptive, 128.	
The cauliflower botis, Botis repetitalis	. 128
Past history, 128—Range, 128—Descriptive, 129.	
The cabbage plutella, Plutella cruciferarum	. 129
Its past history, 129—Habits and natural history, 130.	
Remedies for Cabbage-Worms. Hot water, 131—Pyrethrum, 131—Kerosene emulsions, 131—Othe substances, 131—Paris green, London purple, and white hell bore, 132—Preventive measures, 132—Report of Capt. R. & Lacey, 133—Report of Col. Wright Rives, 134—Poisoning descriptions.	or 8- 3.
vices, 136.	_
Causes of destruction of evergreen forests in New Englan and New York, by A. S. Packard, Jr	. 138
The larch saw-fly, Nematus erichsonii	
History of ite ravages, 138—Its devastations in Maine, 139—I	
ravages in New Hampshire, 141—Its appearance in Massach	
setts, 141—In northern New York, 142—History of the specie	
and its habits, 142—Description of egg, larva, cocoon, and imag	υ,
145—Remedies, 146—Parasites, 146.	140
The spruce-bud tortrix, Tortrix fumiferana	
Its ravages in Maine, 146—Habits and transformations, 147—Do_scription of different stages, 148.	
The spruce nematus, Nematus integer	. 149
Range, 149—Description of different stages, 150.	

2232.	RILEY, C. V.—Continued.	
	CAUSES OF DESTRUCTION OF EVERGREEN FORESTS, ETC.—Continued.	
	The hemlock gelechia, Gelechia "bietisella	150
	Habits, 151—Description of larva, pupa, and moth, 151.	
	EXPERIMENTS ON SCALE INSECTS, WITH PRACTICAL SUGGESTIONS, BY H. G. HUBBARD	152
	Report of progress in experiments, 152—Soap emulsions, 152—Kerosene and soap emulsions, formula, 152—Receipt for making emulsions of kerosene and soap, 152—Introduction and use of the cyclone nozzle, 152—Effects of kerosene emulsions upon plants, 153—Effects of kerosene emulsions varied by change of weather,	
	153—Most favorable season for applying kerosene emulsions, 153—Application of liquid insecticides, 153—The cyclone nozzle, 154—Cost of kerosene wash, 154.	
	Recent experiments	155
	Potash compared with soda as an insecticide, 155—Details of experiments with potash, 155-156—Experiments with soda, 156.	
	Introduction and spread of scale insects	156
	Great vitality of bark-lice, 156—Importation on live trees, 157—	
	Precautionary measures; infection from nursery stock, 157—Pro-	
	tection afforded by hedges and forest trees, 158. MISCELLANEOUS INSECTS	159
	The imported elm-leaf beetle, Galeruca xanthomelana	159
	Rauge, 159—An importation from Europe, 160—Habite and natu-	1170
	ral history, 160—Remedies, 161—Natural enemies, 163—More re-	
	cent-experiments at the Department, 164-Past history of the	
	elms on Department grounds, 164—Condition and characteris-	
	tics of the grove in 1882 and 1883, 164—Extent of injury in 1882	
	and 1883, 164—Preferences of the elm-beetle for certain varieties	
	and species of elms, 165—Effects of arsenical poisons on insect	
	and plant, 165—Preventive effects of the poison, when best applied, 166—Treatment with London purple, 166—Preparation of	
	the poison, 166—Effects of the mixture, 167—Treatment with Paris green, 167—Mechanical means of applying the poison,	
	168—The eddy-chamber nozzle, 168—Hose and bamboo combina- tion, 169.	
	The lesser migratory locust, Calopienus atlanis	170
	Its ravages in the Merrimac Valley, 170—Historical, 170—Characters, 172—Range and life-history, 173—Natural enemies, 174—	
	Remedies, 175—Destruction of eggs, 175—Destruction of young,	
	175—Coal oil, 175—Machines, 175-177—Coal tar, 177—Catching	
	or bagging, 178—Protection of vegetable gardens, 179—Necessity for co-operation, 180.	
2233.	RILEY, C. V. The imported orchard Scolytus (Scolytus rugul	08 W8
	Ratz.). <thomasville 10="" 1883.<="" [ga.]="" november,="" td="" times,=""><td>3b.</td></thomasville>	3b.
	No. 56, p. 182.	
	Answer to inquiry; S. rugulosus bores in twigs of pear; means against i	t.
2234.	RILEY, C. V. Osage orange vs. mulberry for the silk-wo	rm.
	<sci. 17="" 1883,="" 305.="" 49,="" amer.,="" b.="" no.<="" november,="" p.="" s.="" td="" v.=""><td></td></sci.>	
	p. 91. Reprint: <amer. 18,="" 1884,="" 5="" january,="" nat.,="" pp.<="" td="" v.=""><td></td></amer.>	

42, p. 118.

Comparative value of leaves of *Maclura aurantiaca* and of *Morus* as food for *Sericaria mori*; critical review of V. des Lauriers's conclusions.

79. < Prairie Farmer, 22 March, 1884, v. 56, p. 187. S.·b. No.

- 2235. RILEY, C. V. Entomological notes of the year. < Prairie Farmer, 24 November, 1883, v. 55. S.-b. No. 42, pp. 86-87.
 - Progress of experiments for the destruction of scale insects, Coccida; introduction of Aspidiotus rapax into California on apples from New Zealand; strawberries injured by Capsus oblineatus [= Lygus pratensis] and a myriapod in Illinois; occurrence of Leucania unipuncta in several places, and of Cecidomyia destructor in Illinois; extension of culture of Pyrethrum cinerariæfolium in California; occurrence of Anarsia lineatella on strawberryplants in Illinois; of Doryphora juncta and Cassida texana on Solanum melongena in Georgia; of Ceresa bubalus on potato-plants in Pennsylvania; of Epilachna corrupta on wax-beans, etc.
- 2236. RILEY, C. V. A satisfactory remedy for melon bugs, flea-beetles, < Rural New-Yorker, 3 November, 1883, v. 42. S.-b. No. 42, pp. 77-78.
 - Quinn's method of sprinkling the vines with a mixture of tobacco water and soft soap and then powdering with lime is probably the best general preventive against Diabrotica vittata and Halticida; description of J. M. Nicholson's siphon arrangement by which to keep the vines constantly moist with the liquid.
- 2237. RILEY, C. V. The chinch-bug in New York. <Science, 9 November, 1883, v. 2, p. 621. Extract: <Sci. Amer., 22 December, 1883, v. 49, p. 384. S.-b. No. 42, pp. 122-123.
 - Present outbreak of Blissus leucopterus in New York the result of an increase due to favorable conditions rather than an invasion.
- 2238. R[ILEY], C. V. Insects in relation to agriculture. < Stoddart's Encylopædia Americana, 1883, v. 1, pp. 135-142, figs. 1-29.
 - Chapter 9 of article "Agriculture." Brief accounts with numerous illustrations of the insects named below, and of means against them, with cross-references to accounts of other insects in other portions of the work. The headings and subjects of the sub-chapters are as follows:

IMPORTANCE OF ECONOMIC ENTOMOLOGY

INSECTS INJURIOUS TO FRUIT AND FRUIT TREES:

Apple curculio (The), Anthonomus quadrigibbus Say, p. 135, fig. 1-Apple-magget (The), or "railroad-worm," Trypeta pomonella Walsh, p. 135—White-marked tussock-moth (The), Orgyia leucostigma Sm. and Abb., pp. 135-136, fig. 2-Apple-tree tent-caterpillar (The), Clisiocampa americana Harr., p. 136, figs. 3-4-Fall webb-worm (The), Hyphantria textor Harr. [= cunea], p. 136, fig. 5-Oyster-shell bark-louse of the apple (The), Mytilaspis pomicorticis Riley [= pomorum], pp. 136-137, fig. 6-Round-headed appletree borer (The), Saperda bivittata Say [= candida], p. 137, fig. 7-Flat-headed apple-tree borer (The), Chrysobothris femorata Fabr., p. 137—Spring canker-worm (The), Paleacrita vernata Peck, p. 137, figs. 8-9-Full canker-worm (The), Anisopteryx pometaria Harris, p. 137, figs. 10-11—Peach-tree borer (The), Ageria [= Sannina] exitiosa Say, pp. 137-138, fig. 12 - Currant-stalk borer (The), Ageria tipuliformis L., p. 138-Imported current-worm (The), Nematus ventricosus Klug. [= ribesii], p. 138, figs. 13-14-Native currant-worm (The), Pristiphora grossulariæ Walsh, p. 138-Snowy tree-cricket (The), Œcanthus niveus Harr., p, 138, figs. 15-16.



135

2238. R[ILEY], C. V.—Continued.

INSECTS INJURIOUS TO CEREALS AND FORAGE-CROPS:

White-grub (The), Lacknosterna fusca Froh., pp. 138-139, fig. 17—Clover-seed midge (The), Cecidomyia leguminicola Lintner, p. 139—Joint-worm (The), Isosoma hordei Harr., p. 139, fig. 18—Wheat midge (The), Diplosis tritici Kirby, p. 139—Cut-worms, family Noctuida, genera Agrotis, Mamestra, Hadena, and Prodenia, p. 139—Wire-worms, family Elaterida, p. 139.

INSECTS INJURIOUS TO GARDEN VEGETABLES:

Imported cabbage-worm (The), Pieris rapæ Schrank, p. 139, figs. 19-20—Southern cabbage butterfly (The), Pieris protodice Boisd., p. 140—Potherb butterfly (The), Pieris oleracea Boisd., p. 140—Cabbage plusia (The), Plusia brassicæ Riley, p. 140, fig. 21—Harlequin cabbage-bug (The), Murgantia histrionica Hahn, p. 140, fig. 22—Pea-weevil (The), Bruchus pisi Linn., p. 140, fig. 23—Bean-weevil (The), Bruchus fabæ Riley, p. 140, fig. 24—Blister-beetles, meloid genera Macrobasis, Epicauta, etc., p. 140, fig. 25—Striped cucumber-beetle (The), Diabrotica vittata Fabr., pp. 140-141, figs. 26-27.

INSECTS INJURING MISCELLANEOUS FIELD-CROPS:

Cotton-boll worm (The) or corn-ear worm, Heliothis armigera, Hübn., p. 141, fig. 28—Tobacco-worm (The), Sphinx [= Protoparce] carolina L., p. 141.

INSECTS INJURIOUS TO THE VINE.

INSECTS INJURIOUS TO LIVE-STOCK:

Bot-fly of cattle (The), Hypoderma bovis Latr., p. 141—Sheep bot-fly (The), Estrus ovis Linn., p. 141, fig. 29—Horse bot-fly (The), Gastrophilus equi Fabr., pp. 141-142.

142

- 2239. R[ILEY], C. V. Army-worm, Leucania unipuncta, Haw. <Stoddart's Encylopædia Americana, 1883, v. 1, pp. 317-318, 2 figs. Distribution, seasons, habits, and food-plants of and means against Leucania unipuncta; references to the more important articles on this insect; figures of larva and imago.
- 2240. RILEY, U. V. An epidemic disease of *Caloptenus differentialis*.

 <Amer. Nat., December [28 November], 1883, v. 17, p. 1287.</p>
 Note to communication of H. Osborn; the Entomorphinora calopteni a result rather than the cause of disease.
- 2241. RILEY, C. V. The growth of insect eggs. < Amer. Nat., December [28 November], 1883, v. 17, p. 1289.

 Notice of paper of J. A. Osborne: swelling of the eggs of Phanerontera [= Scud-
 - Notice of paper of J. A. Osborne; swelling of the eggs of *Phaneroptera* [= Scudderia] ourvicauda probably connected with embryological development.
- 2242. [RILEY, C. V.] Protective device employed by a glaucopid caterpillar. <Amer. Nat., December [28 November], 1883, v. 17, p. 1289.
 - Notice of paper of F. Müller; general use of shed hairs by larvæ of Arctiidæ in the construction of their cocoons; description of method in which the larva of Eunomia eagrus arranges its shed hairs to form a protection for the pupa.
- 2243. [RILEY, C. V.] Saw-fly larvæ on the quince. <Amer. Nat., December [28 November], 1883, v. 17, p. 1289.

Notice of paper of J. A. Lintner; food-plants of Selandria [= Eriocampa] cerasi.

2244. [RILEY, C. V.] Entomology in New York. < Amer. Nat., December [28 November], 1883, v. 17, pp. 1289-1291.</p>
Review of J. A. Lintner's first annual report as State entomologist; list of

Review of J. A. Lintner's first annual report as State entomologist; list of subjects treated in the report.

- 2245. [RILEY, C. V.] Fruit insects in California. < Amer. Nat., December [28 November], 1883, v. 17, p. 1291.</p>
 Review of M. Cooke's "Injurious Insects of the Orchard."
- 2246. [RILEY, C. V.] Death of Dr. J. L. Leconte. < Amer. Nat., December [28 November], 1883, v. 17, p. 1291.</p>
 Notes the loss to entomologists occasioned by the death of J. L. Leconte.
- 2247. [RILEY, C. V.] Entomological notes. < Amer. Nat., December [28 November,] 1883, v. 17, pp. 1291–1292.

 Brief notices of published articles with items of news.
- 2248. [RILEY, C. V.] Economic notes. < Amer. Nat., December [28 November], 1883, v. 17, p. 1292.

Effect of Phylloxera laws in Europe; use of lime as a means against Macrodactylus; award of the Le Brun prize for the most valuable improvement relating to the silk industry.

2249. RILEY, C. V. [Raspberry canes punctured by Orchelimum glaberrimum.] < Fruit Recorder, Purdy, 1 December, 1883, v. 15. S.-b. No. 42, p. 92.

Description of and means against Orchelimum glaberrimum.

- 2250. [RILEY, C. V.] The Phylloxera in sandy soil. <Rural New-Yorker, 1 December, 1883, v. 42. S. b. No. 42, p. 97. Reprint: <Amer. Nat., January, 1884 [29 December, 1883], v. 18, p. 78. Condition of the grape crop in France; favorable account of the use of American grape-stocks; Phylloxera vastatrix harmless in sandy soils.</p>
- 2251. RILEY, C. V. Bacterial disease of the imported cabbage-worm. <Sci. Amer., 1 December, 1883, v. 49, p. 337. S.-b. No. 42, pp. 91–92. Reprint: <Amer. Nat., January, 1884 [29 December, 1883], v. 18, p. 80.
 - Notice of observations of S. A. Forbes on the death of larvæ of *Pierie rapa* from infection by *Bacterium*; previous mention of this disease.
- 2252. RILEY, C. V. The chinch bug in New York State. <Sci. Amer., 8 December, 1883, v. 49. p. 359. Reprint: <Amer. Nat., 5 January, 1884, v. 18, pp. 79-80.
 - Critical review of papers of J. A. Lintner; the occurrence of *Blissus leucopterus* in New York in unusual abundance in 1882 and 1883 is not warrant for great alarm. See No. 2271.
- 2253. RILEY, C. V. Reports of observations and experiments in the practical work of the division, made under the direction of the entomologist. <Bull. No. 3, Div. Ent. U. S. Dept. Agric., [8 December], 1883, 75 pp., 3 pl.

CONTENTS.

Preface	7
Further notes on the army worm (Leucania unipuncta)	9
Experiments with pyrethrum	16

2253. RILEY, C. V.—Continued.

CONTENTS-Continued.

Notes on forest-tree insects. By A. S. Packard, jr	24
The cotton-worm in South Texas in 1883. By E. H. Anderson	31
Test of machinery for destroying the cotton-worm. By W. S. Barnard.	39
The tree-borers of the family Cossidæ. By J. S. Bailey	49
Tests of silk fiber from cocoons raised at the Department. By W. Mc-	
Murtrie	56

- 2254. RILEY, C. V. Entomology in New York. < Rural New-Yorker, 15 December, 1883. S. b. No. 42, pp. 85-86.
 - Review of J. A. Lintner's first report (for the year 1881) as State entomologist of New York.
- 2255. RILEY, C. V. Lucilia macellaria. <Sci. Amer., 15 December, 1883, v. 49, p. 373. S.-b. No. 42, p. 93.
 Critical review of paper of F. Humbert.
- 2256. RILEY, C. V. Dipterous larvæ in the human body. <Sci. Amer., 22 December, 1883, v. 49, p. 385. S.-b. No. 42, p. 90.
 - References to and notices of several articles on the occurrence of larvæ of Diptera in the human body; synonymy and distribution of Compsomyia [= Lucilia] macellaria; myasis caused solely by larvæ of Sarcephagidæ and Muscidæ; Estridæ occur rarely and exceptionally in the human body.
- 2257. BILEY, C. V. Capitalizing specific names. <Papilio, September-December, 1883, v. 3, pp. 164-166. Separate: <[N.Y.], 3 pp. Comments on reasons given by W. H. Edwards and others for the uniform capitalizing of specific names. See No. 2172.
- 2258. R[ILEY], C. V. Townend Glover. <Papilio, September-December, 1883, v. 3, pp. 167-168.

 Obituary notice.
- 2259. [RILEY, C. V.] The genus Colias. <Amer. Nat., [5] January, 1884, v. 18, pp. 74-76.
 - Review of paper of H. A. Hagen; discussion of the species of Colias in North America; plastic nature and classificatory characters of the genus.
- 2260. [RILEY, C. V.] Larval habits of the dipterous family *Dexida*. <Amer. Nat., [5] January, 1884, v. 18, pp. 76-77.
 - Notice of paper of F. Brauer; parasitism of Decia rustica and of Phorostoma latum on the larva of Rhizotrogus and of Melanophora idiabrotica on Diabrotica rittata.
- 2261. [RILEY, C. V.] Entomological notes. <Amer. Nat., [5] January, 1884, v. 18, pp. 80-81.
 - Carpocapsa pomonella has been introduced into Australia, New Zealand, and Tasmania; occurrence of Myrmioocela ochraceella in ants' nests.
- 2262. RILEY, C. V. Recent outbreaks of the army-worm. <Rural New-Yorker, 12 January, 1884, v. 43, p. 19. S.-b. No. 56, p. 145.

 Rare occurrence of Leucania unipuncta in 1882 and 1883, following its last abundance in 1881; occurrence of its larvæ at East Windsor, Vt., in June, 1883.
- 2263. [RILEY, C. V.] The harlequin cabbage-bug, etc. < Rural New-Yorker, 2 February, 1884, v. 43, p. 70. S.-b., No. 63, p. 41.
 - Habits of and means against Murgantia histrionica; effect of chemicals applied to the soil upon the colors of flowers.

2264. RILEY, C. V. Tribute to the memory of John Lawrence Lecon	ate
Psyche, November-December, 1883 [11 February, 1884], v	
pp. 107-110. Separate: <[Cambridge, Mass., 11 Februs	шу,
1884], pp. 107-110. Notice: <psyche, 119.<="" cit.,="" loc.="" p.="" th=""><th></th></psyche,>	
Biographical notice of J. L. Leconte; his work and writings; his pers	لعور
character; disposition of his entomological collection.	
2265. [RILEY, C. V.] Fruit culture in the South. < Washington Pe	ost,
26 February, 1884. Sb. No. 42, pp. 119-120.	
Interview with a reporter; means against insects infecting the orange.	
2266. RILEY, C. V. Oviposition of the round-headed apple-tree bo	rer.
<rural 1="" 132,="" 1884,="" 43,="" 73.<="" fig.="" march,="" new-yorker,="" p="" p.="" v.=""></rural>	L-b.
No. 42, p. 85; No. 67, p. 3.	
Notice of paper of C. G. Atkins; confirmation of the account of the met	bod
of oviposition of Saperda candida; description of this method; correction	
several statements by W. Saunders in regard to the oviposition of differ	
insects; description and figure of the egg of S. candida; figures of the	our-
rows and hole of exit of this beetle; figures of the pupa and of the posi-	tion
of the egg when deposited.	
2267. [RILEY, C. V., et al.] Third report of the United States En	to-
mological Commission, relating to the Rocky Mountain locu	st,
the Western cricket, the army-worm, canker-worms, and	
Hessian fly; together with descriptions of larvæ of injurio	
forest insects, studies on the embryological development of t	
locust and of other insects, and on the systematic position	
the Orthoptera in relation to other orders of insects; with ma	
and illustrations. < Washington: 1883 [6 March, 1884], p	p.
14+347+12+92, 18 figs., 64 pls., 4 maps.	
	XI III
Preface X	ш
PART I.—IN REFERENCE TO THE ROCKY MOUNTAIN LOCUST (Caloptenus spretu	18).
CHAPTER I:	
Additions to the chronology of locust ravages in 1880 and 1881	3
The locust in 1880 in Texas, 3; in Colorado, 4; in Utah. 4; in 1881, in Texas, 6; in Utah, 7.	
CHAPTER II:	
The Rocky Mountain locust in Montana in 1880. By Lawrence	
Bruner	8
The country from St. Paul to Montana, 8-Between Bismarck and	
Fort Keogh, 9-Burning often impracticable, 9-Destroying lo-	
custs by ditches and kerosene eddies, 13—Ropes dragged to drive	
them, 15—"Drifts of hoppers," 16—Shooting against locust	
swarms, 16—Destroying by ditches and burning straw, 17—No	
more damage for three or four years, 17—Enemies of the locust	
in the Northwest, 17—Topography of western Dakota and Mon-	
tana, 19. Chapter III:	
The Rocky Mountain locust in Wyoming, Montana, etc., in 1881.	

By Lawrence Bruner.....

Letter of submittal of report by Lawrence Bruner, 21 - General report, 22-Brief history of depredations, 22-The earlier

2267. [RILEY, C. V., et al.]—Continued.

CHAPTER III-Continued.

ones, 23—Characteristics of the permanent region, 24—Settlement and other checks against locusts, 25—Distribution of the permanent regions, 25—Their physical peculiarities, 28—The arid region, 29—Permanent breeding grounds, 29—Their locations and relations, 30—The sources of locusts, 30—Interchange of swarms, 31—Regular migratory routes, 31—Agencies controlling migration, 32—Disadvantages of the temporary regions, 33—Period of hatching, 34—Period of growth, 35—Habits of the young, 36—Habits at maturity, 36—Causes of occasional over-increase, 37—Checks, enemies, 38—The efforts of experts baffled, 42—Tree culture, 42—Climatic checks, 43—Tree planting, 43—Flights affected by storms, 45—Number of annual broods, 46—Sub-permanent region, 46—Its relations, 47—Locust movements therein, 48—How to fight this insect, 49—Locust probabilities, 49—Recent swarms, 50.

CHAPTER IV:

Notes on other locusts and on the Western cricket, Anabrus. By Lawrence Bruner

Other locusts, 53—List of North American locusts north of Mexico, 55—The Western cricket, 61.

CHAPTER V:

Data obtained from solar physics and earthquake commotions applied to elucidate locust multiplication and migration, by A. H. Swinton

Importance of the central luminary, wide effects of variation in its potential energy, 65—On the tides, on chemical and organic activities, on the physical forces, on epidemics, etc., 65, 66—Periods of sun-spots, electrical effects of, 66—Thermometric effects of, 67—Volcanic effects of, 68—Entomological effects of, 68—A new sun-spot table, 69—Statement and tabulation of sun-spot dates, 69—Comparison of sun-spot periods and insect periods, 73—American locusts diminished by the spots, 73—European confirmative examples of locusts and other insects, 74—Tabulation of rare insect captures, 79—Explained, 81—Migration and distribution affected, 81—Locusts again predicted in four or ten years, 83—Conclusions, 84.

PART II.—THE ARMY-WORM, CANKER-WORMS, AND THE HESSIAN FLY.

CHAPTER VI: By C. V. Riley.

The army-worm, Leucania unipuncta. By C. V. Riley

Nomenclature, 89—Other army-worms, falsely so called, 89—Geographical distribution, 91—Injuries by, 92—Past history of, 92—Descriptive characters of, 101—The egg, 101—The larva, 102—The pupa, 103—Adults and sexual differences, 103—Habits and natural history, 105—Oviposition, 105—Habits when young, 108—Duration of larval life, 109—Traveling habits, etc., 110—Time of appearance, 112—Sudden appearance and disappearance, 114—Food-plants, 116—The pupa state, 117—Habits of the moth, 117—Flight, 118—Position at rest, 118—Number of broods yearly, 118—Hibernation, 122—Natural enemies, 125—Remedies, 128—Burning old grass, etc., 128—Predictions, meteorological influences on the species, 129—Ditching, coal-tar, poisoning, 130—Rolling, fencing, roping, 131—Report of observations by L. O. Howard, 132—Ex-

53

65

2267. [RILEY, C. V., et al.]—Continued.

CHAPTER VI-Continued.

tent of country injured, 133—Crops injured, 133—Amount of damage, 133—Previous season and crop, 134—Number of broods, 134—An accompanying cut-worm, 1.35—Natural enemies, 135—Army-worm correspondence in spring of 1882, 136—The invasion of 1880 in New Jersey, by Rev. Samuel Lockwood, 139—Performances of the worms, 139—The direction of travel, 141—Origin, eggs, etc., 143—Breeding spots, 143—Recapitulation, habits, and remedies, 145—Notes from Prof. C. V. Riley, the number of broods, hibernation, seasonal influences, 147—Bibliography, 148.

CHAPTER VII: By C. V. Riley.

Canker-worms, Paleacrita vernata, Anisopteryx pometaria. By C. V. Riley

Classification, 157—Two distinct insects concerned, 158—Differences between them, 159—Nomenclature, 162—Past history, 165—The spring canker-worm, 170—Range of the species, 170—Characters, 171—Habits and natural history, 172—Appearance of the worms, 173—Food-plants, 174—Modes of distribution, 175—Enemies, 175—Destructiveness of canker-worms, 178—The fall canker-worm, 179—Range of this species, 179—Its characters, 180—Habits and natural history, 181—Oviposition, 182—Season of appearance, 182—Pupation, 183—Food-plants, 183—Remedies and preventive measures applicable to both species, 183—Sticky substances, 183—Hanging tin band, 185—Troughs of oil, 186—Precautions and classification of contrivances, 189—Jarring and burning, 191—Washes and dustings, 191—Paris green, 192—Muriate of lime, 192—Sulphur plugged in trees, 193—Fall plowing, 193—Birds and parasites, 195—Different measures against the species, 196.

CHAPTER VIII:

PART III.—SCIENTIFIC RESULTS.

CHAPTER IX:

Descriptions of larvæ of injurious forest insects. By A. S. Packard, jr. Buprestidæ, 251—The flat-headed apple-boter, Chrysobothris femorata, 251—Chalcophora virginica, 252—Melanophila sp., 253—The flat-headed spruce-borer (Melanophila?), 254—The flat-headed peach- and cherry-borer, Dicerca divaricata Say, 255—Buprestid under hemlock bark, 255—Cerambycidæ, 256—Longicorn larva under hemlock bark, 256—Saperda on the willow, 256—The lesser pine-borer, Asemum mæstum Haldeman, 256—The oak-borer, Elaphidion parallelum Newman, 257—The common oak clytus,

157

198

263

273

277

2267. [RILEY, C. V., et al.]—Continued.

CHAPTER IX-Continued.

Xylotrechus colonus Fabr., 257—Clytus (?) larva on black-birch, 259—The ribbed rhagium, Rh. lineatum Oliver, 259—The lesser prionus, Orthosoma brunneum De Geer, 260—Unknown longicorn borer from an oak log, 261—Unknown longicorn borer in sycamore, 262.

CHAPTER X:

The embryological development of the locust. By A. S. Packard, ir. Formation of the blastoderm, 263-Origin of the primitive amæboid cells, 264—The blastodermic disc or primitive band, 264—The germinal groove, 264-Origin of the cellular or germinal layers, 264—Diagrammatic view of the origin of these layers, from Graber, 265-Embryonal membranes, 265-Serous membrane, 265-Amnion, 265—Tabular view of the eight embryonal layers, 265— Division of the embryo or primitive band into body segments, 266-Development of the appendages, 266; of the nervous system, 266; of the alimentary caual, 266-The stomodæum, 266-The proctodeum, 267—The mesenteron or stomach, 267—The pyloric appendages of the stomach, 267—The Malpighian tubes. 267-The genital glands, 267-The traches and salivary glands, 267—Spinning glands, 267—Development of the wings, 268—Criticism of Gegenbaur's views, 269—Fritz Müller's views, 269— Speculation on the primary origin of wings, 270-Correlation of metamorphosis with the acquisition of wings, and of the latter with the ripening of the sexual organs, 271-Differentiation of mesonotum and metanotum dependent on the presence of wings, 271-Development of a pupal stage, 271-Genealogy of the orders of insects, 271.

The embryological development of orthopterous insects. By A. S. Packard, jr

Rapid development of the embryo in autumn, 272—Embryology of Gryllotalpa vulgaris, 272—Formation and contraction of the embryonal skins, 272—Formation of the digestive canal, 272—Functions of the mid-gut and hind-gut, 273—Origin of the Malpighian vessels, salivary glands, tracheæ, and ovaries, 273—The sections of eggs studied by the author, prepared by N. N. Mason, 273.

2267. [RILEY, C. V., et al.]—Continued.

hatched Hylurgops, 281.

CHAPTER X-Continued.

280

The number of segments in the head of winged insects. By A. S. Packard, jr

Views generally entertained on this subject, 282—Head composed of four segments, 283—The procephalic lobes form the antennal segment, 284—The clypeus and labrum are the tergal portion of this segment, 284—The epicranium is the pleural portion, 284—The occiput is the tergal portion of the labial segment, 284—The gular region probably the base of the labium, 285.

CHAPTER XI:

The systematic position of the Orthoptera in relation to other orders of insects. By A. S. Packard, jr.....

Review of the characteristics of the four lowest orders of winged insects, 286—Probable descent of Orthoptera, Pseudoneuroptera, and Dermatoptera from a Thysanuran form, 286-Method of the present study, 286-Characters of the Phyloptera, 287-Mouthparts, 287-Thorax, 287-Wings, 288-Abdomen, 288-Metamorphosis, 288-Nomenclature of external parts of Arthropoda, 288-Sequence of orders of Phyloptera, 288-Characters of Dermatoptera, 289; of Orthoptera, 289; of Pseudoneuroptera, 290-Want of uniformity in characters of Pseudoneuroptera, 290-The diagnostic characters superficial, 290-Structure of labium, 291-Relative proportions of head-parts, 291-Subdivisions of the order. 292—Characters of Platyptera, 292; of Odonata, 292—Of Ephemerina, 293-Characters of Neuroptera, 293-Ligula, 293-Thorax, 293-Wings, 293-Abdomen, 294-Subdivisions of the order, 294-Tabular view of the grand divisions of winged insects, 294-Diagrammatic view of the genealogy of the insects, 295—Genealogy of the Hexapoda, 295; of Thysanura, 295; of Dermatoptera, 295; of Orthoptera, 296; of Pseudoneuroptera, 296; of Hemiptera, 297; of Neuroptera, 297—Derivation of cruciform larvæ, 297; of Sialidæ, 297; of thysanuriform larvæ, 297-Diversity of Neuropterous larvæ, 297-Larvæ of Sialidæ, 297; of Hemerobiidæ, 297; of Trichoptera, 298; of Panorpidæ, 298-Significance of the abdominal legs of Panorpid larvæ, 299-The hypermetamorphosis of Mantispa a key to the origin of a complete metamorphosis, 299—The aspect of the Neuroptera comparatively specialized and modern, 299-The Neuropterous labium a secondary product, 299-Origin of the Coleoptera, 299-The free, active larvæ of the carnivorous groups most nearly allied to the primitive form, 300-The scavenger and phytophagous larvæ show increasing degradation of development, 300-The relative form of the maxillæ a good index of the general development of the body in Coleoptera, 300-Structure of maxillæ in the several families, 300—Close resemblance of the mouth-parts of Elaterid

346

[3]

[8]

2267. [RILEY, C. V., et al.]—Continued.

CHAPTER XI-Continued.

larvæ to those of Carabid larvæ, 301-The hypermetamorphosis of Meloidæ furnishes a clew to the probable origin of the different types of Coleopterous larvæ, 301-Brief description of the metamorphoses of Meloe, 301-Of Epicauta, 302-Comparison of the larval stages of Meloids with the different types of Coleopterous larvæ, 302-Origin of the Diptera, Lepidoptera, and Hymenoptera, 303-Presence of temporary abdominal appendages on each segment of Lepidopterous, Coleopterous, and Orthopterous embryos, 304—Description of the structure of Dermatoptera (Forficula), 304—Labia, 308—Larva of Forficula, 308—Orthoptera, 309-Blattariæ, 309-Close relation of Blatta and Termes, 310-Structure of Mantidæ, 310-Blatta the stem-form of the Orthoptera, 312-Mantis connects the Acrydii and the Blattariæ, 312-Structure of Phasmida, 312—Diapheromera, 312—Prisopus, 313— This genus connects the Phasmida with the Acrydii, 314-Structure of Acrydii, 314-Caloptenus, 314-Proscopia, 316-Conocephalus, 318-Locustariæ (Anabrus), 318-Gryllidæ, 319-Gryllus, 319—Gryllotalpa, 320—Œcanthus, 322—Sequence of families of Orthoptera, 322-Structure of Pseudoneuroptera, 322-Corrodentia, 322-Perlide (Pteronarcys), 322-Psocide (Psocus), 325-Termitidæ, 326-Odonata, 329-Ephemerina, 333-Neuroptera, 335—Planipennia, 335—Sialidæ, 335—Hemerobiidæ, 338—Panorpidæ (Panorpa), 342-Trichoptera (Limnophilus), 344.

CHAPTER XII:

Note on the geographical distribution of the Rocky Mountain locust, illustrated with a colored zoö-geographical map of North America. By A. S. Packard, jr......

APPENDICES.

APPENDIX I:

Early references to the occurrence of the Hessian fly in North America.

Early newspaper references, [3]—In 1784-785, at Long Island and
West Chester, [3]—In 1788, in Long Island, Delaware, New Jersey, and Pennsylvania, [3]—The fly resisted by certain varieties of wheat, [3-5].

APPENDIX II:

B. Observations of the new crop gall-gnat. By Dr. Balthasar Wagner, Fulda, 1861. Translated by Carl F. Gissler.....

Literature, [8]—Preliminary consideration of the subimago and imago stages, [8]—Occurrence at Fulda, [8]—Breeding under nets, [9]—Seasonal and sexual differences, [9]—The fly short lived, [10]—Description of the adult, [10]—Of the male, [12]—Of the female, [13]—The winter generation, [14]—The egg, [15]—The maggot, [16]—The pupa or chrysalis, [19]—The perfect insect, [20]—Seasonal and sexual differences, [21]—Only the maggots hibernate, [22]—Single and double brooded, [23]—Summer generation, [23]—The new cereal gall-gnat as compared with Cecidomyia secalina Loew, [24]—Comparison of C. secalina Loew with C. destructor Say, [24]—Views of Loew, Fitch, and Herrick

21 ENT

2267. [RILEY, C. V., et al.]—Continued. APPENDIX II-Continued. discussed, [25]-The rye gall-gnat, C. secalina and C. destructor, all of one species, [28]—The name Hessian fly not justified, [28]— Historical evidence thereon, [28]-Attempt at a new theory of the origin of the insect, [31]-Of Asiatic origin?, [31]-Carried to North America from France or Spain, [32]—Parasites, [33]— Remedies, [34]—The removal of the barley after-growth, [34]— Picking by hand, [35]—Grazing, [35]—Mowing, etc., [36]—Sowing grain as bait, [36]-Plowing, burning, rolling, [37]-Richness of soil, [37]—Resistive varieties of wheat, [38]—Sowing late or early, [38]. APPENDIX III: The Hessian fly in Silesia in 1869. By Professor Dr. Ferd. Cohn.... [39] The Hessian fly, Cecidomyia destructor, and other Diptera observed destroying grain in Silesia, [39]. APPENDIX IV: Koeppen's account of the Hessian fly. By F. T. Koeppen, St. Petersburg, 1880 [41] The Hessian fly recently detected extensively distributed in Russia, [41]-Two generations each season, [41]-Number of eggs at each deposition, [41]-Parasites, [41]-Doubtful species of Cecidomyia, [42]—Descriptions of species, [42]. APPENDIX V: The Hessian fly not imported from Europe. By Dr. H. A. Hagen, [43] Cambridge, Mass.... Conclusions of the author and others as to the nativity of Cecidomyia destructor, [43]—Not imported by the Hessian troops, [43]— Probably here before the war, [45]—Not known in Germany before 1857, [46]-Mistakes corrected, [46]-A critical review of its history, [47]-Marked difference between C. destructor and C. secalina, [49]. APPENDIX VI: Report on the Rocky Mountain locust in 1880. By John Marten... [50] Report to Dr. Cyrus Thomas of observations, [50]—Damages from locusts and wheat flies in Minnesota, [50]—Previous damages from locusts in Dakota, [51]—In Montana, [52]. APPENDIX VII: Report of notes made in 1880. By A. J. Chipman..... No locusts in Kansas, [55]—Suffering from drought, [55]—Favorable climatic changes, [55]—The locusts in Colorado, [56]. APPENDIX VIII: [57] Miscellaneous locust notes..... Dates and localities of locust swarms that have attained the coast of Great Britain. Notes by A. H. Swinton, [57]-Occurrence of the migratory locust in Japan, [59]—Early notices of locusts in

ates and localities of locust swarms that have attained the coast of Great Britain. Notes by A. H. Swinton, [57]—Occurrence of the migratory locust in Japan, [59]—Early notices of locusts in North America, [59]—Locusts at Fort Frontenac or Cataraqui, Canada, [59]; at Puerto Seguro Harbor, California, [60]—The locust in California in 1830, [60]; in Nevada, [60]—Locusts and coffee-trees in Guatemala, [60]—History of locusts in other countries, [61]—Locusts probably never will do so much damage again in North America, [61]—Utilization of locusts as food, [61]; of crickets as manure, [62]—Ravages of locusts in Russia in 1879 and 1880, [62]; in the Caucasus, [63]; in Elizavetopol, [63]; in the district of Rassachs, [64]—Plowing as a means of killing the

2267. [RILEY, C. V., et al.]—Continued.

APPENDIX VIII—Continued.

locusts, [64]—Destruction of locusts in the district of Gori, [64]—Locusts in China in 1878, [65]—Oil as a means of destroying them, [65]-2,000,000 catties of eggs collected, [65]-Article on the extermination of the locust in China, [65-68]—Chinese bibliography of the subject, [65]—Chinese regulations regarding the extermination of locusts, [67]-Locusts in Cape Colony, South Africa, [68]—Ravages of unfledged locusts, [68]—Locusts as food, [69]-Barriers as means of warding off and capturing locusts, [69]—Egg laying and hatching, [72]—Protection against flying locusts, [72]—Locusts in the Philippine Islands in 1878-'79, [72]—The locust plague in Bolivia, [72]—Locusts in India, [73]; in the Ottapidaram Taluq, [73]; in Sattur, [73]—Request of the Madras Government for information as to the movements of the locusts, [74]-Locusts in Kulapurum, Comaralingum, Kolumam. and in the Pulni Taluq, [74]—Difficulties in coping with locusts in southern India, [74]—Means against unfledged locusts, [75]— Locusts driven into the sea at Tuticorin and Trichendur, [76]-Recommendation of rewards for gathering locusts, [76]-Locusts at Peryapatty, [76]; in the Bellary district, [77]-Recommendation of means against locusts, [78]-Locusts at Madura, [78]-Destruction of young locusts by fire at Pothanore, [78]—Locusts at Tiroomoorthypovil, Dhully, and Jellipatty, [79]; in various parts of the Punjab, [79]—Means against the locusts, [79]—Remuneration for destruction of locusts, [80]—Locusts in Cuddapah district, [81].

APPENDIX IX:

Experience with the spring canker-worm. By Martin A. Howelf, jr.. Means employed to save an orchard from the ravages of cankerworms, [62]—Pruning, manuring, and washing of the trees, [62]—Spraying of the leaves with soft soap and arsenite of soda in water, [63]—Belting the trees with sulphur and adhesive mixtures, [63]—Endurance of the moths, [64]—Variation of color in eggs, [65]—Most practical means of relief, [65]—Adulterations of Paris green, [65]—Good force-pumps, [65]—Distribution of canker-worms by the wind, [65].

- 2268. [RILEY, C. V.] The silk industry in the United States. <Sci. ence, 7 March, 1884, v. 3, pp. 290-292. S.-b. No. 42, pp. 82-84-Review of article of W. C. Wyckoff; early history and present status of silk culture in North America; criticism of the estimates of the amount of silk raised in early times; relations of silk-culture and silk-manufacture to free trade and protection in the United States.
- 2269. RILEY, C. V. The army-worm. <Washington, 1883 [20 March, 1884], pp. 89-152, pl. 1-2, with 1 p. expl. of pl. Advance reprint of principal portions: <Rept. [U. S.] Com. Agric. for 1881-1882, January, 1883, pp. 89-106.
 - Chapter 6 of the Third Report U. S. Entomological Commission. See No. 2267 for synopsis of contents.
- 2270. RILEY, C. V. Canker-worms. < Washington, 1883 [20 March, 1884], pp. 157-198, pl. 3, with 1 p. expl. of pl. Extract: < Amer.

- 2270. RILEY, C. V.—Continued.
 - Cult., 1884, v. 46: 29 March, p. 1; 5 April, p. 1. S.-b. No. 42, pp. 118-119; No. 57, pp. 86-87.
 - Chapter 7 of Third Report U. S. Entomological Commission. See No. 2267 for synopsis of contents.
- 2271. [RILEY, C. V.] The chinch-bug. Another entomologist expresses his views regarding the farmers' pest. <Watertown [N. Y.] Daily Times, 27 March, 1884, v. 24. S-b. No. 57, pp. 42-43. Reprint: <Ogdensburg Journal, 1884.
 - Notice of articles of J. A. Lintner; the occurrence of *Blissus leucopterus* in New York State in unusual abundance in 1882 and 1883 not a cause for great alarm; means against this insect. See No. 2252.
- 2272. BILEY, C. V. Notes on North American *Psyllidæ*. < Proc. Biol. Soc. Wash., 1884, v. 2, pp. 67-79. Separate: <[Washington], 10 April, 1884, pp. 67-79.
 - Notice of earlier studies of European and North American Psyllidæ; list of `18 nominal species described from North America; systematic and synonymical list of the same; descriptions of Packypsylla n. g. [p. 71], P. venusta, P. celtidis-mamma, and P. [Blastophysa n. s-g., p. 75] celtidis-gemma n. sp. [p. 74]; dichotomic table of the three species of Packypsylla; descriptions of Ceropsylla n. g. [p. 76] sideroxyli n. sp. [p. 76], Rhinopsylla n. g. [p. 77] schwarsii n. sp. [p. 78].
- 2273. BILEY, C. V. Remarks on the bag-worm, *Thyridopteryx ephemorasformis*. <Proc. Biol. Soc. Wash., 1882–1884, 1884, v. 2, pp. 80–83, figs. 1–3. Separate: <[Washington], 10 April, 1884, pp. 80–83, figs. 1–3.
 - Explanation of the more important and less known facts in relation to the life-history of Thyridopteryx ephemeræformis, especially in reference to the act of coition; figures of all stages and of the external male genitalia.
- 2274. RILEY, C. V. The use of naphthaline as an insecticide. <Science, 11 April, 1884, v. 3, pp. 455–456.
- Review and summary of article of E. Fischer.

 2275. BILEY, C. V. Entomography of *Hirmoneura*. <Science, 18
 - April, 1884, v. 3, p. 488.

 Review of F. Brauer's observations on the life-history of *Hirmoneura obscura*.
- 2276. RILEY, C. V. Orthoptera. <Standard Natural History (The) ...
 Boston, Cassino, 1 May-27 July, 1884, v. 2, pp. 167-203, 1 pl.,
 figs. 243-285. Reprint: <Riverside Natural History (The) ...
 Boston and New York, Houghton, Mifflin & Co., 1888, v. 2, pp.
 167-203, 1 pl., figs. 243-285.
 - Systematic position, limitations, classificatory characters, transformations, molts, and geographical and geological distribution of the order; characters, habits, and distribution of the several families; mention of the principal North American and of some foreign species, with more particular accounts of many species, especially in regard to their stridulation and oviposition, with figures of a few species and of their eggs. Ectobia germanica, Phasmomantis carolina, Diapheromera femorata, Ecanthus niveus, E.latipennis, Orocharis saltator, Camptonotus scudderi, Cyrtophyllus concavus, Microcentrum retinerve, and Caloptenus spretus are particularly mentioned.
- 2277. RILEY, C. V. Reports of observations and experiments in the practical work of the division, made under the direction of the

entomologist, together with extracts from correspondence on miscellaneous insects. <Bull. No. 4, Div. Ent. U. S. Dept. Agric. [3 May], 1884, 192 pp., 4 figs.

CONTENTS.

Introduction	7
Report upon cranberry and hop insects. By John B. Smith	9
Observations on the Rocky Mountain locust and other insects in the	
Northwest during the summer of 1883. By Lawrence Bruner	51
Preliminary report of observations upon insects injurious to cotton,	
orange, and sugar-cane in Brazil. By John C. Branner	63
Report on the effects of cold on the scale insects of the orange in	
Florida. By Joseph Voyle	70
Extracts from correspondence	74
Additional notes on the cultivation of pyrethrum in the United States.	95
RILEY, C. V. Silk-culture in the colonies. <science, 1<="" 9="" td=""><td>fay,</td></science,>	fay,

- 2278. RILEY, C. V. Silk-culture in the colonies. <Science, 9 May, 1884, v. 3, pp. 562-563.
 - Explanation of several technical terms used in silk-culture.
- 2279. RILEY, C. V. Maple-tree insects. <Sci. Amer., 24 May, 1884, v. 59, p. 325. S.-b. No. 63, p. 45.
 - Answer to inquiry of R. H. Cutler; food-plants of and means against Pulvinaria innumerabilis.
- 2280. RILEY, C. V. Acronycta betulæ n. sp. <Bull. Brooklyn Ent. Soc., May, 1884, v. 7, pp. 2-3, fig.
 - Description of Acronycla betulæ n. sp. [p. 2], reared from Betula nigra; seasons and habits of larva; comparison with related species.
- 2281. RILEY, C. V. Recent advances in economic entomology. <Proc. Philos. Soc. Washington, 1884, v. 7, pp. 10-12. Separate: <Washington, 1884, pp. 10-12. Reprint: <Kansas City Review of Science, May, 1884, pp. 13-15. S.-b. No. 63, pp. 43-44. Characteristics of economic entomology as a science and difficulties of its prosecution; the chief insecticide substances and their applicability; recent progress in mechanical appliances.
- 2282. RILEY, C. V. Department of insects. <Ann. Rept. of Regents Smith. Inst. for 1882, 1884, pp. 215-216.
 - List of accessions to the U.S. National Museum, with comments; note on the Glover plates and manuscripts.
- 2283. RILEY, C. V. Quelques mots sur les insecticides aux États Unis et proposition d'un nouveau remède contre le Phylloxera. Communication faite à la Société d'Agriculture de l'Hérault à la seance du 30 Juin, 1884. < Montpelier, 1884, t p. cover+8 pp. Reprint: < Messager Agricole, 10 July, 1884, ser. 3, v. 5, pp. 255-265. < La Vigne Amer., July, 1884, v. 8, No. 7, p. 207.
 - Resistant vines better than the best insecticides as a means against *Phylloxera* vastatrix; principal insecticides in use, especially arsenical substances, petroleum and pyrethrum; method of action and mode of application of these insecticides, especially application by the cyclone nozzle, and the subterranean application of kerosene enulsion as a means against *Phylloxera* vastatrix; tobacco vaper, under favorable conditions, would probably kill this insect.

2284.	[RILEY, C. V.] Remedies for various insects. < Prairie Fari 26 July, 1884, v. 56, p. 470. Sb. No. 59, p. 36. Comparative efficacy and other qualities of several insecticides, espec	
	arsenical substances, tobacco vapor, petroleum emulsions, and pyreth	
2 285.	RILEY, C. V. Habits of Grapholitha olivaceana. <ent. m<="" mo.="" td=""><td></td></ent.>	
	August, 1884, v. 21, p. 67. Grapholitha olivaceana bred from curled tips of Solidage.	
0000		•
2286.	RILEY, C. V. Rhyssa not lignivorous. Science, 28 Novem 1884, v. 4, p. 486.	Der
	Parasitic habits of Rhyssa [= Thalessa] upon Tremex columba.	_
2287.	RILEY, C. V. The apple-root borer. <rural 13="" 1884,="" 43,="" 50.="" 63,="" 831.="" <i="" against="" cember,="" larva="" means="" new-yorker,="" no.="" of="" p.="" sb.="" the="" v.="">Priorus laticollis.</rural>	De
2288.	RILEY, C. V. A new insect injurious to wheat. <bull. bro<="" td=""><td>ook</td></bull.>	ook
2200.	lyn Ent. Soc., December, 1884, v. 7, pp. 111-112. Separa <[Brooklyn, N. Y., December, 1884], p. 111.	
	Description of larva, pupa, and female image of <i>Isosoma grande</i> n. sp. [p. reared from wheat stems in Indiana.	111
2289.	RILEY, C. V. The insects of the year. <science, 1884,="" 26="" 4,="" 565-568.<="" december="" pp.="" td="" v.=""><td>ber,</td></science,>	ber,
	Entomological calendar for each month of the year.	
2290.	RILEY, C. V. On the care of entomological specimens. < ence, 9 January, 1885, v. 5, p. 25.	
	Reply to editorial comments on the care of entomological specimens in National Museum.	the
0001		6 1
4491.	RILEY, C. V. Report of the entomologist. <ann. [u.<="" rept.="" td=""><td>-</td></ann.>	-
	Commissioner Agric. for 1884, 1885, pp. 285-418, 10 pl. Serate: < Washington: 31 January, 1885, pp. 9+pp. 285-41	_
	pp. 9, 10 pl.	
	Introduction	286
	Scope of the report, 285—Climate and insect phenomena during the year, 285—Silk-culture, 286—Field force of the Bureau, 286—Suggested additions to field force, 287—Office force and	
	its work, 287—Trip to Europe, 287—International Forestry Exhibition at Edinburgh, 287—Phylloxera in France, 288—Ex-	
	hibit of the Bureau at the New Orleans Exposition, 288—Third	
	report of the United States Entomological Commission, 288—	
	Special reports, 288—Suggestions, 288—Illustrations, 288.	
	CABBAGE INSECTS	289
	Cabbage cut-worms.	289
	General characters, habits, and natural history, 289—Natural en-	
	emies, 290—Damage to cabbage, 290. The dark-sided cut-worm, Agrotis messoria	290
	Synonymy, 290—Spread, 290—Climbing habits, 290—Natural his-	
	tory, 290—Larva and moth popularly described, 290.	
	The granulated cut-worm, Agrotis annexa	291
	Spread, 291-Number of annual generations, 291-Hibernation,	
	291—Eggs and early states described, 292.	
	The shagreened cut-worm, Agrotis malefida	292
	Distribution, 292—Points of difference from Agrotis annexa, 293— Early states described, 293.	

RILEY, C. V.—Continued. CABBAGE INSECTS—Continued.	
The W-marked cut-worm, Agrotis clandest	lina
Food-plants, 293—Habits, 294—Larva a scribed, 294.	
The greasy cut-worm, Agrotis ypsilon Synonymy, 294—Distribution, 294—Hi 294—Hibernation, 295—Larva and m 295—Description of egg, 295.	abits, 294—Food - plants,
The speckled cut-worm, Mamestra subjunct Distribution, 296—Habits, 296—Appear	
The glassy cut-worm, Hadena devastatrix. Past history, 297—Natural history, 297-	
The variegated cut-worm, Agrotis saucia Distribution, 297—Food-plants, 297—Ej scribed, 298—Rapidity of developmen	gg and early states de-
Remedies for cabbage cut-worms	the stem, 298—Trapping
Other cabbage insects	
The imbricated snout-beetle, <i>Epicarus im</i> Distribution, 300—Food-plants, 300— 301—An allied imported insect, said t	Habits, 301—Remedies,
The wavy-striped flea-beetle, <i>Phyllotreta</i> of Food-plants, 301—Habits, 301—Injury, Remedies, 302.	rituta
Zimmermann's flea-beetle, <i>Phyllotreta sim</i> Food-plants, 304—Habits of larva, 307— with the wavy-striped flea-beetle, 305 306—Other species of the group, 307-	-Characters as compared -Early states described, -Their habits, 307—Nat-
ural enemies, 307—Pleurotropis phyllor The Colorado cabbage flea-beetle, Phyllon How it differs from the preceding spe Pacific coast species, 308.	reta albionicaecies, 308—Mention of a
The Colorado potato-beetle, <i>Doryphora</i> 10 Mentioned as sometimes eating cabbage	
The harlequin cabbage-bug, Murgantia his Distribution, 309—Its spread to the No plants, 310—Natural history, 310—At Natural enemy, 311—Remedies, 311—cultivation, 311—Destroying the earl 312.	rth and East, 309—Food- mount of damage, 311— Applications, 311—Clean
The tarnished plant-bug, Lygus pratensis. Distribution, 312—Food-plants, 312—8 carnivorous habit, 312—Effects of its in color, 313—Habits, 313—As an ener Remedies, 314—Preventive measures,	ynonymy, 312—Supposed puncture, 313—Variation my to strawberries, 314—
The false chinch-bug, Nysius angustatus Synonymy, 315—Food-plants, 316—Hab	
The cabbage plant-louse, Aphis brassics. An imported insect, 317—Damage, 317 tory, 318—Natural enemies, 318—Ren	' Habits, 317Life-his-
The cabbage Anthomyia, Anthomyia brass Introduced from Europe, 319—Life-hi mies, 320—Remedies, 321.	ica

2291. RILEY, C. V.—Continued.	
CABBAGE INSECTS—Continued.	
The cabbage Oscinis, Oscinis brassica	32
First found at St. Louis, 322—Habits, 322—Descriptive, Oscinis	
brassica n. sp., 322.	
The Rocky Mountain locust, Caloptenus spretus	32
Successful introduction of a parasite (Apanteles glomeratus) of the	
imported cabbage-worm	323
GENERAL TRUTHS IN APPLIED ENTOMOLOGY	323
Introductory, 323—Benefits derived from insects, 324—Influence of civilization upon insect increase, 324—Losses from insects, 324—Knowledge which the economic entomologist should possess, 325—Habits of the grape phylloxers, 325—Number of species of insects, 326—The different insecticides, 326—Arsenical compounds, 327—Petroleum, 327—Kerosene emulsions, 328—Pyrethrum, 328—Its cultivation in California, 328—Its	
influence on mammals, 329—Insecticides to be used against	
root-feeding insects, 329—Mechanical contrivances, 329—The	
eddy-chamber or cyclone nozzle, 330.	~~~
Their importance as insecticides, 330—Use in orange groves, 330—Warning against imperfect emulsions, 331—Formula for kerosene and soap emulsions, 331—Importance of a stable emulsion, 331—Use of a force-pump in making emulsions, 331—Trees injured by unemulsified oil, 332—The use of ley washes	330
in California, 333.	
Miscellaneous insects	334
The American Cimbex, Cimbex americana	334
Injury to willows, 334—A new habit, 334—Eggs and mode of oviposition, 335—Habits of young larva, 335—Remedies, 336—Other willow enemies, 336.	٠
The streaked cotton wood leaf-beetle, Lina scripta	336
Value of the cottonwood in the West, 336—Damage by the beetle in 1884, 336—Other food-plants, and former injuries, 337—Natural history, 338—Remedies, 338—Apparatus for poisoning, 339—Varieties of the beetle, 340.	
The Southern buffalo-gnat, Simulium. pecuarum	340
Losses in former years, 340—Damage done by European species, 341—Early states of European species, 342—Early states of American species, 342—Recent ravages in the Southwest, 343—Remedies, 344—Smudges, 344—Body applicatious, 344—Suggestions, 345.	
The angoumois grain-moth, Geleckia cerealella	345
Abundance, 345—Origin, 345—Past history, 346—Natural history and method of work, 346—Number of annual generations, 347—Results of its work, 347—Parasites, 348—Remedies, 348—Baking and concussions, 348—Quarantine, 348—Bisulphide of carbon, 349—Preventive measures, 349—Description of eggs, 350.	
The cottony maple scale, Pulvinaria innumerabilis	350
Distribution, 350—History and synonymy, 350—Life history, 351—Food-plants, 352—Mode of spreading, 353—Natural enemies, 354—Remedies, 355.	
The cranberry fruit-worm, Acrobasis vaccinii	35 5
Natural history, 355—Remedies, 356—Descriptive, Acrobasis vac- cinii n. sp., 356—Systematic position, 357.	

2291.	RILEY, C. V.—Continued.	
	MISCELLANEOUS INSECTS-Continued.	
	The larger wheat-straw Isosoma, Isosoma grande	357
	The phytophagic habits of the genus, 357-Mr. Webster's obser-	
	vations, 357—Descriptive, Isosoma grande n.sp., 358—Oviposition,	
	358—Only females known, 358.	
	SILK CULTURE	359
	Sending out eggs, 359—Reports, 359—The manual of instructions,	
	359-Appointment of Philip Walker as special agent, 359-	
	Diseases, 359—Pasteur's process, 359—A Government reeling	
	establishment, 360-Correspondence, 360-The Serrell reel,	
	360—Home reeling, 361—Herbelin's filature, 361.	
	REPORTS OF AGENTS	36
	Rust of the orange. By H. G. Hubbard	36
	Discoloration of the fruit, 361—Time of appearance, 361—It is	
	not a fungus, 369—The mite on the leaves, 362—First appear-	
	ance of mites on the fruit, 363-Attacks of mites always fol-	
	lowed by rust, 363—Interval between disappearance of mites	
	and appearance of rust, 363—Description of the mite, 363—Life	
	history, 364—Food, 364—Wandering habits, 365—Numerical	
	abundance, 365-Rings of rust on fruit, 366-Influence of	
	weather, 366—Agencies which assist in the distribution of the	
	mites, 366—Food-plants, 367—Effects of attacks on foliage, 367—	
	Rusted fruit, 367—Introduction and spread of the mite, 367—	
	Periods of increase, 368—Geographical distribution, 368—Reme-	
	dies, 368—Influence of soil and methods of cultivation, 368—	
	Preventive measures, 368—Application of insecticides, 369—	
	Whale-oil soap, 369—Sulphur, 370—Kerosene, 372—Creosote,	
	372—Potash, 372—Pyrethrum, 373—Lime, 373—Ashes, 373—	
	· Caution, 373.	
	Second report on the causes of the destruction of the evergreen and	
	other forest trees in northern New England and New York. By A.	
	8. Packard, jr	374
	Localities visited, 374—Destruction of spruces in northern New	
	York, 374—Probably caused by bark-boring beetles, 375—De-	
	struction of spruces in northern Maine, 376—Discussion of	
	causes, 376—Condition of the hackmatack in 1884, 377—The	
	reddish-yellow spruce-bud worm, 378—Its injuries, 378—Its	
•	classification, 378—Its habits, 378—Remedies, 379—The com-	
	mon longicorn pine-borer, 379—Abundance, 379—Mode of ovi-	
	position, 380—Description of egg, 380—Habits and damage	
	done, 381—The sugar-maple borer, 382—Its burrows, 382—Its	
	eggs and larvæ, 382—Kerosene as a remedy, 383. Insects affecting fall wheat. By F. M. Webster	383
	The wheat-straw Isosoma, 383—Oviposition, 384—Description of	30.
	its egg, 385—Number of generations, 385—Effect of its work on	
	wheat-straw, 386—Table o position of larvæ in straw, 386—	
	Natural enemies, 387—Wheat and grass saw-fly, 387—Appear-	
	ance, 388—Parasites, 388—Wheat midge, 389—Natural enemies,	
	389—Dates, 389—American Meromyza, 389—Dates, 390—The	
	flea-like negro-bug, 390—Dates, 390—Description of egg, 390—	
•	The tarnished plant-bug, 391—The soldier-bug, 391—The field-	
	cricket, 391—Cutting off wheat, 391—Insects affecting rye, 392—	
	Insects affecting oats, 392—The stalk-borer, 392—Dates, 393.	

	3 00	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	
	2291.	RILEY, C. V.—Continued.	
		REPORTS OF AGENTS—Continued.	
		Report upon insects affecting the hop and the cranberry. By J. B. Smith	904
		The hop Aphis, 393—Its first appearance, 393—The cranberry	393
		fruit-worm, 394—The fire-worm, 395—Cranberry leaf-folder,	
		396—Its dimorphism, 396—A cecidomyid larva infesting pears	
		at Meriden, Conn., 396—Description of larva and infested fruit.	
		396—Pupation, 397—Distribution, 397—Varieties of pear in-	
		fested, 397—Probably an imported pest, 398—Remedy, 398.	
		Notes from Nebraska. By Lawrence Bruner	396
		The Rocky Mountain locust in 1884, 398—Other species of locusts,	
		399—The snowy tree-cricket, 399—The chinch-bug, 399—Cut-	
		worms, 400—The imported cabbage-worm, 400—Its natural ene-	
•		mies, 400—Other cabbage-worms, 401—False caterpillars on	
		grass and sedge, 401—Cimber americana on willow hedges, 401—	
		Injury to cotton woods by the striped beetle, 402—The Colorado	
		potato-beetle, 403.	
		NOTES OF THE YEAR	403
		Chinch-bug notes	403
		The so-called "invasion" of St. Lawrence County, New York,	
		in 1883, 403—Mr. Lintuer's prediction and his reasons therefor,	
		403—Reasons for dissenting from this view, 404—The result in	
		1884, 405.	
		Notes on the grape Phylloxera	405
		Regarding the grape Phylloxera in Russia, 405—In reference to	
		the treaty of Berne and the prohibition of the introduction of	
		bulbs and cuttings from the United States into Germany, 407—	
		The grape Phylloxera in graperies; legal questions arising, 408.	
	•	Miscellaneous notes	410
		The box Psylla found in the United States, 410—The dwarfing of	
		oaks by Mallodon melanopus, 410—The clover-seed midge, 411—	
		The potato-stalk weevil, 411—The red-humped prominent, 411— The varying anomals, 412—White-lined morning-sphinx, 412—	
		The apple-tree tent caterpillar, 412—Corn bill-bugs, 413—A	
		swarming mite, 413—A new enemy to white roses, 413—A bec-	
		tle eating peach-leaves, 413—Effect of cold on the eggs of bark-	
		lice, 413—Ravages of grain-weevils in Florida, 414—Fuller's	
		rose-beetle, 414—The blood-sucking Conorhinus, 414—The ca-	
		talpa sphinx, 415-Notes on cotton worms, 415-Pyrethrum,	
		416.	
		Notes from Missouri. By M. E. Murtfeldt	416
		The striped flea-beetle, 416—The Colorado potato-beetle, 416—	
		The imported cabbage-butterfly, 416—The cottony maple scale,	
		417—The stalk-borer, 417—Grapholitha prunivora, 418—The	
	•	grape saw-fly, 418—A saw-fly on plum, 418—A saw-fly on ash,	
		418.	
	2292.	RILEY, C. V. General truths in applied entomology. Est	8ay.
			e pa -

rate: < Macon, Ga., 1884, t.·p. cover, pp. 153-159. print: <Rept. [U. S.] Commissioner Agric. for 1884, pp. 323-330. <Gardener's Chronicle, 1885, v. 23; 20 June, pp. 785-786; 27 June, p. 818. S.-b. No. 61, p. 6; 10-11. See: <Ga. Crop

- 2292. RILEY, C. V.—Continued.
 - Rept., June, 1884, p. 33. <Sci. Amer. Suppl., 7 February, 1885, v. 19, p. 7588. S.-b. No. 61, p. 12.
 - See No. 2291 for synopsis of contents.
- 2293. R[ILEY], C. V. Codlin moth. <Suppl. Encyclopædia Britannica, 9th ed., 1884, v. 2, p. 281, fig.

 Natural history, remedies, and figures of Carpocapsa pomonella.
- 2294. B[ILEY], C. V. Colorado potato-beetle. <Suppl. Encyclopædia Britannica, 9th ed., 1884, v. 2. p. 311, fig.

 Summary of the eastward spread of *Doryphora* 10-lineata; natural history, enemies, remedies, and figures of the same.
- 2295. R[ILEY], C. V. Cotton-worm. <Suppl. Encyclopædia Britannica, 9th ed., 1884, v. 2, pp. 432-433, figs.

 History, ravages, life-history, remedies, and figures of Aletia xylina.
- 2296. R[ILEY], C. V. Curculio, plum. <Suppl. Encyclopædia Britannica, 9th ed., 1884, v. 2, p. 494.</p>
 Natural history and means against Conotrachelus nenuphar.
- 2297. RILEY, C. V. Insect named. < Daily Gate City [Keokuk, Iowa], 18 February, 1885. S. b. No. 63, p. 31.

 Brief notes of Olfersia americana.
- 2298. BILEY, C. V. A new remedy for the imported cabbage-worm.

 <Rural New-Yorker, 28 February, 1885, v. 44, p. 132. S.-b.
 No. 63, p. 50. See: <Colman's Rural World, 5 March, 1885,
 v. 38, p. 76. S.-b. No. 63, p. 8. <Prairie Farmer, 7 March,
 1885, v. 57, p. 153. S.-b. No. 61, p. 26. <Amer. Farmer, 1
 April, 1885, v. 4, p. 105. S.-b. No. 67, p. 60. <Sci. Amer., 23
 May, 1885, v. 52, p. 322. S.-b. No. 61, p. 19. <Indiana Farmer,
 30 May, 1885. S.-b. No. 61, p. 22.
 - Ice-cold water applied during the heat of the day causes the death of the larva of *Pieris rapæ*.
- 2299. BILEY, C. V. The collection of insects in the National Museum.

 <Science, 6 March, 1885, v. 5, pp. 188-189. S.-b. No. 61, pp. 7-8.

 Answer to criticisms of C. H. Fernald; the insects in the National Museum cared for by the entomologist of the Department of Agriculture.
- 2300. RILEY, C. V. Ants and aphides. <Rural New-Yorker, 14 March, 1885, v. 44, p. 171. S. b. No. 61, p. 19.</p>
 Answer to inquiry of J. McFarland; habits of ants in relation to plant-lice;
 - subterranean plant-lice attended by ants; ants in northern States beneficial rather than injurious; ground can be cleared of them by injection of bisulphide of carbon or a strong kerosene emulsion.
- 2301. RILEY, C. V. Parasites of the larva of Lachnosterna fusca.

 <Psyche, 1884 [16 March, 1885], v. 4, p. 224.

 Criticism of statements of O. Lugger; larval habits of Tiphia inormata and Rhipiphorus sp.
- 2302. RILEY, C. V. Jumping spiders. <Rural New-Yorker, 11 April, 1885, v. 44, p. 250. S.-b. No. 63, p. 33.

 Answer to inquiry of H. B. S.; habits of Attus sp.

- 2303. RILEY, C. V. Circular No. 9 [U. S. Department of Agriculture, Division of Entomology]. <[Washington:] 1 May, 1885, 1 p., fo., 1 fig. Reprint: <Pacific Rural Press, 16 May, 1885. S.-b. No. 63, p. 27. <Weekly Times-Democrat [New Orleans], 23 May, 1885. S.-b. No. 61, p. 23. <Rural Californian, June. 1885, v. 80, p. 122. <S.-b. No. 61, pp. 20-21. <Sci. Amer. Suppl., 6 June, 1885, v. 19, p. 7859. S.-b. No. 63, pp. 25-26. <Pacific Rural Press, 16 May, 1885, v. 29, p. 469. S.-b. No. 63, p. 27. See: <Farmer's Review, 21 May, 1885, v. 14, p. 362. S.-b. No. 63, p. 29. N. Y. Weekly Sun, 27 May, 1885. S.-b. No. 63, p. 29. N. Engl. Farmer, 16 May, 1885. S.-b. No. 63, p. 29.
 - Conditions under which the Government will buy silk-worm eggs; how to select moths to produce eggs.
- 2304. RILEY, C. V. The imported elm-leaf beetle. Its habits and natural history and means of counteracting its injuries. <Bull. No. 6, Div. Ent. U. S. Dept. Agric. [14 May], 1885, 20 pp., 1 pl., 1 fig.
 - Treats of Galeruca xanthomelena; its habits and natural history; description, figures, remedies, and methods of application of insecticides.
- 2305. RILEY, C. V. Destroying Cicadas: Scurfy apple bark-louse. <Rural New-Yorker, 23 May, 1885, v. 44, p. 353. S.-b. No. 61, p. 24.
 - Answer to inquiries of J. A. K.; means against Cicada [= Tibicen] septendecim and Chionaspis furfurus.
- 2306. RILEY, C. V. [Poisonous spider.] <Rural New-Yorker, 23
 May, 1885, v. 44, p. 354. S.-b. No. 61, p. 23.
 - Reply to statements of Dr. W. H. G.; no spider known whose foot-tracks kill leaves or trees.
- 2307. [RILEY, C. V.] Expected advent of the locust. <Sci. Amer., 23 May, 1885, v. 52, p. 320. S.-b. No. 61, p. 9. <Farmer's Home Journ., 13 June, 1885. S.-b. No. 63, p. 18. <Orange Co. [N. Y.] Farmer, 28 May, 1885. S.-b. No. 63, p. 23. <See: <Sci. Amer., 20 June, 1885, v. 52, p. 389. S.-b. No. 63, p. 28. Simultaneous appearance of a 17-year (Giorda [- Tibicar] sentenderim) and a
 - Simultaneous appearance of a 17-year (Cicada [= Tibicen] septendecim) and a 13-year (C. [= T.] tredecim) brood; localities of the two broods; life-history and habits.
- 2308. RILEY, C. V. Rust of orange. <Rural New-Yorker, 16 May, 1885, v. 44, p. 355, figs. 196, 197. S.-b. No. 61, pp. 18-19, figs. 196-197. Reprint: <Colman's Rural World, 18 June, 1885, v. 38, p. 195, 2 figs. S.-b. No. 61, pp. 31-32, 2 figs.
 - Description, habits, and means against *Phytopius oleivorus*; figures the same and its eggs; figure of a rusty orange.
- 2309. [RILEY, C. V.] Red-ants. <Sci. Amer., 21 May, 1885, v. 52, p. 183. S.-b. No. 66, p. 35.
 - Recommends pyrethrum, kerosene, hot water, and naphthaline as means against red-ants in houses.

- 2310. [RILEY, C. V.] Ridding the ground of cut-worms. <Rural New-Yorker, 30 May, 1885, v. 44, p. 368. S.-b. No. 63, p. 22.

 Answer to inquiry of V. R. R.; means against the larvæ of Agrotidæ.
- 2311. [RILEY, C. V.] An entomological breakfast. <N. Y. Times, 2 June, 1885. S.-b. No. 63, p. 1. See: <Belvidere [Ill.] Standard, 23 June, 1885. S.-b. No. 63, p. 7. Cicadas and Acridida as food.
- 2312. RILEY, C. V. The periodical or seventeen-year Cicada. <Harper's Weekly, 6 June, 1885, v. 29, p. 363, 4 figs. S.-b. No. 61, pp. 24-26, 4 figs.
 - Records and map showing distribution of brood XXII (septendecim) and of brood VII (tredecim); habits, enemies, and figures, with details of structure of Ctcada [= Tibicen] septendecim; figure of twig punctured by the same and of twig healed after the puncture.
- 2313. [RILEY, C. V.] The winged pests of the West. <St. Louis Globe-Democrat, 9 June, 1885. S.-b. No. 61, p. 11. Prediction that the northern States of the Mississippi Valley will escape serious damage from locusts this year; reasons for this opinion; ravages of Campula pellucida in California.
- 2314. BILEY, C. V. The periodical or seventeen year Cicada. < Amer. Grange Bulletin, 11 June, 1885. S.-b., No. 63, pp. 16-17. Chronological record, natural history, and popular names of Cicada [= Tibican] septendecim.
- 2315. RILEY, C. V. The periodical Cicada. An account of Cicada septendecim and its tredecim race, with a chronelogy of all broods known. <Bull. No. 8, Div. Ent. U. S. Dept. Agric., [17 June], 1885, 46 pp., 8 figs. Second edition, 13 July, 1885.
- 2316. RILEY, C. V. Notes on joint worms. <Rural New-Yorker, 20 June, 1885, v. 44, p. 418, figs. 215-218.</p>
 Habits, means against, and figure of Isosoma hordei; criticism of A. J. Cook's views on Isosoma; figures I. tritici and I. grande.
- 2317. RILEY, C. V. Beetles in the corn-fields. < Daily Gate City [Keokuk, Iowa], 23 June, 1885. S.-b. No. 61, p. 17.

 Answer to letters of J. M. Evans and J. M. Schaffer; Agonoderus pallipes injurious to seed corn; recommends soaking the seed corn in Paris green or London purple.
- 2318. RILEY, C. V. Notes on the periodical Cicada. Science, 26 June, 1885, v. 5, pp. 518-521. Reprint: Sci. Amer. Suppl., 27 June, 1885, v.19, pp. 7905-7906. S.-b. No. 61, pp. 47, 48; No. 63, p. 74. Record showing distribution of broad XXII (septendecim) and of broad VII (tredecim) in 1885; the specific value of the different forms; long period of underground development; its life-history; food of the larva, methods of burrowing, and transformations; the Cicada versus civilization.
- 2319. RILEY, C. V. The Chester onion pest. <Orange County [N. Y.]
 Farmer, 2 July, 1885. S.-b. No. 61, pp. 1-2. See: <Rural New
 Yorker, 5 December, 1885, v. 44, p. 829. S.-b. No. 61, p. 151.
 Extract from Rept. U. S. Ent. for 1884; remedies against cabbage cut-worms;

the poisoned ball system and kerosene emulsion effective against the onion

cut-worm; formula for a soap-kerosene emulsion.

- 2320. RILEY, C. V. Premature appearance of the periodical Cicada.

 <Science, 3 July, 1885, v. 6, pp. 3-4.
 - Criticism of L. F. Ward's record of the occurrence and song of Cicads [= Tibicen] septendecim. See No. 2326.
- 2321. RILEY, C. V. Periodical Cicada in Massachusetts. <Science, 3
 July, 1885, v. 6, p. 4.
 - The occurrence of Cicada [= Tibicen] septendecim in southeastern Massachusetts needs confirmation.
- 2322. RILEY, C. V. Destructive insects of the year. <Rural New Yorker, 11 July, 1885, v. 44, p. 464. S.-b. No. 61, p. 8.

 Injuries and means against Agrotis malefida, Anthonomus musculus and Eurycreon rantalis; localities and food-plants of the last.
- 2323. [RILEY, C. V.] [Grasshopper ravages in California.] <Rural New-Yorker, 11 July, 1885, v. 44, p. 470. S.-b.: No. 61, p. 2. Ravages committed by Melanoplus [= Caloptenus] devastator and Caloptenus differentialis; recommends the use of coal-oil pans against them.
- 2324. RILEY, C. V. Pests of the strawberry. <Rural New-Yorker, 18
 July, 1885, v. 44, p. 484. S.-b. No. 61, p. 17.

 Answer to inquiry of J. H. J.; means against Agrotis tricosa, Phoxopteris
 fragariæ, Eccopsis permundana and Emphytus [= Harphiphorus] maculatus.
- 2325. RILEY, C. V. The imported elm-leaf beetle. < Harper's Weekly, 18 July, 1885, v. 29, p. 463, 1 fig. S.-b. No. 61, pp. 21–22, fig. Natural-history descriptions and figures of all stages of Galoruca xanthomelans; means against the same.
- 2326. RILEY, C. V. Premature appearance of the periodical Cicada.

 <Sci. Amer. Suppl., 15 August, 1885, v. 20, p. 8021.

 Criticism of L. F. Ward's record of the occurrence and song of Cicada [=Tibicen] septendecim; an introductory note states that this article was prepared for Science, but that it appeared in that journal in a mutilated and weakened form. See No. 2320.
- 2327. RILEY, C. V. The cyclone nozzle. <Rural New-Yorker, 22 August, 1885, v. 44, p. 567. S.-b. No. 61, p. 347.

 Facts relative to the invention of the cyclone nozzle.
- 2328. RILEY, C. V. A new remedy against destructive locusts. <Rural New-Yorker, 29 August, 1885, v. 44, p. 577. S.-b. No. 61, p. 33.

 Recommends the use of poisoned bait; formula of the same.
- 2329. RILEY, C. V. The influence of climate on Cicada septendecim. <Entom. Amer., August, 1885, v. 1, p. 91.
 - Records the transfers of eggs of brood XXII (septendecim) of Tibicen septendecim to the extreme southern States where no septendecim brood is known to occur, and of brood VII (tredecim) to northern States where no tredecim brood is known to occur.
- 2330. RILEY, C. V. Department of insects. <Ann. Rept. of Regents Smith. Inst. for 1883, 1885, pp. 239-244.
 - Brief mention of work accomplished in the U.S. National Museum; list of accessions to the collection.

- 2331. RILEY, O. V. Insects of the year. <Prairie Farmer, 3 September, 1885, v. 57, p. 567. S.-b. No. 63, p. 37; 134. See: <Entom. Amer., December, 1885, v. 1, pp. 176-177.
 - Ravages of Agrotis messoria, Anthonomus musculus, Eurycreon rantalis, Cooldomyia destructor, and Acrididæ on the Pacific coast.
- 2332. BILEY, C. V. On the parasites of the Hessian fly. <Proc. U. S. Nat. Mus., 14-17 September, 1885, v. 8, pp. 413-422, pl. 23. Abstract: <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, pp. 332-334. <Amer. Nat., November, 1885, v. 19, pp. 1104-1105.
 - Description, synonymy, habits, and figures of Merisus destructor (Say); M. subapterus n. sp. [p. 416, pl. 23, f. 2: p. 333]; Eupelmus allynii French; Tetrastichus productus n. sp. [p. 419, pl. 23, f. 5: p. 333]; and Platygaster herrickii Packard.
- 2333. RILEY, C. V. Enemies of the black-walnut and willow. <Rural New-Yorker, 19 September, 1885, v. 44, p. 632. S.-b. No. 61, p. 35.
 - Habits, description of eggs, larva, and adult of Datana ministra; description of eggs, larva, and adult of Cimbex americana; means against both species.
- 2334. RILEY, C. V. The song-notes of the periodical Cicada. <Science, 25 September, 1885, v. 6, pp. 264-265. Reprint, with additions: <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, pp. 330-332. Translation: <Stett. Ent. Zeit., 1886, jahrg. 47, pp. 158-160. See: <Science, 11 September, 1885, v. 6, p. 225. <Kansas City Review, October, 1885, p. 171.
 - Description of the three prevalent notes of Cicada [= Tibicen] septendecim.
- 2335. RILEY, C. V. The probabilities of locust or "grasshopper" injury in the near future, and a new method of counteracting their injury. <Colman's Rural World, 29 October, 1885, v. 38, p. 348. S.-b. No. 63, pp. 32–33. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, pp. 519–520. <Proc. 6th Meeting Soc. Prom. of Agric. Sci., Manhattan, Kans., 1886, pp. 38–39. See: <Mich. Christian Herald, 3 September, 1885. S.-b. No. 63, p. 39. <Amer. Grange Bull., 5 November, 1885. S.-b. No. 63, p. 34. <Prairie Farmer, 10 October, 1885, v. 57, p. 669. S.-b. No. 61, pp. 34–35.
 - The advance of settlement and cultivation will prevent such widespread injuries of Caloptenus spretus as occurred between 1874 and 1877; use of poisoned baits.
- 2336. RILEY, C. V. The present status and future prospects of silk-culture in the United States. <Entom. Amer., October, 1885, v. 1, pp. 139-140. Reprint: <Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, p. 516. See: <Science, 11 September, 1885, v. 6, p. 236.
 - Abstract; headings under which the subject is discussed; summary of conclusions.

33 6	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.
2337.	RILEY, C. V. The grain moth. <rural 1885,="" 44,="" 56.="" 63,="" 7="" 744.="" against="" and="" asopia="" descriptions,="" farinalis.<="" habits,="" means="" new-yorker,="" no.="" november,="" p.="" ravages;="" sb.="" td="" v.=""></rural>
2338.	[Riley, C. V.] Profits of silk-culture. <rural 1885,="" 26="" 44,="" 50-51.="" 63,="" 885.="" actual="" answer="" cost="" december,="" different="" in="" inquiry="" j.="" m.="" new-yorker,="" no.="" of="" p.="" parts<="" pp.="" producing="" s.;="" sb.="" silk="" td="" to="" v.=""></rural>
•	of the United States.
2339.	RILEY, C. V. Aletia xylina vs. A. argillacea. <entom. 1,="" 161-163.<="" 1885,="" amer.,="" december,="" pp.="" td="" v.=""></entom.>
	Reasons for considering Aletia xylina the correct name for the cotton-worm; advance print from No. 2343.
2340.	[RILEY, C. V.] [Euphanessa mendica, etc.] <entom. 1,="" 170-171,="" 173,="" 176.<="" 1885,="" amer.,="" december,="" pp.="" td="" v.=""></entom.>
	Remarks on papers read at the A. A. A. S.; Euphanessa mendica considered a geometrid; some characters of the Lepidoptera; life-habits of Egerida.
2341.	RILEY, C. V. On the hitherto unknown mode of oviposition in the <i>Carabidæ</i> . Abstract: <proc. 1884,="" 1885,="" 33,="" 538-539.<="" adv.="" amer.="" assoc.="" for="" pp.="" sci.="" td="" v.=""></proc.>
	History of Chalænius impunctifrons traced from egg to imago; eggs laid singly on leaves of trees inclosed in mud or clay.
2342.	RILEY, C. V. [Report of the Edinburgh International Forestry Exhibition.] < Rept. U. S. Dept. Agric. for 1884, 1885, pp. 167-179.
	Brief mention of some of the insect collections exhibited.
2343.	RILEY, C. V. Fourth report of the United States Entomological, Commission, being a revised edition of Bulletin No. 3, and the final report on the cotton-worm, together with a chapter on the
	boll-worm, with maps and illustrations. <washington: 1885,="" 1886],="" 2="" 38+399+147,="" 45="" 64="" [3="" february,="" figs.,="" maps.<="" pl.,="" pp.="" td=""></washington:>
	LETTER OF SUBMITTAL XVII PREFACE XIX INTRODUCTION XXIII
	Chapter I: Classification and nomenclature. Destructiveness
	Popular and scientific names, 1—Most desirable popular name, 1— Different technical names, 1—History of these names, 1—The name Aletia xylina the correct one, 1—Synonomy of the insect,
	2—Classificatory position, 2—Importance of the family Noctui- dæ, 2—Destructiveness of the worm, 2—Causes which increase this destructiveness, 2—Regions where losses are greatest, 2;
	and least, 3—Tabular statement of losses, 3—Previous statements, 3—Statement of loss in 1881, 4. CHAPTER II:

CHAPTER II-Continued.

7-Habits of worm at different stages, 7-Jumping habit, 7-Only known to feed on cotton and one another, 8-Odor of the worm, 8-Migrations of worm, 8-Method of pupation, 8-Formation of cocoon, 8-The worm never burrows in the ground, 8-Distinctive characters of the chrysalis, 9-Duration of chrysalis state, 9-Distinctive characters of the moth, 9-Sexual differences in the moths, 9-Different habits of the moths at night and by day, 10-Their strong flight, 10-Their position at rest, 10-How soon the female begins to lay, 10-Her prolificacy, 10-Food of the moth, 10-It injures fruits, 11-Structure of the tongue, 11-Mention of other Noctaida which injure fruits, 11-Time elapsing from one generation to another, 11-This will average about one month, 12-Time of year when the first worms appear, 12-Former opinions erroneous, 12-Dates of earliest appearance variable, 12-Worms of all sizes found in Florida and southern Georgia in the latter part of March, 12-First worms always few and scattered, 12-They multiply and spread irregularly, 12-Their progress governed by the season and latitude, 13-The third generation often called the first, 13-Number of annual generations, 13-There are at least seven in southern Texas, 13—Gradual progress and succession of broods. 14-The second generation exceptionally very injurious, 14-Why not usually so, 14-Extent of migratory flights of the moths, 15-Probable existence of northern food-plants, 15-Causes and seasons of migration, 15-Behavior of migrating moths, 15-The question of hibernation, 15-Different former opinions and beliefs concerning hibernation, 16-Discussion of these opinions, 16-The chrysalides are killed by a temperature below 22° F., 16-Parasited chrysalides can bear greater cold, 16-Fallacy of the belief that the chrysalis winters underground, 16-Ease with which erroneous conclusions can be drawn from mistaken identity, 17-The chrysalis of Aspila virescens mistaken for that of the Aletia, 17-Chrysalides of other cotton larvæ found underground in abundance, 17-The chrysalis of Aletia killed by burial, 17-Ability of the moth to survive the winter, 18-Unreliability of most testimony as to the hibernation of the moth, 18-Other moths easily mistaken for it, 18-Hypena scabralis, 18-Phoberia atomaris, 19-Leucania unipunota, 19—Absence of testimony to the survival of the moths beyond March, 19-Theory of annual introduction of the species from some southern foreign country, 19-Statement and discussion of Grote's arguments and of others in support of this theory, 20-Arguments in favor of the hibernation of the moth, 21-Both immigration and hibernation may occur, 21-Summary of the evidence, 22-Aletia hibernates only as a moth and only in the extreme south, especially in Texas, 22.

CHAPTER III:

338	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.
2343.	RILEY, C. V.—Continued.
	CHAPTER III—Continued.
	remedies, 35, 36-Arsenic, 36-Paris green, 36-38-Planting jute,
	38-Machinery, 38-London purple, 38-Pyrethrum, 38.
	Chapter IV:
	The cotton-worm in other countries.
	Geographical distribution of Aletia xylina, 39—Other insects injurious to cotton in the Eastern Hemisphere, 39—Occurrence of Aletia xylina on the Pacific coast of Mexico, 39-40—On the Gulf coast of Mexico, 40-41—In Yucatan, 41—In the West Indies, 42—In the northern countries of South America, 42-43—In Brazil, 43-44.
	Chapter V:
	On the anatomy of Aletia. By Edward Burgess and C. S. Minot
	Circumstances under which this chapter was prepared, 45—External anatomy of larva, 45—True legs, 45—Prolegs, 46—Colored markings, 46—Stigmata, 46—Internal anatomy of larva, 47—Head, 47—Ganglia, 47—Digestive canal, 47—Malpighian vessels, 47—Salivary glands, 47—Dorsal vessel, 47—Stomach, 48—External anatomy of imago, 48—Method of preparing the exceleton for examination, 48—Head and appendages, 48—Prothorax, 48, 49—Mesothorax, 49—Metathorax, 49, 50—Supposed organ of hearing, 50—Abdomen, 50–51—Spiracles, 51—Legs, 51—Scales, 51–52—Scale-pores, 52—Proboscis, 52—Spines, 53—Internal anatomy of imago, 53—Digestive canal, 53—Pharynx, 53–54—Salivary glands, 54—Esophagus, 54—Food reservoir, 54–55—Stomach, 55—Malpighian vessels, 55—Intestines, 55—Aorta, 55—Nervous system, 55—Ganglia, 55—Terminal body segments and organs of reproduction, 56—Brush-sac, 56—Mále organs of reproduction, 56—Testes, 57—Vasa deferentia, 57—Penis, 57—Female organs of reproduction, 57—Ovaries, 57—Sebaceous glands, 57—Vagina, 58—Copulatory pouch, 58—Terminal abdominal segments of the female, 58.
	CHAPTER VI:
	The cotton belt. By Prof. E. A. Smith
	A. General features of the cotton States
	B. Description of the agricultural subdivisions
	The alluvial region, 67-68—The lower prairie region, 68-69—The long-leaf pine region, 69-71—The oak uplands region, 72-73—The upper prairie region, 73-75—The red and brown loam region, 75-77—The sandy and siliceous lands of the older formations, 77-79—The gneissic region, 79-80.
	CHAPTED VII.

Terrestrial and meteorological influences affecting the worm...... Condition of soil and plant connected with the appearance of the first worms, 81-The earliest worms are confined to the "low lands," 81; and to luxuriant plants, 82; and to the vicinity of winter shelters, 82; and to the same localities, 82-Influence of wet weather on the development of the worms, 83-Severe rains with gales destroy both worms and crop, 83-Late cold rains do the same, 84-Frequent summer rains favor the development of the worms, 84-Hot, dry weather destructive to them, 84-Artifi59

67

CHAPTER VII-Continued.

cial drought produces the same effect, 84—Indirect influences of wet weather in favoring the development of the worms, 85—Immunity of the worms from their enemies in wet weather, 25—Drowning of ants by heavy showers, 85—Wet weather prevents poisoning and working the cotton, 85—Letter from J. W. Du Bose on the influence of winds on Alstia, 85–86.

CHAPTER VIII:

Natural enemies General remarks, 87-Importance lessened by the use of arsenical poisons, 87—Vertebrate encinies. 87—Quadrupeds, 87—Birds, 87— English sparrow, 88-Toads and lizards, 89-Invertebrates, 89-Spiders, 89-Ants, 90-Species of ants destroying Aletia, 90-Hubbard's observations on ants, 92-The leaf-cutting ant, 94-Wasps, 94—Colcoptera, 95—Tiger-beetles, 95—Ground-beetles, 95—Lady-birds, 96—Soldier-beetles, 96—Heteropters, 97—List of species, 97—Diptera, 99—Orthoptera, 99—Neuroptera, 100—Lepidoptera, 100-Parasites, 101-List of species, 101-The Trichogramma egg-parasite, 102-The cotton-worm Microgaster, 104-Comstock's Euplectrus, 105-Elachistus suplectri, a secondary parasite, 106-The common flesh-fly, 107-Cyrtoneura stabulans, 108—Is it parasitic?, 108—Tachina-flies, 109—Their habits, 109— The watchful Pimpla, 111—The ring-legged Pimpla, 113—Cryptus nuncius, 113-The ovate Chalcis, 114-The devouring Tetrastichus, 115-Species that are easily mistaken for parasites of

CHAPTER IX:

Preventive measures

Aletia, 115—Hexaplasta zigzag, 115—Phora aletia, 116.

Mode of cultivation, 120—Improving cotton seed, 120—Forcing the young plants, 120—Transplanting from hot-beds, 120—Objections, 121—Frequent cultivation, 121—Topping the cotton, 121—Fertilizers, 121—Sulphuric acid on seeds, 121—Late planting, 121—"Worm-proof cotton," 121—Diversified agriculture, 122—Rotation of crops, 122—Jute as a protection, 122—Other supposed protective plants, 123—Protection of natural enemies, 123—Immunity of cotton under trees, 124—Preventing oviposition of the moth, 124—Futility of decoctions for this purpose, 125—Road dust, 125—Early application of direct remedies, 126—Concerted action, 126—Early poisoning and hand-picking, 126—Cotton-worm warnings, 127—Destroying chrysalides accidentally carried into gin-houses, 127—False theories, 127—Burning the stalks, 127—Winter plowing, 127—Salt, 127.

CHAPTER X:

Remedies; means of coping with the insect; substances used for its destruction

Destruction of the eggs, chrysalides, and moths, 128—Destruction of the eggs impracticable, 128—Little chance for successful destruction of the chrysalides, 128—Destruction of the moth, 129—Lights and fires for attracting the moths, 129—Indifferent success with lamps used at Columbus, Tex., 130—Importance of using lamps early in the season, 130—Apparent success with lamps near Hearne, Tex., 131—Great attractiveness of the electric light, 131—Movable lights, 131—Poisoned sweets and fluids, 131—Fondness of the moths for ripe fruit, 132—Killing moths by poisoned fruit,

87

120

CHAPTER X-Continued.

132-Method of using poisoned liquids, 132-No results from using poisoned baits late in summer, 133-Poisoning the glands of the plant, 133—Cotton-leaf essence and its attraction to the moth, 134—Hand-picking, 135—Mechanical means of killing the worms, 136—Shaking off the worms, 136—Poisoning the worms, 136— Progress in the use of insecticides, 137—Classification of insecticides, 137-Importance of preparing materials in advance, 138-Arsenical compounds, 138—Safety in their use, 138—Difficulty in determining minimum quantities, 139—Overdoses of poisons, 139 -General rules in dry applications, 140-Mixing devices, 141-Ingredients, 141—Wet application, 142—Principles to be followed and ingredients to be used, 142-Comparison of dry and wet applications, 142-Paris green, 143-Historical data, 143-Liquid application, 143—Dry application, 144—Minimum quantities, 144 -Patents on Paris-green combinations, 146-Arsenic, 147-Commercial arsenic, 147—Arseniate of soda, 147—Fowler's solution, 148—Johnson's dead-shot, 148—Texas cotton-worm destroyer, 149 -London purple, 149-Manufacture and analysis, 149-History of its use, 150—Experience in Alabama in 1880, 150—Advantages and disadvantages, 151-Dry application, 151-Wet application, 152-Other mineral substances, 153-Salt and saltpeter, 153-Sulphur, 154—Red lead, 154—Road dust, 154—Oils and allied substances, 155—Kerosene, 155—Former methods of application, 155 -Invention and perfection of emulsions, 156-Method of preparing emulsions, 157—Formula for perfected emulsion, 158—Experiments with imperfect emulsion in 1890, 158-Experiments with perfected emulsions, 160-Oil of creosote, 162-Oil of tar, 162-Gas-tar water, 162-Carbolic acid, 163-Cotton-seed oil, 163-Vegetable insecticides, 164—Pyrethrum, 164—History, 164—Mode of cultivation, 165-Success in cultivating the plant in America, 166-Preparation of the powder, 167-Its use as an insecticide, 168-Advantages and disadvantages, 168-Active principle in Pyrethrum, 169-Its effects on cotton-worms, 169-Imported vs. Californian powder, 170-Dry application, 170-Minimum quantities of dry powder, 171-Dry powder mixed with flour and other ingredients, 172-Application in fumes, 174-Alcoholic extract, 174-Experiments with extract obtained by distillation, 174-Experience with extract obtained by repercolation, 176-Use of the powder in simple water solution, 177-Tea or decoction of Pyrethrum, 178-Effect of Pyrethrum upon other insects, 179-Prospects for the use of Pyrethrum for the cotton-worm, 180-Ox-eye daisy powder and its uselessness as an insecticide, 180-Extracts and decoctions from various plants, 181-Difficulties in the way of discovering new vegetable insecticides, 182-Mode of preparing the extracts and diffusions, 183-List of the plants experimented with, 184-Effect of alcohol upon the worms, 187-Yeast ferment and fungus infection, 188-Dr. Hagen's recommendations of the use of yeast as an insecticide, 188-Objections to Dr. Hagen's plan, 189-Negative results obtained by the Commission, 190.

CHAPTER XI:

Machinery and devices for the destruction of the worm. By W. S. Barnard, Ph. D

CHAPTER XI-Continued.

- I. Spray nozzles, classification, preferred kinds, 191-Many-punctwred nossles, 191-196-Preferred construction, 191-193-Straining and cleaning vs. clogging, 192-Eddy-roses, their operation, construction, and leading importance, 192-193-Plug-roses, action and construction unsatisfactory, 193-194-Johnson's, Melcher's, Dawson's, Foster's, 193-Lynch's, 194-Colliding jets, gas-jets, superiority, Daughtrey's, Weber's, Prouty's, 194-T-roses, 194-Yeager's, Warner's, improvements, 195-Divided rose-heads, clutch-head of Mast, Foos & Co., Foss', Fox's, Barrows', Vose's, 195—Peripheral roses [divided], Melcher's, Yeager's, Ruhmann's, 196-Rose-combinations of Barry, Prentice, etc., 196-Slot nozzles, 196-205-Operation, 196-Preferred construction, 196-201-Disadvantages, improvements made, 197-201—Eddy-chambered, 197-199-Lip construction, 198-199-Inside cleaner, 199-Simple slotnozzles, Fowler's, Mallory's, Iske's, 201-202-Plug slot-nozzles, Allen's, Ruhmann's, Johnson's, "The Niagara," Pinter's, 202-Removable slots of Long, Vestal, and Merigot, 203-Jawed slots, "The Boss Nozzle," Raymond's and Perkin's, Smith's, Moffet's, 203—Williams', Ruhmann's, Pinter's, 204—Stanton's, 205—Side slots, Schier's, Melcher's, 205-Deflector nozzles, 206-211-Definition, use, applicability, relative merits, 206-Drip-waste, clogging, 206-Simple constructions made, 206-208-Conformations for narrow, wide, and even sprays, 206-207-Removable deflectors, 207-Hollings', Douglas's, Nickerson's, 208-Hayden's, Killam's, Lewis', Schier's, Barrett's, 209-Ruhmann's, Binkley's, Schier's compound, 210—Schier's and Polansky's, 211—Centrifugal nozzles, 211-221-Operation, kinds, choice, 211-The new and most perfect sprayers, 212—Eddy-chambered, 212-219—Conformation and operation, 212-Clogging, cleaning, construction principles, 213—Forms described, 214—Whistle-jets, 215-216—For blast-atomizers, 216-Eddy-jets proper, 216-219-Involute form, cone form, 216-Convex and concave forms, 217-Direct discharge, proximal diagonal discharge, 217—Distal diagonal discharge, 218-Centrifugal nose-pieces, double-cone or doublechambered form, and chambered plug form, 218-Direct spray and solid jet, 219-Fistular spray nozzles, hose-pipes, 219-With rotary segment, Hotz's, Clifford's, Gielow's, Johnson's, Hoyer's, Clarke's, 220-With cross-plug, McGaffey's, Johnson's, Gray's, Gielow's, Hosford's, 221-Spray-wheels, 221.
- II. Centrifugal throwers, 221-226—Their character and operation, 221—Rotated orifices, pronged reels, brush poison-throwers, 222—Brush fibers, 222-223—Rotary polishing brushes, improved construction, fiber strength, spring and density, 223—Feeding the brushes with liquid, 223-225—With powder, 224—A simple brush thrower of powder, 224-225—Advantages over-sifters, velocity of rotation, 225—Wisewell's, 225-226—Whisps and brooms, rotated recesses, 226.
- III. Blowers of poison, 226-252—Rotary blowers, 226-235—Kinds and success of, 226-227—For blowing powder, 227-232—Hoppers and adjustable feeders, 227-228—Blower encasements and blast-pipes, 228-229—Blast forks and deflectors, 229—Improved light rotary blowers, 229-230—Hauled blowers, compound rotary blower, Aframe and legged swivel wheels, 230—Rotary velocity, 231—

CHAPTER XI-Continued.

Hurd's rotary powder blower, 231-232—For blowing liquids, 232-235—Improved feeders of liquid to rotary blower, 232-233—Dripcatcher, forked blast spray, 233-Darnell's rotary liquid-blower, 233-234-Hurd's ditto, and Perl's rotary fume-blower, 234-Force-blast rotary blowers-Oscillating blowers, 235-251-Bellows power, construction, durability, improvements, 235-236-For blowing powders, 236-243—Hermetic powder-box, 236—Feeders of powder to blasts, 236-237-Pipes, forks, and nozzles for these powder blasts, 237-238-Hauled compound bellows powderer, to wagon, with motor, etc., 238-Cultivator bellows-blower, 238-939-Knapsack and horse-back bellows powderers, 239-Improved small hand-bellows powderers, 239-241-With forks, 241-Allen's powderer, 241-Common powder bellows, Woodsson's, Hendley's improved, 242—Stelle's, etc., 243—For blowing fluids, 243-249—Importance, resuction, feeding by blast-suction, blast-pressure, gravitation, gauges, 243-244-Blast spray conductors and nozzles, reverberatory, 244-Whistle-jets and agitation chambers, 245—Common blast atomizers, 245—Improved automatic-feeding blast sprayers, 246-249-Reatomizing, reverberatory nozzles, 247-Nether blast sprays, directing pipes and nozzles, 248—Compound combinations, 249--Peck's blast sprayer, and Wallace's, 249—Reciprocating or pistoned blowers, 249-251— Improved air-pump apparatus, 249-250-Air-pumps of Humphryville, Rumsey, etc., 251—Generator blowers, 251-259—Steinmann's vaporizer, 251.

CHAPTER XII:

Machinery and devices for the destruction of the worm—continued. IV. Pneumatic compression squirters, 253-261—Kinds, advantages, available fire-extinguishers, 253-254—Carbonic anhydride, apparatus, practicability, 255—Danger, antidotes, safety constructions, 255-256—Simple generators, 257—Rotary force-blast, compression ejectors, 257—Cecillating bellows, pneumatic compression ejectors, 257—Reciprocating or pistoned ditto, 258-261—Simple kind to construct, beer forcers applicable, Worswick Co.'s, Weindel's, Rumsey's, Douglas's, 258—Daughtrey's underspray theory and machine, 258-261—The author's devices, 259—Weber's nozzles, 260.

V. Solid compression squirters of poison, 261-283—Kinds, 261— Rotary force-pumps, 261-262—Hydraulie bellows, 262-Oscillating force-pumps, 262—Reciprocating force-pumps, 262-283—Kinds, 262— Hydronettes and fountain pumps, 262-269-May's patent, Tyler's, Servants', Staples', Whitman's fountain pump, 263—Hydronettes, double-acting, Deakin's, Rumsey's, cost, 264-Author's improvements in knapsack apparatus, 264-265—In knapsack fireextinguishers, Condict's and Doty's, 265-In horseback apparatus, Warner's, author's, 265-In cart or wagon, 265-268-Calahan's carts, legged wheels, tongue or shafts, 266—Wagon use, Trelease's report, 266-268-More economic devices, 268-Other pumps not in trade, 268-269-Aquapults, aquajects, aquarius, excelsior, hydropult, 269-271-Operation, cost, 269-Aquapult, Johnson's patents, Douglas', Prouty's, 269-270-Johnson's syringe, Douglas' aquarius, Rumsey's aquaject, 270 — Deakin's excelsior, Vose's hydropult, 271-Bucket pumps and knapsack

CHAPTER XII-Continued.

pumps, 271-274-Lewis's, 271-Korth's and less desirable kinds, Stoner's, Mallory's Kaiser's, Dix's, 272-Crandal's, Holland's knapsack extinguishers, Douglas's, Stanton's, 273-Bucket poisoners, McDonald's, Allen's watering-pot improvements, 273-Amor's and Lane's can-syringe, Hull's, Wisner's, 274-Barrel and tank pumps and appurtenances, 274-283—Single acting, discharging below the piston, Melcher's, 274—Voglesang's, Ruhmann's, 275— Polansky's, Schier's, Butman's, Ball's agitator, 276-Evenden's, Helmecke's, 277—Yeager's pump and nozzle, 277-278—Pinter's, 278—Single-acting, discharging from above the piston, Chipley's, Weith's, 278—Available cistern or well pumps, 278-279—Barrow pumps, truck pumps, garden engine pumps, windmill pumps, Blunt's Lotus pump, the Pendulum and Index pumps, counter pumps, 279-Double-acting, force-pumps proper, 279-283-The best, 279-Kinds characterized, Vose's hydropult, the Champion, 280-Ramsden's, reduction-cylinder pumps, patentees and manufacturers, 281-282-Author's agitator barrel pump, found most satisfactory, 282-283-Conduits, frames, portage, and combinations of appurtenances, 283-Johnson's cotton-spraying machine, 284-Jones', Binkley's, '285—Goodin's, Wolfram's, 286—Author's underspraying accessories, 288-Plan, skid, mixing, and straining funnel, 288-Adjustments of pipes, forks and nozzles, 289-297; fork modifications, 290-292-Pendent pipes, 292, conformability, lightness, cheapness, 292-293-Tube substances, 293-Author's Aframe machines, 293-297—Maximum width undersprayed, 295— Pipe adjustments to row widths, flexible systems superior, 296-

VI. Gravitational distributors, 297-309—For liquid, 297-302—Kinds, author's tripod automatic sprinkler, 297-298—Schank's sprinkler, Taylor's, 298—Robinson's, 299—Suction force-pumps and windlass elevators, 299—Bilge-pumps, 300—Horseback automatic sprinklers, watering-pot method, Willie's sprinkler, 300—Ramsey's, 301—Knapsack automatic sprinklers, 301—Gray's, Ruggle's, Townsend's, 302—Automatic hand-sprinklers, watering-pots afoot and on horseback, 302—For dry poisons, sifters, kinds, 302-309—Disadvantages, reciprocating sieve machines, hand-sieves, 303-304—Sifting bag, Hurd's sifter and blower, Goodheart's duster and sprinkler, 304—Rotary-sieve machines, 304-307—Robinson's duster and sprinkler, Davis' duster, Levy's, 305—Taylor's duster and sprinkler, 306—Reciprocating-stirrer sifters, Willie's, 307—Rotary-stirrer sifters, Young's, 307—Smith's, 308—Eldridge's, 309.

CHAPTER XIII:

Machinery and devices for the destruction of the worm—concluded.

VII. Insect manipulators, mechanical treatment, dislodging, crushing or stifting the worms or chrysalids, 310—Sweeping, knocking, or jarring off the worms; friction drags, fringes; beaters, 310—Collecting and despatching means, 311—Crushing, Helm's sweeper and crusher, 311—Ewing's sweeper and stifler, 312—moths, Wood-Smith's, 313—Iske's catching trays, 314—Trapsforthe 314-321—Kinds of lures, lighttraps, 314—Lewis's, McQueen's, Rigel's, 315—Walker's and others, 315-316—Lamps in motion, Leblanc's, Fordtran's, 316—Bait traps, Heard's, Garrett's, 317—Traps combining light and bait, author's net trap, 317-319, thought

CHAPTER XIII-Continued.

the best, 318—Indiscriminate killing wrong, the best bait, 319—Stith's trap, 319—Pugh's, Garrett's, 320—Binkley's, 321.

322

355

CHAPTER XIV:

CHAPTER XV:

Insects liable to be mistaken for Aletia..... Confusion of Aletia with other moths, 345—With Aspila virescens, 345—Characters of the most important of these moths to be illustrated in this chapter, 345—List of these moths, 345—Account of Anomis erosa Hübner, 345—Its geographical distribution, 346— Structure of its eggs, 346—Distinguished from egg of Aletia xylina. 346—Its seasons, 346—Habits of larva, 346—Characters of moth, 347—Detailed description of egg, 348—Larva, 348—Pupa, 349— Seasons and food-plants, 349—Account of Anomis texana n. sp., 350-Habitat of A. exacta, 350-Of A. texana, 350-Larva distinguished from that of Aletia xylina, 350-Pupa distinguished, 350—Account of Leucania unipuncia Haworth, 350—Reference to discussions about this species, 350—Its geographical distribution, 351-Oviposition, 351-Food-habits of worms, 351-Pupation, 351 -Number of broods, 351-Hibernation, 351-Account of Aspile virescens, 351—Synonymy, 351—Confounded with Aletia xylina only in pupa state, 351-Pupæ of the two species distinguished, 352-Food-plants of A. virescens, 352-Moth distinguished from that of Aletia xylina, 352—Account of Drasteria erecktea (Cramer), 352—Its geographical distribution, 352—Variations in size of moth, 352—Food-plants of larva, 352—Habits of moth, 352—Number of broods, 352—Oviposition, 353—Colors of larva, 353—Account of Laphygma frugiperda, 353—Its food-habits, 353—Synonyms, 353—Account of Platyhypena scabra (Fabr.), 354—Geographical distribution, 354—Food-plants, 354—Number of broods, 354 -Reference to descriptions of larva, 354-Euplectrus platyhypenæ Howard, bred from larva, 354-Account of Phoberia atomarie (Hübner), 354—Reference to descriptions and figures of moth,

CHAPTER XVI:

354—Food-plant of moth, 354.

The boll-worm (Heliothis armigera Hübner)

Introductory, 355—One of the foremost of our injurious insects, 355—Extent of its ravages, 355—Nomenclature, 357—Synonyme, 358—Popular names, 358—Geographical distribution, 358—Foodplants, other than cotton, 359—Corn, 359—Tomato, 361—Tobacco and other Solanaceæ, 362—Leguminosæ, 362—Cucurbitaceæ, 363—Malvaceæ, 363—Other food-plants, 363—Characters and transformations, 364—The egg, 364—The larva, 365—The pupa, 370—The imago, 371—Number of broods, 372—Hibernation, 373—Summary of the distinguishing points compared with Aletia, 374—Egg, 374—Larva, 374—Pupa, 374—Adult, 374—Natural enemies, 375—Remedies, 377—Early planting, 378—Low corn vs. high corn, 378—Fall plowing, 378—Destruction of the moths, 379—Lights

1 0040 Dryme C V Continued	
2343. RILEY, C. V.—Continued.	
CHAPTER XVI—Continued.	
and poisoned sweets, 379—Hand-picking, 380—Poisoning, 381— Pyrethrum, 381—Bibliography, 382.	
EXPLANATION TO PLATES	385
Plate.I-III, 385—IV-VII, 386—VIII-XI, 387—XII-XVI, 388—XVII-	•••
XX, 389—XXI-XXV, 390—XXVI-XXVIII, 391—XXIX-XXXII,	
392—XXXIII-XXXVIII, 393—XXXIX-XLII, 394—XLIII-XLVII,	
395—XLVIII-LIII, 396—LIV-LVII, 397—LVIII-LXI, 398—	
LXII-LXIV, 399.	
APPENDICES	[1]
Preface to appendices	[3]
APPENDIX I:	
Report of H. G. Hubbard	[5]
Report of observations made in 1881 upon Aletia and other insect	
enemies of cotton, in the State of Florida, [5]—Cotton injured	
more by Dysderous suturellus than by Aletia xylina, [6]-And	
more by rust-mite than by either, [7]—Great destruction of	
Aletia eggs by Trickogramma pretiosa, [7]—Successions of broods	
of Aletia, [7]—Distribution of worms on plant, [8]—Tabular	
statement of periods of time occupied in each stage of growth,	
[10]—Locality of deposition of eggs on plant [12]—Proportions	
of light and dark colored worms, [12]—Effect of shade in protect-	
ing cotton, [12]—Periods of different stages, [13]—Copulation,	
[13]—Position of moths at rest, [13]—Experiments with poisons,	
[14]—Pyrethrum, [14]—London purple, [15]—Poisoning the	
moths, [15]—Yeast, [16]—The boll-worm, Heliothie armigera	
Hübn., [16].	
APPENDIX II:	
Report of Prof. R. W. Jones	[17]
The cotton boll-worm, Heliothie armigera, [17]—Its importance, [17]	
-Food-plants, [17]-Natural history, [18]-The egg, [18]-The	
larva, [18]—Chrysalis, [19]—The moth, [19]—Weather, [19]—	
Natural enemies, [19]—Cotton army-worm, Aletia xylina, [19]—	
First appearance observed in 1880, [19]—Experiments with	
poisoned sweets, [20]—With adhesive sweets, [20]—Gradual dis-	
appearance of moths in October, [20]—Preparation of vegetable	
substances as insecticides, [20]—Pyrethrum, [21]—Experiments	
with Pyrethrum, [22].	
APPENDIX III:	
Report of J. P. Stelle	[25]
First appearance of Aletia xylina and Heliothie armigera in Texas	
in 1880, [25]—Cotton blight, [25]—Its symptoms, [25]—Its	
causes, [26]—Boll rot, [26]—Its symptoms, [27]—Its cause [27]	
—The flare, [27]—Its symptoms, [27]—Its probable cause, [27]	
-Other food-plants than cotton for Aletia, [27]-None found for	
the larva, [27]—Many for the moth, [27]—Food of larva of	
Heliothie armigera, [28]—Annoyances to the cotton-worm, [28]—	
Common salt, [28]—Saltpeter, [28]—Road dust, [28]—Open	
spaces, [28]—Trees, [29]—Shade, [29]—Natural enemies, [29]—	
Birds, [29]—Ants, [29]—Other insects, [30]—The yeast forment	
remedy, [31]—Pyrethrum powder, [31]—Other vegetable poisons,	
[32]—Arsenical poisons, [32]—London purple, [32]—Paris green,	
[33]—Arsenic, [33].	

[37]

[55]

[59]

2343. RILEY, C. V.—Continued.

APPENDIX IV:

Reports of Dr. E. H. Anderson For 1880, [37]—Seasons and habits of A. xylina, [37]—Of Heliothis armigera, [37]—Localities of first appearance of A. xylins in the season, [37]—Reasons for this first appearance, [37]-[38]—Hibernation of this insect in some form, [38]—Causes of undue multiplication of the worms, [38]-Influence of ants on the number of worms, [38]—Hahits of ants, [38]—Other insects gathering sweets from the cotton-plant, [39]—Jute growing amidst cotton has no effect on the worms, [39]—Saturnia io found eating cotton, [39]—An enemy of the cotton-worm found, [39]—Experiments with London purple, [39]—With Pyrethrum extract, [39] -With yeast, [40]-Eggs of insects found on costen-leaf, [40]-Characteristics of season of 1860, [40]—Report for 1881, [40]— Diary of observations on issuance, egg-laying, and hibernation of moths, and on weather, from October 3 to December 31, [40]-[44]—Observations of Heliothis armigers, [41]-[43]—Report for 1882, [44]—Observations on auts and aphides, [45]—Abundance and sudden disappearance of Laphygma frugiperda, [45]—Scarcity of cotton-worms, [45]—Experiments with Pyrethrum, [45]-[48]—Weather in September, [46]-[47]—Scarcity of Heliothis armigera in cotton, [46]-[47]—Its preferred food-plants, [47].

APPENDIX V:

APPENDIX VI:

APPENDIX VII:

Reports of consuls and consular agents on the cotton crop and its

2343.	RILEY, O. V.—Continued.	
	APPENDIX VII-Continued.	
	injurious to the cotton-plant in the Republic of Mexico, by D.	
	H. Strother, U. S. consul-general, [70].	
	APPENDIX VIII:	
	Answers to Circular No. 7	[71]
	From D. M. Hamilton, St. Francisville, West Feliciana Parish, La.,	
	[71]—From R. A. Lee, Evergreen, Ala., [73]—From J. M. Wol-	
	kom, Henderson, Tex., [74]—From P. S. Clarke, Hempstead,	
	Waller Co., Tex., [75]—From L. D. Hoyt, Livingston, Sumter	
	Co., Ala., [77]—From F. S. Shielda, Lake Concordia, Concordia	
	Parish, La., [79]—From G. E. Gillespie, M. D., Natchitoches,	
	La., [81]—From H. O. Dixon, Jackson, Miss., [84]—From F. L. Yoakum, Larissa, Cherokee Co., Tex., [85]—From O. H. Perry,	
	Perry Co., Ala., [86]—From J. W. Grace, Walterborough, Colle-	
	ton Co., S. C., [86]—From F. M. McMeekin, Jamestown, Alachus	
•	Co., Fla., [87]—From H. P. Bee, San Antonio, Tex., [88]–[89]—	
	From E. H. Anderson, Kirkwood, Miss., [90]—Condensed sum-	
	mary of the habits of the worm, from Dr. D. L. Phares, Wood-	
	ville, Miss., [92].	
	Notes	[93]
	Note 1	[95]
	Hübner's description of Aletia argillacea, with translation, [95].	
	Note 2	[96]
	Reasons for rejecting Hübner's description of Aletis argillacea, [95]	
	—Wherein this description and the figures differ from A. xylina, [96]—Doubts of authors concerning A. argillacea, [95]—Search	
	for the type of A. argillacea, [96]—Condition of the Sommer collec-	
	tion, [96]—How specimens of A. xylina are labeled therein, [96].	
	Note 3.	[96]
	No published, full, and complete description of the earlier states of	,
	A. xylina extant, [96] -Reprint of descriptive portion of Dr. C.	
	W. Capers' article "On the cotton caterpillar," [97]—History	
	of ravages of A. xylina prior to 1828, [97]—Sudden abandonment	
	of cotton-plants, [97]—His description of moth, [98]—Egg, [98]	
	-Larva, [98]-Smell of larva, [98]-Habits, [98]-Food-plant,	
	[98]—Pupation, [98]—Description of pupa, [98]—Remedies,	
	[99]—Detailed description of egg, [99]—Six stages of larva,	
	[99]—Pupa, [100].	C1003
	Note 4	[Im]
	Anomis terms, [100].	-
	Note 5	Г10 0 Т
•	Resemblance of larva of Plusia dyaus to that of Aletia xylina, [100]—	,
	Proportion of light and dark specimens early and late in the sea-	
	son, [100].	
	Note 6	[100]
•	Cotton leaves blotched by young larva of Spilosoma acrea, [100].	
	Note 7	[100]
	Larva of Aletia xylina fed on Ipomesa commutata, [100]—Abutilon	
	and Phytolacca defoliated by different larvæ, [100]—Said to feed	
	on "salve bush," [100].	
	Note 8	[100]

2343.	RILEY, C. V.—Continued.
	Notes-Continued.
	Note 9
	Notice of Wm. Trelease's "Nectar: what it is and some of its uses,"
	[101]—Teleology and dysteleology of nectar glands of cotton-
	plant, [101].
	Note 10 [101]
	Rapidity with which the broods of A. xylina follow one another in
`	midsummer, [101]—Time of first appearance of worms, [101]—
	Number of broods, [101]—Prolificacy of moth, [101]—Importance
•	of natural checks upon its increase, [101].
	Note 11
	Influence of winter temperature on time of first appearance of
	worms, [101].
	Note 12
	Theories of hibernation of A. xylina, [101]—Proof of hibernation
	of moth, [102]—importance of this proof, [102].
	Note 13
	Influence of latitude upon time of hatching of insects, [102].
	Note 14
	Number of broods of A. xylina previously recognized, [102].
	Note 15
	some besides cotton, [102]—Failure to find any other, [102]—
	Feeding of larva of Anomis cross on Urena lobata, [102]—Value
	of fiber of <i>U. lobata</i> , [102]—Geographical distribution of <i>U. lobata</i> ,
	[103]—Eggs and larva of Anomis erosa distinguished from those
	of Aletia xylina, [103]—Examination of malvaceous plants in
	herbarium of U. S. Department of Agriculture, [103]—Disad-
	vantages of such an examination, [103]—Plants on which eggs
	were found in the herbarium, [103]-Petition for aid in obtain-
	ing evidence of the food-plant of A. xylina in the more north-
	ern States, [103]-List of malvaceous plants growing in these
	States, with localities, [103].
	Note 16 [104]
	Description of larva of Aspila virescens, [104]
	Note 17
	Travels of E. A. Schwarz in 1878 and 1879, [104]—Reference to
	published accounts of the results of his investigations and those
	of others on hibernation of A. xylina, [104].
	Note 18[104]
	Platyhypena scabra, [104]—Characters of larva, [104]—Food-plants
	of larva, [104]—Pupation and hibernation, [104]—Characters of
	pupa, [104].
	Note 19 [104]
	Seasons of larva of Phoberia atomaris, [104].
	Note 20
	Criticism of paper by A. R. Grote on hibernation of A. xylina,
	[104]—Grote's arguments against hibernation, based on experi-
	ence gained in the same regions which furnish arguments against
	the theory of annual immigration, [105]—Possibility of hiber-
ī	nation admitted by Grote, [105].
	Note 21

2343. RILEY, C. V.—Continued. Notes-Continued. Note 22 References to discussion of J. P. Stelle's claim to have first recommended publicly the use of Paris green for A. xylina, [105]. Note 23 [105] References to descriptions and figures of brush-sacs of moths, [105]. .Worms worse in wet weather than in dry, because more protected from enemies, [105]-Localities of first appearance of worms are those of least molestation, [105]. Appetite of swine for cotton-worms, [106]—Worms eaten by dogs and cats, [106]. Reference to list of birds of southern States, [106]. Note 27 [106] Range of English sparrow in United States, [106]-In hotter portions of the country it is confined to towns and villages, [106]. Note 28 [106] Report by Dr. Geo. Marx on spiders found on cotton, [106]-Preliminary list of spiders which destroy insects noxious to agriculture, [106]—List of spiders observed to devour larvæ of Aletia xylina, [106]—How they capture their victims, [106]—Food-habits of Theridula spherula, [107]—Observations on habits of Oxyopes viridans, by H. G. Hubbard, [107]-Anta captured by larvæ of a Cicindela, [107]. Description of image of Trichogramma pretiosa, [107]. Metamymar n. g. and M. aleurodis n. sp., provisionally named, Note 30...... [108] Description of image and larva of Apanteles aletia, [108]. Apanteles aletiæ parasited by Eupelmus sp., [108]. Description of image of Euplectrus comstockii, [108]. Note 33...... [108] Description of image of Elachistus suplectri n. sp., [108]. Note 34 [109] Sarcophaga sarracenia distinct from S. carnaria, [109]-Points of difference between the Sarcophage of America and Europe gen: erally, [109]—Additional specific characters of S. sarraceniæ, [109]. Note 35 [109] Description of image of Tachina aletia, [109]. Note 36 [109] Description of image of Tachina fraterna, [109]. Note 37 [109] Differences between tachinid larva and that of Sarcophaga, [109]-Differences of puparia, [109]-Reference to description of larva of Senometopia atropirora, [110]; of larva and pupa of Tachina villica, [110]-Description of larva and puparium of Sarcophaga sarraceniæ, [110]; of larva of Belvoisia bifasciata, [110]-Difference of larva of B. bifasciata from that of Tachina concinnata,

242	RILEY, C. V.—Continued.	
20 2 0.	· ·	
	Notes—Continued. [110]—Structure of spiracles in normal form of tachinid puparium, [110]—Description of the puparium, [110]; of puparium	
	of Belboisia bifasciata, [111].	
	Note 38 Synonymy of Cryptus conquisitor, [111]—C. pleurivinetus erroneously	[111]
	given as a synonym of C. annulloornie, [111].	F 3 1 4 3
	Note 39	[III]
	Synonymy of Cryptus samiæ, [111]. Note 40	C1117
	Description of larva and pupa of Chalois ovata, [111]—List of Lepidoptera parasited by Ch. ovata, [111]—Variations in size or imago, [111].	[***]
	Note 41	F111 7
	Synenymy of Tetrastichus seurus, [111]—Description of imago, F1111.	
	Note 42	[111]
	Description of image of Hexaplasta signag, [111]—Difficulty of defining the families Chalcidida, Proctotrupida, and Cympida, [112]—Food-habits of these families, [112].	•
	Note 43	[112]
	Phora aletic not a true parasite, [112]—Habite of this species, [112].	
	Note 44	[118]
	The vast majority of the moths attracted to light said to be males, [112]—If this is so, the usefulness of fires and lights as a remedy is almost nothing, [112].	
	Note 45	[112]
	Antidotes for arsenical poisoning, [112].	
	Note 46	[112]
	Non-fertilized blossoms destroyed by morning showers or spray-	
	ing, [112]—When to make wet applications of poison, [112]. Note 47	F1101
	Test of purity of Paris green, [113].	[113]
	Note 48	Γ1131
	Proportions of Paris green mixture, [113].	,
	Note 49	[113]
	Effect of kerosene and kerosene emulsion on cotton-plants, [113].	
	Note 50	[113]
	Pyrethrum willemoti probably a synonym of P. roseum, [113]—Reference to Willemot's paper on P. willemoti, [113].	
	Note 51 Growth of productive pyrethrum industry in California, [113]— Effect of pyrethrum on warm-blooded animals, [113]—Pyrethrum recommended as a disinfectant and germicide, [113].	[113]
	Note 52	[113]
	Cost of production compared with price of pyrethrum, [113].	• •
	Note 53	[113]
	Enemies of Aletia xylina killed by poisoning the worms, [113].	
	Note 54	[113]
	Positiveness with which various plants have been recommended as insecticides, [113].	
	Note 55	[114]

2343. RILEY, O. V.—Continued.			
Notes—Continued.			
Note 56	[114]		
ances for underspraying apparatus, [115]—Management of apparatus, [115]—Rapidity of service, [115]—Quantity of poison required, [115]—Success of stirrer-pump device, [115]—Construction of stirrer-pump, [115]—Construction of nozzle-pipes, [116]—Adjustment of these pipes, [116]—Success of Eddy-chamber nozzles, [116]—A closed system of pipes advisable, [116]—			
High pressure should always be used, [117]—Adjustment of descending pipes, [117]—Flexile joints recommended, [117]—Advantages of flexile pipes, [117]—Lifting of descending pipes in turning, [118]—Devices for lateral shifting of pipes, [118]—Summary of conclusions from experiments, [118].			
Recommendation to use Paris green in 1872, [119]—Extract from report made on this subject by J. P. Stelle, in 1880, [119]—Comment on the same, [120].	[119]		
Note 58 Description of image of Anomis texana n. sp., [120].	120]		
Note 59	[120]		
Note 60	[121]		
Note 61	[121]		
Note 62 Remarks on specimens sent from Bahia, by R. A. Edes, in 1880, [121]—Characters of egg, larva, and pupa of Anomis sp., [121]— Insects found in cotton bolls, [121].	[121]		
Note 63	_		
Note 64	[121]		
2344. RILEY, C. V. Reports of experiments with various insection substances, chiefly upon insects affecting garden crops, number the direction of the entomologist. <bull. 11,="" 1886,="" 26],="" 34="" [february="" agric.,="" dept.="" ent.="" no.="" pp.<="" s.="" th="" u.=""><th>nade</th></bull.>	nade		
CONTENTS.			
Experiments with insecticides	5 9 23 27 ner.,		
Remarks made at a meeting of the A. A. A. S.; books useful for beginned)F8.		

- 2346. [RILEY, C. V.] [Habitat of Mezium americanum.] < Proc. Eat. Soc. Wash., [30 March], 1886, v. 1, p. 14.

 Occurrence of Mezium americanum in old hay.
- 2347. [RILEY, C. V.] [Arctic insects.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, pp. 14-15.

 Remarks on collection of insects made at Point Barrow, Alaska; prevalence in Arctic regions of species common to America and Europe: occurrence of

in Arctic regions of species common to America and Europe; occurrence of Urocerus flavicornis in Alaska.

- 2348. [RILEY, C. V.] [Habits of Isosoma.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 15.

 Phytophagic habits of the genus Isosoma; I. grande observed ovipositing.
- 2349. [RILEY, C. V.] [Cranberry fruit-worm.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 15.

 Names Acrobasis vaccinii as the well-known cranberry fruit-worm.
- 2350. [RILEY, C. V.] [Rhyssa lunator.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 15.

 States that Rhyssa [= Thalessa] lunator is parasitic and not lignivorous.
- 2351. [RILEY, C. V.] [Tiphia and Rhipiphorus.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 15.

Refers to O. Lugger's statements in regard to the habits of Tiphia and Rhipiphorus as manifestly incorrect.

- 2352. [RILEY, C. V.] [Insects attracted to light.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, pp. 15-16.

 Disfigurement of buildings by the insects and spiders attracted to the electric light on the dome of the Capitol.
- 2353. [RILEY, C. V.] [Parasitic Coleoptera.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 16.

 Parasitic habits of Aleochara unthomyia [= nitida].
- 2354. [RILEY, C. V.] [Scenopinus.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 16.

 Larva of Scenopinus sp. infesting the blanket of a Navajo Indian.
- 2355. RILEY, C. V. Annual address of the president. < Proc. Ent-Soc. Wash., [30 March], 1886, v. 1, pp. 17-27. Brief comments upon the histories or injuries of Agrotis fennica, Hadena de-

Brief comments upon the histories or injuries of Agrotis fennica, Hadena devastatrix, Nomatus erichsoni, Phytonomus punctatus, P. nigrirostris, Pulvinaris innumerabilis, Systachus leucophaus, and Phylloxera vastatrix; notes on the egg parasites of the Acridida; mode of oviposition of some Carabida; advantages of Washington from an entomological standpoint.

- 2356. [RILEY, C. V.] [Gall-making moths.] < Proc. Ent. Soc. Wash.,
 [30 March], 1886, v. 1, p. 30.
 - Walshia amorphella, previously known as a gall-maker on Amorpha frutices, bred from roots of loco weed; Euryptychia saligneana distinct from Padiess scudderiana.
- 2357. [RILEY, C. V.] Sphida, Grote.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 30.

Considers S. obliquata synonymous with Arzama densa.

- 2358. [RILEY, C. V.] [Food of Calopteron and Photinus.] < Proc. Ent. Soc. Wash., [30 March], 1886, v. 1, p. 31.
 - Larva of Calopteron terminale gregarious and lignivorous; Photinus pyralis feeds on snails.
- 2359. RILEY, C. V. A carnivorous butterfly larva. <Science, 30 April, 1886, v. 7, p. 394.
 - Distribution and recorded food-habits of Feniseca tarquinius; Apkidida the normal food of the same.
- 2360. RILEY, C. V. Entomology. Professor Riley to Dr. Shaffer. <Daily Globe [Keokuk, Iowa], 2 May, 1886. S.-b. No. 63, p. 30.</p> Injuries of and means against Abia caprifolium [= Zaraa inflata].
- 2361. BILEY, C. V. A carnivorous butterfly larva.—Plant-feeding habit of Feniseca tarquinius. < Amer. Nat., June, 1886, v. 20, pp. 556-557.
 - Supposed food-plants of the larva of Feniseca tarquinius; first publication of proof that it feeds on plant-lice; Schizoneura tessellata, Pemphigus fraxinifolii, and P. imbricator the species preyed upon.
- 2362. RILEY, C. V. Thrips-Leaf hoppers. < Gardener's Mo. and Hortic., June, 1886, v. 28, p. 174. S.-b. No. 61, p. 56. Letter to Duncan Rhind; remedies against Erythroneura vitifex [= Typhlocyba vitis].
- 2363. RILEY, C. V. Report of the entomologist. <Ann. Rept. [U.S.] Commissioner Agric. for 1885, 1886, pp. 207-343, 1 map, 9 pl. Separate: < Washington: June, 1886, pp. 10+137+10.

TABLE OF CONTENTS.

Introduction	207
Silk-culture, 207—Review of the divisional work in silk-culture, 207—Necessity for a home market for cocoons, 205—Proposed es- tablishment of a filature at Washington, 203—Silk culture in	
France, 209—Destructive locusts, 209—Injurious insects of the year, 209—Correspondence of the Division, 210—Work of the Di-	
vision in economic ornithology, 210—Establishment of an api- cultural station, and the work of the Division in apiculture, 211—	
Work of other divisional agents, 212—Relations of the Division to the National Museum, 213—Publications of the Division, 213—Office force, 213.	
Silk-culture	214
General work of the division, 214—Distribution of eggs, 214—States in which greatest interest is felt, 214.	
Distribution of mulberry trees	214
Number distributed, 214—Danger of indiscriminate distribution, 214—Osage orange, 214—Russian mulberry, 215.	
Distribution of eggs	215
Variety distributed, 215—Designation of races, 215.	
Home-raised vs. imported eggs	215
Feeling against importation, 215—Reasons for importation last year,	
215-Provisions for the next distribution, 216-Eggs rejected,	
216—Labor involved in silk-culture, 216.	
Establishing of filatures—Cost of producing reeled silk	216
23 ENT	

MILEI, O. V.—Conwinger,	
ILK-CULTURE—Continued.	
Importance of filatures, 216—Establishing of stations, 217—Work	
at Philadelphia station, 217-Work and results at New Orleans	
fliature, 217—The Serrell automatic silk reel, 218—Application of	
the saving of this reel to New Orleans figures, 219.	
Work on the Pacific coast	219
Last season's experimental crop, 219—Incorporation of the Ladies'	~1.
Silk-Culture Society of California, 219—The station at Piedmont,	
219—Proposed work at the station, 220—Work of the State board,	
220—Attempt to form a stock company, 220.	
On the precautions necessary in the production of pure silk-worm	
eggs and in properly wintering them	221
Pasteur's classification of diseases, 221.	
Flaccidity (Flacherie)	221
Symptoms and consequences, 221—Action of worms, 221—Color,	
221—Odon, 221—Nature and treatment, 222—Microscopic diagno-	
sis, 222—The bacillus of flaceidity, 222—Examination of the	
chrysalis, 222.	
Pébrine	22
Symptoms, 223—The black spots, 223—Microscopic diagnosis, 223—	
Appearance of chrysalis, 224—Tests for the determination of	
pébrine, 224—Difficulty of using the worm for test, 224—Advantage	
of using the moth, 224—Isolation and examination of the moths,	
224—Maillot's method, 224—Examination of chrysalides, 224—	
Selection of cocoons, 225—Moth cells, preparation, 225—Fasten-	
ings for cells, 225—Preparation of moth for examination, 226—	
The corpuscle of pébrine, 226.	
	227
Wintering the eggs.	221
Colors of eggs, 227—The micropyle, 227—Proper temperature and	
humidity, 227—Hibernating boxes, 228—Acknowledgments, 228.	~~
MISCELLANEOUS INSECTS	228
Destructive locusts or "grasshoppers," Acridida	228
Unusual damage, 228—The Rocky Mountain locust, 223—Proba-	
bilities for 1886, 229—The California migratory locust, 229—	
Abundance the present year, 229—The destructive species in	
previous years, 230—Geographical range, 230—Characters of the	
species, 231—Remedies, 232—New remedy, 232—Non-migratory	
species, 232—The red-legged locust, 232—The differential locust,	
233—The two-striped locust, 233—The lesser locust, 233.	
The periodical Cicada, Tibicon septendecim	233
Preparation of a Bulletin, 233—A seventeen-year and a thirteen-	
year race, 234—Our announcement in 1868, 234—Dr. Smith's man-	
useript, 234-Dr. Phares's earlier discovery, 234-Two distinct	
forms or varieties; specific value of the different forms, 234-	
First discovery, 234—Points of difference, 235—How the species	
should be catalogued, 235—The long period of underground de-	
velopment, 235—The food of the larva, 236—Method of burrow-	
ing of the larva, 237—The transformations, 237—Rising of the	
pupe, 237—Five phases of ecdysis, 238—Time required for trans-	
formation, 238—The Cicada vs. civilization, 238—Influence of	
domestic animals, 238; of building of towns, 238; of the En-	
glish sparrow, 238—Song notes of the Cicada, 239—Notes of other	
insects, 239—First or phar-r-r-raoh note, 239—Second note or	
"screech," 240—Third or "chirping" note, 240—Other notes,	

258

265

270

276

283

2363. RILEY, C. V.—Continued.

MISCELLANEOUS INSECTS-Continued.

240-Variation in time of appearance, 240-Enemies of the Cicada, 241—Supposed sting of the Cicada, 241—Oviposition, 242—Injury caused to fruit trees; remedies and preventive measures, 242-Small injury in larva state, 243-Destruction of the insects as they issue from the ground, 243; by Pyrethrum powder, 243; by Pyrethrum water, 243; by kerosene emulsion, 244— Experiments with substances that kill by thwarting exuviation, 244—Carbolic acid, 244—Acetic acid, 245—Alcohol, 245—To prevent ovipositing, 245—Kerosene emulsion, 245—Consideration of the two 1885 broods, 246-Circular letter, 246-Brood VII, 247-Brood XXII, 248—Points of contact of the two broods, 249— Geographical distribution of Brood VII, 250-Distribution of Brood XXII, 251-Summary of distribution and future appearance of different broods, 252-Influence of climate upon the races, 254—Experiment in the transfer of eggs, 255—Persons assisting, 255-Description of the placing of the different lots of eggs, 256-The Cicada in 1886, 257.

The leather beetle, or toothed Dermestes, Dermestes vulpinus.......
Injury to boots and shoes, 258—History of its occurrence at St.
Louis, 258—Unrecorded points in its habits and natural history,
259—Food of larvæ, 259—The eggs, 259—Growth of larvæ, 260—
The pupa, 261—The beetle, 261—Litigation growing out of its
injuries, 261—History of the Savannah case, 262—Remedies,
263—Care and cleanliness, 263—Use of poisons, 263—Descriptive,
264—Mature larva, 264—Differences between young and old larvæ, 264.

The dark-sided cut-worm, Agrotis messoria

Injury to onions; new habit, 270—Habits and natural history,
270—Number of broods, 270—Hibernation, 271—Remedies, 272—
Treatment of land in spring, 272—Poisoned balls, 272—Kerosene emulsions, 272—Report of John B. Smith, 273—Report of Thomas Bennett, 274.

The pear midge or pear Diplosis, Diplosis pyrivora.....

356	BIBLIOGRAPHY OF ECONOMIC ENTOMOLOGY.	
2363.	RILEY, C. V.—Continued.	
D.	AISCELLANEOUS INSECTS—Continued.	
	Decisive steps for eradication recommended, 283-Life-history and	
	habits, 283—Laying of eggs and growth of larvæ, 283—Habits	
	of larvæ, 284-The cocoon, 284-One annual generation; dates	
•	of issuance of adults, 284—Parasites, 285—Remedies, 285—Is it	
	an introduced species?, 285—Reasons for so considering it, 285—	
	Schmidberger's account of the European pear midge, 286-Dr.	
	Joseph Mik's opinion, 287—Descriptive, 287—Imago, 287—Larva,	
	238—Pupa, 238—The classificatory value of the genitalia, 235.	
	REPORTS OF AGENTS	26
	Report on the locusts of the San Joaquin Valley, Cal. By D. W.	
	Coquillett	28
	Letter of transmittal, 289—Letter of instructions, 289—Location,	
	290—Work of the locusts in general, 290—Species most destructive,	
	291—The devastating locust, 291—The ash-colored locust, 296—	
	292—Where did these locusts hatch?, 292—Consideration of sur-	
	rounding topography, 293—How the young locuste may be dis-	
•	tinguished, 293—Influence of submersion on hatching, 294—Cause	
	of the abundance in this valley in 1885, 294—Early hatching, 294—	
	Lack of rain, 294—The differential locust, 295—Abundance, 295—	
	Egg-pod, 295—Oviposition, 296—Flight, 296—The yellow locust,	
	Abundance, 296—Flight, 296—Cannibalism, 296—Oviposition,	
	297—Other species of locusts, 297—Table showing relative abun-	
	dance, 297—Injury committed by the locusts, 297—To grape-vines,	
	298—To grain-fields and vegetable-gardens, 298—Natural enemics,	
	996—Birds and poultry, 298—Wasps, 298—Mites, 299—Flies, 299—	
	Remedies, 299—Collecting in windrows of dry stems and then	
	burning, 299—Rolling, 300—Kerosene emulsion, 300—Burning sulphur, 300—Horse troughs, 300—Bran, arsenic, and sugar trap-	
	poisoning, 300—Buhach, 302.	•
	Report on the abundance of the Rocky Mountain locust in 1825. By	
	L. Bruner	303
	Letter of submittal, 303—The great abundance of native species,	
	303—Direction of flight of the Rocky Mountain species, 304—	
	Probabilities for 1986, 305—Effect of weather upon probabilities,	
	305—Scarcity of parasites, 305—Abundance of Melanophus atlanis	
	and Camula pellucida, 306—List of locuste noticed in larger num-	
	bers than usual, 307.	
	Notes on locusts at and about Folsom, Cal. By A. Koebele	30t
	Methods adopted by the Natoma Company, 308-Monner of flight,	
•	309—Time of appearance, 309—Losses, 310—Parasites, 310—Ponl-	
	try, 310.	
	Insects affecting fall wheat. By F. M. Webster	311
	Isosoma tritici Riley and I. grande Riley, 311-Notes of breeding, 312-	
	Selection of plants, 314—Facts concerning oviposition, 314—	

Influence of temperature, 314—No males found, 315—The grain Sphenophorus, 315—Found in rye and wheat, 315—Oviposition, 316—Damage to straw and roots, 316—The pupa, 316—Dates of emergence, 316-The white-grub, 316-Distinguished from Sphenophorus, 317-The tarnished plant bug, 317-Found feeding on kernels of wheat, 317—Euschistus fissilis, 317—Dercocoris [= Calocoris] rapidus, 317—Diedrocephala flaviceps, 318—The eggs, 318— Injurious to oats, 318—The chinch-bug, 318—Abundance in fall wheat, 318—On bottle-grass, 318—Parisitized by Mormis, 318—The wheat midge, 318—Breeding notes, 318—Date of emergence, 319.

319

2363. RILEY, C. V.—Continued.

REPORTS OF AGENTS-Continued.

332—Descriptive, 333.

Third report on the causes of destruction of the evergreen and other forest trees in northern New England. By A. S. Packard General review, 320—Condition of the spruce on the coast of Maine, 321—Exemption from insect attack, 321—Condition of the hackmatack in 1885, 321-Verification of last year's predictions, 322-The white-pine weevil, 322—Eggs, 322—Larva, 323—Fupa and its cell, 393—Distinguished from other species, 323—Life-history in brief, 324—Effects of its work, 324—Prevention, 325—The spruce Epizeuxis, 325-Discovery of food-plant, 325-Compared with other species, 325—Descriptive, 326—The spruce plume-moth, 326— Habits, 326—Descriptive, 326—The pine Pheocyma, 327—The evergreen Cleora, 327-Habits, 327-Descriptive, 327-The fir Paraphia, 328—Descriptive, 328—The apruce Therina, 328—Descriptive, 329—The pine Therina, 329—The pine Amorbia, 329— The V-marked Cacoccia, 329—Habits, 329—Descriptive, 330—The hickory Eccopsis, 330—Habits, 330—Descriptive, 331—The variegated Eccopsis, 331-Descriptive, 331-The white-heart hickory Gelechia, 331—Habits, 332—Descriptive, 332—The willow Teras, 332—Descriptive, 332—The purple willow Gracilaria, 332—Habite,

Report on experiments in apiculture. By N. W. McLain......

Letter of submittal, 333—Economy in the production of wax, 334—
Experiments in wax-feeding, 334—Device for feeding sugar syrup and for other purposes, 334—Description, 334—Wintering bees, 335—Loss from thirst, 335—"Roaring in the hive," 336—Device for furnishing water, 336—Bees vs. fruit, 336—House built for experiment, 336—Details of experiments, 337—Conclusions, 338—Fruits used, 339—Experience of Richard Rees, 339—Artificial fertilization, 339—Desirability of accomplishing it, 339—Recent experiments, 340—Experiments with larval and pupal queens unsuccessful, 340—Experiments with image queens, 341—Conclusions and ground for hope, 342—Bee forage, 342—Improved races, 342—Importance of apiculture, 342—Statistics, 343—Necessity and scope, 343—Desirability of accurate crop reports of apiarian products, 343.

2364. RILEY, C. V. Miscellaneous notes on the work of the Division of Entomology for the season of 1885, prepared by the Entomologist. <Bull. No. 12, Div. Ent. U. S. Dept. Agric. [13 July], 1886, 46 pp. 1 plate.</p>

CONTENTS.

Report on the production and manufacture of Buhach. By D. W. Co-	
quillett	7
Additions to the third report on the causes of the destruction of the evergreen and other forest trees in northern New England. By A. S.	
Packard, jr	17
The periodical Cicada in southeastern Indiana. By A. W. Butler	24
Notes of the year	32

2365. R[ILEY], C. V. Hessian-fly. <Suppl. Encyclopædia Britannica, 9th Ed., 1886, v. 3, p. 324, fig.

Natural history, remedies, and figures of Ceoidomyia destructor.

333

- 2366. R[ILEY], C. V. Locust. < Suppl. Encyclopædia Britannica, 9th Ed., 1886, v. 3, pp. 626-628, fig., map.
 - Ravages, life-history, enemies of, and means against Caloptenus spretus; migrations and their cause; map of the region overrun; figure of adult.
- 2367. RILEY, C. V. Some popular fallacies and some new facts regarding Cicada septendecim L. < Proc. Amer. Assoc. Adv. Sci. for 1885, [August], 1886, v. 34, p. 334.
 - Variety cassinii is not the race tredecim of Thicen septendecim; twigs with eggs do not necessarily break off or die to insure the hatching of the larva.
- 2368. RILEY, C. V. Report of the curator of the department of insects in the U. S. National Museum for 1884. <Ann. Rept. of Regents of Smith. Inst. for 1884, 1885 [1886], pp. 185-188.
 - List of accessions, with notes; recommendations for utilizing the Glover plates; needs of the Department.
- 2369. RILEY, C. V. Notes on Feniseca tarquinius, Fabr. <Ca. Ent., October, 1886, v. 18, pp. 191-193.</p>
 Comments on article by W. H. Edwards; records made at the Department of Agriculture proving the carnivorous habits of the larva of Feniseca tarquinius.
- 2370. RILEY, C. V. Two useful lives. <Sci. Amer., 29 January, 1887, v. 56, p. 64. S.-b. No. 63, p. 122. See: <Wine and Fruit Grower, January, 1887, v. 9, p. 10. S.-b. No. 61, pp. 55-56. Brief obituary notices of Louis Bazille and Jules Lichtenstein.
- 2371. RILEY, C. V. Mr. Hulst's observations on *Pronuba yuccasella*.

 <Entom. Amer., March, 1887, v. 2, pp. 233-236.

 Criticism of the observations of G. D. Hulst; summary of the method of ovi-
- 2372. [RILEY, C. V.] Fruit pest extermination. <San Diego Mirror, 5 April, 1887. S.-b. No. 63, pp. 88-89.

 Means against Coccidæ.

position of Pronuba yuccasella; pollination of Yucca by the same.

- 2373. [RILEY, C. V.] [Californian orange insects.] < Daily Herald [Los Angeles], 9 April, 1887. S.-b. No. 63, p. 88.

 Means against Coccidæ.
- 2374. [RILEY, C. V.] Our bugs. <San Francisco Daily Examiner, 16
 April, 1887. S.-b. No. 63, p. 80; 86-87.
 Food-plants, ravages, and means against the white-scale.
- 2375. [RILEY, C. V.] [Remedies and appliances.] <Press and Horticulturist, 16 April, 1887. S.-b. No. 61, pp. 82-84.

 Replies to inquiries made at the annual convention of the State Board of Horticulture of California; means against the codlin-moth; methods of spray
 - ticulture of California; means against the codlin-moth; methods of spraying; differences between allied scales; their methods of dispersal and means against them.
- 2376. RILEY, C. V. Bumble-bees vs. red-clover. < Rural New-Yorker, 23 April, 1887, v. 46, p. 270. S.-b. No. 61, pp. 36-37. Revival of unsettled questions; red-clover sterile in the absence of bumble-bees in New Zealand; some fertilization by other insects probable; introduction of bumble-bees into New Zealand; their rapid propagation there;</p>

beneficial effects on red-clover.

2377. RILEY, C. V. Young grasshoppers. <San Francisco Examiner, 25 April, 1887. S.-b. No. 63, p. 85.

Interview with reporter; ravages of and means against Melanoplus [= Caloptenus] devastator.

2378. RILEY, C. V. Our shade trees and their insect defoliators. Being a consideration of the four most injurious species which affect the trees of the capital; with means of destroying them. <Bull. No. 10, Div. Ent. U. S. Dept. Agric. [7 May], 1887, 69, pp. 27 figs.

CONTENTS.

LETTER OF SUBMITTAL	5
Introduction	7
FOUR PRINCIPAL LEAF-EATERS	8
The imported elm leaf-beetle, Galeruca xanthomelana	8
An importation from Europe	8
Habits and natural history	8
Remedies	11
More recent experience at the Department	14
Past history of the elms in question, 14—Condition and charac-	
teristics of the grove in 1882 and 1883, 14—Extent of injury in	
1882 and 1883, 14-Preferences of the elm beetles for certain	
varieties, 15-Effects of arsenical poisons on insect and plant,	
15-Preventive effects of the poison, 16-Treatment with Lon-	
don purple, 17—Preparation of the poison, 17—Effects of the	
mixture, 17-Treatment with Paris green, 18-Mechanical	
means of applying the poison, 19.	
The bag-worm, Thyridopteryx ephemeræformie	22
Habits and natural history	22
The eggs, 22—The larva and its bag, 23—Pupation, 25—The	
imago or perfect insect, 25.	
Geographical distribution	26
Food-plants	27
Enemies	27
The white-marked tussock-moth, Orgyia leucostigma	29
Habits and natural history	29
The eggs, 29—Development and characters of the larva, 29—	
Habits of the larva, 30—Pupation, 30—The imago, 30—Hiberna-	
tion, 31—Number of annual generations, 31.	
Food-plants	31
Natural enemies and parasites	31
Geographical distribution	33
The fall web-worm, Hyphantria cunea	33
Natural history	33
Limitation of broods, 33—The eggs, 34—The larva, 35—Pupa and	
cocoon, 36—The moth, 36.	
Injury done in 1886	37
Proportionate injury to different plants and shade trees	40
Peculiar effect of defoliation upon some plants	42
Enemies of the web-worm other than insects	43
Predaceous insect enemies	44
Fungus diseases of the web-worm	46
Experiments to obtain percentage of diseased caterpillars, 47.	

2378	B. RILEY, C. V.—Continued.	
	CONTENTS—Continued.	
	The parasites of the web-worm	48
	Telenomus bifidus Riley, 48—Meteorus hyphantria Riley, 49—Apan-	
	teles hyphantriæ Riley, 50—Limmeria pallipes Prov., 51—Tachina	
	sp., 52.	
	SUMMARY OF THE HABITS OF THE FOUR SPECIES	53
	REMEDIES AND PREVENTIVE MEASURES	55
	Winter work.	55
	One simple preventive remedy for all	55
	Pruning and burning	59
	Mulching	60
	Influence of tree-boxes	60
	Whitewashing of trunks	61
	Birds: The English sparrow	62
	THE FUTURE OF OUR TREES. PRUNING	63
	TREES WHICH ARE UNINJURED	64 64
	GOOD AND BAD EFFECTS OF OUR TREES	65
	Prospects the coming season. Conclusion	67
	Index	
2379	9. RILEY, C. V. Remarks on the insect defoliators of our sha	de-
	trees. <new 1887,="" pp.<="" t-p.+12="" th="" york,=""><th></th></new>	
•	Report of address made before the New York farmers, 10 March, 1887; rava	ges
	and means against Galeruca xanthomelana, Thyridopteryx ephemerafors	
	Orgyia leucostigma and Hyphantria cunea.	•
2380	0. RILEY, C. V. Variable moulting in <i>Orgyia</i> . <ent. ma<="" mo.="" th=""><th>g.,</th></ent.>	g.,
	May, 1887, v. 23, p. 274.	
	The female of Orgyia leucostigma undergoes four molts, the male underg	066
	three; tendency of individual larvæ of all orders to vary from the nor	nal
•	number in the species; whenever there is a discrepancy in size of the se	X68
	the smaller undergoes a less number of molts; number of molts wh	
	not sexual dependent on food supply; molting correlated with rate	of
	growth and nutrition.	
2381	I. RILEY, C. V. Pedigree moth-breeding. <ent. mag.,="" mo.="" mo.<="" th=""><th>AV.</th></ent.>	AV.
	1887, v. 23, pp. 277–278.	-J 1
		41.
	Recommends Sericaria mori as the most favorable insect to experiment wi its tendency to vary under new conditions.	GD;
2382	2. RILEY, C. V. Reports of observations and experiments in	ho
	practical work of the division, made under the direction of t	
	- '	
	entomologist. <bull. 13,="" agr<="" dept.="" div.="" ent.="" no.="" s.="" th="" u.=""><th>nc.</th></bull.>	nc.
	[3 June], 1887, 78 pp., 4 figs.	
	CONTENTS.	
	Introduction	~
	Introduction	7
	Report on locusts in Texas in the spring of 1886. By L. Bruner	9
	Fourth report on insects injuring forest and shade trees. By A. S. Pack-	~
	ard, jr	20
	Report on Nebraska insects. By L. Bruner	33
	Tests with insecticides on garden insects. By W. B. Alwood	38 48
	- The state of the	-50

2382. RII	LEY, C. V.—Continued.
Ar s	rents—Continued. record of some experiments relating to the effect of the puncture of some hemipterous insects upon shrubs, fruits, and grains, 1886. By F. M. Webster
Ap	icultural experiments. By N. W. McLain
] I Fo	LEY, C. V. A new apple pest. <sci. 111;="" 112.="" 138;="" 148.="" 18="" 185.="" 1887,="" 216.="" 23="" 29,="" 384.="" 40,="" 50.="" 56,="" 61,="" <colman's="" <gardener's="" adult="" amer.,="" and="" arsenical="" by="" description="" habits,="" haltica="" hortic.,="" july,="" june,="" larva="" mo.="" no.="" od-plants,="" of="" p.="" poisons.<="" punctipennic="" rural="" sb.="" species="" subdued="" td="" the="" v.="" world,=""></sci.>
2384. RII]] Ra	LEY, C. V. A destructive cricket in Louisiana. <florida Dispatch, 20 June, 1887, v. 7, p. 576. Sb. No. 61, pp. 64-65. Beprint: <insect 1,="" 1888,="" 2="" 87-88.<br="" life,="" october,="" pp.="" v.="">vages and means against Gryllus sp., destructive to cotton, peas, tobacco, weet and Irish potatoes.</insect></florida
1 Let	LEY, C. V. Strawberry borers. < Pacific Rural Press, 25 June. 1887, v. 33, p. 559. Sb. No. 61, p. 90; No. 63, p. 110. 10 tter to I. A. Wilcox; recommends bisulphide of carbon as a means against Exercia impropria.
2 6 Fre	LEY, C. V. Life-history of the Icerya. < Pacific Rural Press, 25 June, 1887, v. 33-34, p. 565; 2 July, p. 9. Sb. No. 61, pp. 50-64, 9 figs. om advance proof of Report of U. S. Commissioner of Agriculture for 1886. See No. 2394 for synopsis of contents.
	LEY, C.V.] Cut-worms. < Pacific Rural Press, 25 June, 1887, v. 33, p. 578. Sb. No. 63, p. 108. cans against cut-worms.
I	LEY, C. V. Reports of observations and experiments in the practical work of the division made under the direction of the entomologist. <bull. 1="" 14,="" 1887,="" 3="" 4="" 62="" agric="" august],="" dept.="" div.="" ent.="" figs.<="" no.="" pl.,="" pp.,="" s.="" td="" u.=""></bull.>
	CONTENTS.
Re 1 Re	port on insects injurious to garden crops in Florida. By W. H. Ashnead
(Curculio-proof. By D. B. Wier

2389. RILEY, C. V. The Icerya or fluted scale, otherwise known as the cottony cushion-scale. <Bull. No. 15, Div. Ent. U. S. Dept. Agric. [18 August], 1887, 40 pp. <San Diego Union, 16 April, 1887. S.-b. No. 61, pp. 90-97. <Los Angeles, 13 April, 1887.

S. b. No. 61, pp. 97-104. <Los Angeles Tribune, 14 April, 1887. S. b. No. 61, pp. 104-111. See: <Pacific Rural Press, 23 April, 1887, v. 33, pp. 361-362; 364. S. b. No. 61, p. 48; No. 63, p. 76. <Riverside Daily Press, 12, 13 April, 1887. S. b. No. 61, pp. 65-68; No. 63, pp. 98-105. <Press and Horticulturist, 16 April, 1887. S. b. No. 61, pp. 74-82. <Florida Dispatch, 2 May, 1887, v. 7, pp. 385-388. S. b. No. 63, pp. 75; 77. <Pacific Rural Press, 2 July, 1887, v. 34, p. 9.

CONTENTS.

	Letter of submittal	5
	Introduction	7
	The scale-insects of the orange in California, and particularly the Icerya or fluted scale, etc	9
	Notes on Icerya—Its probable origin the Islands of Bourbon and Mauritius.	27
	The use of gases against scale-insects	35
0300	IRITER C. V. 1 [Classification of insects] < Entom Amer	San-

2390. [RILEY, C. V.] [Classification of insects.] < Entom. Amer., September 1887, v. 3, p. 102.

Comments on the address of J. H. Comstock before the A. A. A. S.

- 2391. [RILEY, C. V.] [Pronuba and its connection with the pollination of Yucca.] < Entom. Amer., September, 1887, v. 3, pp. 107-108.

 Record of the results of recent experiments on the pollination of Yucca and the agency of Pronuba in this work.
- 2392. RILEY, C. V. Beschreibung einer den Birnen schädlichen Gallmücke († *Diplosis nigra* Meig.). <Wiener Entomol. Zeit., September, 1887, v. 6, pp. 201–206, 3 figs.
 - Discovery of the species in America; its habits and life-history; detailed description of all stages; possible difference between European and American specimens; suggests *D. pyrivora* for the latter, if distinct.
- 2393. RILEY, C. V. Some important discoveries in the life-history of the hop-plant louse (*Phorodon humuli* Schrank). <Soc. for Prom. Agric. Sci., September, 1887, 1, No. 9, p. 205. Reprint: <Sci. Amer. Suppl., 24 September, 1887, v. 24, p. 9781. S.-b. No. 61, pp. 117-119. <Gardener's Mo. and Hortic., October, 1887, v. pp. 309-311. S.-b. No. 61, p. 124.
 - Résumé of recent discoveries in the life-history of *Phorodon humuli*; proof of its migration from plum to hop; life of the egg on plum in winter; spring migration to hop; number of broods thus far observed; probable course of later broods.
- 2394. RILEY, C. V. Report of the entomologist. <Ann. Rept. [U. S.] Commissioner Agric. for 1886, 1887, pp. 459-592, 11 pl. Separate: <Washington, September, 1887, pp. 459-592+6+9, 11 pl.

TABLE OF CONTENTS.

Fruit interests of the Pacific coast, 459—Cottony cushion-scale
(Icerya purchasi), 459—Kerosene emulsions, 459—Caustic soda

459

TABLE OF CONTENTS-Continued.

and caustic potash, 459—Remedy for the California red-scale (Aspidiotus aurantii), 460—Experiments made by Mr. Coquillett and Mr. Koebele, 460—Emulsifying oil, 460—Southern buffalo gnats, 461—Results of investigations made on the Southern buffalo gnat and the turkey gnat, 461—Common fall web-worm (Hyphantria cunea), 461—Imported elm-leaf beetle, 461—Spraying appliances, 461—Alternation of generation with the jointworms, 462—Injurious insects of the year, 462—Hop Aphis (Phorodon humuli), 462—Serrell automatic reel for silk, 462—Apicultural station at Aurora, Ill., 462—Work of the Division of economic ornithology, 462—International exhibition of machinery and contrivances for applied remedies against fungiand insects at Florence, 464—Publications of the Division during the year, 464—Proposed publications, 464—Work of field agents, 465—Office force, 465.

MISCELLANEOUS INSECT;.....

466 466

The cottony cushion-scale, Icerya purchasi Introductory, 466—Geographical distribution, 466—In Australia, 466-In Cape Colony, 467-In New Zealand, 467-Importation of the species into California, 468-Its spread and present limitation in California, 469-Food-plants, 471-Original food-plant of Icerya purchasi, 471—Its food-plants in South Africa, 471— Its food-plants in New Zealand, 472—Its food-plants in California, 472—Characters and life-history, 474—The egg, 475— The female larva, first stage, 475—Female larva, second stage, 476-Female larva, third stage, 476-The adult female, fourth stage, 477-The egg-sac, 478-The male larva, probable second stage, 478-Male larva, third stage, 479-The male pupa and cocoon, 479-The adult male, 480-Rate of growth of the different stages, 481-Habits, 481-Exudation of the honeydew, 482-Mode of spread and distribution, 483-Natural enemies, 484—Birds, 484—Predaceous insects, 484—Parasites, 487— Remedies and preventive measures, 488-Importation of parasites, 488-Preventive action, 489-Spraying with insecticides, 489—Fumigating, 491—Bandages around the trunk, 491—Conclusion, 491.

492 493

2394. RILEY, C. V.—Continued. Table of Contents—Continued.

BLE OF CONTENTS—Continued.	
The fall web-worm, Hyphantria cunea	518
Natural history, 518—Limitation of broods, 518—The eggs, 519—	•
The larvæ, 519—Pupa and cocoon, 520—The moth, 520—Injury	
done in 1886, 521—Proportionate injury to different plants and	_
shade trees, 522-Peculiar effect of defoliation upon some	
plants, 525-Enemies of the web-worm other than insects,	
525-Predaceous insect enemies, 526-Fungus disease of the	
web-worm, 527-Experiments to obtain percentage of diseased	
caterpillars, 529—True parasites of the web-worm, 530—Sec-	
ondary parasites, 530—The Telenomus egg-parasite, 531—The	
Meteorus parasite of the web-worm, 531—The Microgaster par-	
asite of the web-worm, 533-The Limneria parasite of the web-	
worm, 534-The Tachina parasite of the web-worm, 534-Rem-	
edies, 535—Pruning, and burning, 535—Mulching, 536—Arsen-	
ical poisons, 536—Emulsions of kerosene, 538—Naphtha, 538.	
Joint-worms, Isosoma sp.	539
The common joint-worm, 539—Its occurrence in Virginia in 1885,	
539—Other recent appearances, 540—Its identity with I. nigrum	
Cook, 541—Parasites, 542—The wheat-straw Isosoma, 542—	
Dimorphism of I. tritici and I. grande, 543—Occurs in California,	
544—In Kansas, 545—Parasites, 546.	
SILK CULTURE	546
Appropriations for the current year, 546—Establishment of a fil-	-
ature at Washington, 546—Osage orange vs. mulberry, 546—	
Rendition of osage cocoons, 547—Mr. Serrell's opinion, 547—	
Satinage, 547—The Serrell reel; cost of work up to the present	
time, 548—Expenses tabulated, 548—Chances for improvement,	
549—Distribution of eggs, 549—Reasons for purchasing foreign	
eggs, 549—Improper choking of cocoons, 550—A simple appa-	
ratus for choking, 551—Cocoons produced in the United States	
in 1886, 551—Tabulated by States, 552.	
REPORT OF AGENTS	552
Report on remedies for the cottony cushion-scale. By D. W. Co-	000
quillett	552
Letter of transmittal, 552—General considerations, 553—Caustic	•••
potash, 554—Caustic soda, 555—Hard soap, 555—Soft soap, 555—	
Kerosene emulsions, 556—Tobacco, 556—Sheep dip, 557—To-	
bacco soap, 557—Vinegar, 557—Paris green, 557.	
Report upon supplementary experiments on the cottony cushion-	
scale, followed by a report on experiments on the red-scale.	
By A. Koebele	558
Letter of transmittal, 558—Introductory, 558—Soap solutions,	•••
558—Preparation of soap, 559—Resin compounds, 559—Lye so-	
lution, 560—Bisulphide of carbon, 560—Kerosene emulsion,	
560—Experiments, 560—Experiments on funigation with bi-	
sulphide of carbon, 569—Experiments on radingation with bi-	
aurantii), 569.	
Insects affecting small grains and grasses. By F. M. Webster	573
TOO OF THE CONTRACT STREET OF THE PROPERTY OF	

Insects affecting fall wheat.....

The wheat-straw Isosoma (Isosoma tritici, Riley), 573—The American Meromyza (Meromyza americana, Fitch), 574—The com-

573

Letter of transmittal, 573.

panion wheat fly (Oscinis? sp.), 574.

0204 Print C V Continued	
2394. RILEY, C. V.—Continued. TABLE OF CONTENTS—Continued.	
Treate official hallon	575
The white-grub (Lachnosterna fusca, Frohl.), 575—The barley	, 010
root-louse (Schizoneura sp.), 575—The grain Aphis (Siphonophora avenæ, Fab.), 576.	
Insects frequenting or depredating upon buckwheat	576
Supposed exemption from insects, 576—Mode of observation, 576—Signification of symbols, 576—List of insects, 576.	
Insects affecting timothy, 578.	
The glassy cut-worm (Hadena devastatrix, Brace), 578—Outbreak	
in Indiana in 1886, 578—Doubt as to species concerned, 578—	
Method of work, 579—Traveling, 579—Parasites, 579—Damage,	
580—Condition of fields in October, 580—The grain Spheno-	
phorus (Sphenophorus parvulus, Gyll.), 580.	
Insects affecting white-clover	580
The flavescent clover weevil (Sitones flavescens, Allard), 580-	
Infests alsike, 580-Mode of attack, 580-Distribution, 580-	
Eggs and larvæ, 580—Hibernation, 580—Summary of life-his-	
tory, 581—The clover-stem magget (Oscinis sp.), 582—Descrip-	
tive, 581-Time and place of oviposition, 581-Number of	
broods, 581.	
Report on experiments in apiculture. By N. W. McLain	583
Letter of transmittal, 583—The "quaking disease," 583—Nature	
of the disease, 583—Odor of infested hives, 583—Treatment	
with brine and soda, 583—Another form of the disease, 584—	
The foul-brood disease, 584—It is contagious, 584—Live pollen	
the medium, 584—Treatment, 584—Results of treatment, 585—	
How the disease spreads, 587—The control of reproduction,	
587—Difficulties, 587—Experiments in 1885, 588—Device for	
holding the queen, 588—Differences in drones, 589—Proportion	
of impotent individuals, 589—Mr. Baldwin's experiments at	
Clarksville, Mo., 589—Comments, 590—Fertilization in con-	

2395. RILEY, C. V. The Hessian-fly in England; its origin; its past; its future. <London Times, 17 October, 1887. S.-b. No. 61, pp. 144-147.

Proposed experiments, 591.

Date of the introduction of *Cecidomyia destructor* into England; probability that it has been introduced from continental Europe since the time of Curtis; conditions in Great Britain unfavorable for its increase.

finement, 590—Apparatus, 590—Results of experiments, 590—

- - Life-history, migrations, and mode of hibernation of Phorodon humuli.
- 2397. RILEY, C. V. On the luminous larviform females of the *Phengo-dini*. <Ent. Mo. Mag., [December], 1887, v. 24, pp. 148-149.

 Structural characters of the larval *Phengodini*; food of *Zarhipis*; characters of eggs, young larvæ, and female larvæ of *Zarhipis* and *Phengodes*; female *Phengodini* considered an archetypal hexapodal form; relations between phosphorescence and differentiation of the sexes.

- 2398. RILEY, C. V. [Introduction of the Hessian-fly into England.]

 <Trans. Ent. Soc. London, [December], 1887, pp. 45-48, Proc.

 Discussion of the date of introduction of Cocidomyia destructor into America and England; concludes from a study of the historical evidence, the distribution and parasites of the insect, that it was probably introduced into England about three or four years ago.
- 2399. RILEY, C. V. Poisonous insects. <Reference Handbook of the Medical Sciences, 1887, v. 5, pp. 741-760, figs. 2971-3020.

 An exhaustive illustrative review of the Arachaida, Myriapoda, and Hexapoda which secrete a poison injurious to man; descriptions of their life-histories; the manner in which the injury is inflicted and the remedies for the same.
- 2400. RILEY, C. V. The problem of the hop-plant louse [Phorodon humuli, Schrank] in Europe and America. <Rept. Brit. Assoc. Adv. Sci., 1887, pp. 750-753. Separate: <pp. 1-3. See: <Nature, 13 October, 1887, v. 36, pp. 566-567. <Gardener's Chronicle, 17 September, 1887. S.-b. No. 61, pp. 333-334. Life-habits of Phorodon humuli; effects of extreme heat and of meterological conditions; natural enemies and means against the Phorodon.
- 2401. RILEY, C. V. On *locrya purchasi*, an insect injurious to fruit trees. <Rept. Brit. Assoc. Adv. Sci., 1887, p. 767. Separate: <p. 1. See: <Nature, 20 October, 1887, v. 36, p. 592. </p>
 Summary statement of food-plants, original home, synonomy, and means against *locrya purchasi*.
- 2402. RILEY, C. V. On the luminous larviform females in the *Phengodini*. <Rept. Brit. Assoc. Adv. Sci., 1887, pp. 760-761. Separate: <pp. 1-2. See: <Entom. Amer., September, 1887, v. 3, p. 107. <Proc. Amer. Assoc. Adv. Sci., 1887 [May, 1888], v. 36, p. 262.
 - Résumé of facts relating to the history, characters, and life-habits of the luminous larviform females in the *Phongodini*; bearing of these facts on the theory of evolution.
- 2403. RILEY, C. V. [Scale on Euonymus latifolia?] <Sci. Amer., 14

 January, 1888, v. 58, p. 27. S.-b. No. 61, p. 148.

 Means against Chionaspis euonymi and other Cocoidæ; formulæ of kerosene emulsions.
- 2404. [RILEY, C. V.] [Larval habits of Lixus.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 33.

 Lixus macer bred from stems of Chenopodium hybridum and from Helianthus; larva of L. parcus a gall producer on stems of Amelanchier.
- 2405. [RILEY, C. V.] [Girdling habits of *Pædisca obfuscata*.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 33.

 Larva of *P. obfuscata* spins a web over the orifice at the amputated end.
- 2406. [RILEY, C. V.] [Early stages of Aphorista vittata and Epipocus punctatus.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 37.
 - Comparative characters between corresponding stages of the two species.

- 2407. [RILEY, C. V.] [Food-habits of Feniseca tarquinius.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, p. 37.

 List of Aphidide preyed upon by larve of F. tarquinius.
- 2408. [RILEY, C. V.] Notes on *Phengodes* and *Zarhipis*. < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, pp. 62-63.</p>
 Structural characters of the luminous larve of *Phengodes* and *Zarhipis* and of an unnamed form from Nevada; difference between the perfect female and the larva.
- 2409. [RILEY, C. V.] [Trees injured by spiders.] <Proc. Ent. Soc.. Wash., [13 March], 1888, v. 1, p. 84.
 Growth of trees retarded by spider-webs.
- 2410. [RILEY, C. V.] [Remarks on exhibited specimens.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, pp. 84-86.
 - Notes on the life-habits of Ageriida. Habits and early stages of Melittia gloriosa, Sciapteron robinia, Phemonos 5-caudata, Ageria impropria, A. albicornis, and A. pyri.
 - 2. Color variation in the larva of Agraulis vanilla. Colorational variation between eastern and western larva of A. vanilla.
 - 3. Miscellaneous insects. Food-plants of Eumonia atala and Cloantha derupta; habitat of Dendrotettix querous n. g. et sp.
- 2411. RILEY, C. V. Further notes on *Phengodes* and *Zarhipis*. <Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, pp. 86-87.</p>
 Comparative characters of the larvæ of *Phengodes* and *Zarhipis*; life-habits and metamorphoses of the same; colorational and structural characters of the larva and larviform female of *Zarhipis*; description of the eggs of *Zar-*
- 2412. [RILEY, C. V.] [Remarks on exhibited specimens.] < Proc. Ent. Soc. Wash., [13 March], 1888, v. 1, pp. 87-89.
 - Notes on the eversible glands in larvæ of Orgyia and Parorgyia, with notes on the synonomy of species. Presence of glands, probably scent organs, in the larvæ of Orgyia and Parorgyia; synonomy and food-plants of some species of Parorgyia.
 - 2. Further remarks on *Phongodes*. Comparative characters of larva and larviform females of *Phongodes laticollis*.
 - Interesting Lepidoptera. Characters of Syntomeida sp., and of an undetermined moth.
- 2413. RILEY, C. V. The British pest. Worthlessness of the sparrow as an insect-killer. <National Tribune, 26 April, 1888.

 Result of the examination of the stomach contents of 522 sparrows, of which 92 only, or 17 h per cent., contained insects, a large proportion of these being innoxious or actually beneficial species; review of recorded observations in North America.
- 2414. RILEY, C. V. Elm-tree depredators. <Newark [N. J.] Press and Register, 10 May, 1888.
 - Report of an address before the Newark Board of Trade; life-history and means against Galeruoa xanthomelæna.
- 2415. RILEY, C. V. On the original habitat of *Icerya purchasi*. < Pacific Rural Press, 12 May, 1888, v. 35, p. 425.
 - Australia probably the true home of Icerya purchasi; its distinctness from I. sacchari,

2416. RILEY, C. V. The buffalo-goat problem in the lower Mississippi Valley. Abstract: <Proc. Amer. Assoc. Adv. Sci. for 1887, [May], 1888, v. 36, p. 362.

Result of late investigations on species of Simulium.

2417. RILEY, C. V. Systematic relations of *Platypsyllus*, as determined by the larva. <Sci. Amer. Suppl., 2 June, 1888, v. 25, pp. 10356–10358, 4 figs.

Review of the literature concerning the systematic position of *Platypsyllus* castoris; additional facts confirming G. H. Horn's view of the coleopterous nature of the insect.

2418. RILEY, C. V. Report of the entomologist. <Ann. Rept. [U. S.]

Commissioner Agric. for 1887, 1888, pp. 48-179, 8 pl. Separate:

<Washington: June, 1888, pp. 48-179+6, 8 pl.

TABLE OF CONTENTS.

Damage by the chinch-bug, 48—Codling-moth, 48—Hop Aphis, 48—Work of Division in regard to silk-culture in the past year, 49—Recent experiments on the fluted or cottony cushion-scale of California, 49—Hydrocyanic acid gas and arseniuretted hydrogen, 49—Possible introduction into California of the parasites of Icerya from Australia, 50—Recent experiments on other scale insects, 50—Professor Osborn's report on the more important insects of the season in Iowa, on the chinch-bug, on hop insects in Wisconsin, 50—Mr. Bruner's report, 50—Mr. Webster's report on the southern buffalo-gnat, 50—The year as a whole one of com-

parative immunity, 50—Apiculture, 50.

THE CHINCH-BUG, Blissus leucopterus. By L. O. Howard...... Introductory, 51—Past history, 51—Barliest appearance, 59—First described by Say, 52-Called "Mormon louse" in Illinois in 1840, 52—First recorded appearance in Missouri, Iowa, Indiana, and Wisconsin, 52—Its damage in the years 1864 and 1868, 52—Estimates of loss in 1871, 52-Great injury in 1874 in Missouri and adjoining States, 52-First injurious appearance in New York in 1882, 53-Damage in succeeding years, 53-Geographical distribution, 53-Indigenous east of the Rocky Mountains, north and south, 53-States in which it does the greatest damage, 53-In Cubs, 54-West of the Rocky Mountains, 54-Specimens from California and Mexico, 55-Injury during 1887, 55-Review of localities and damage this season by the statistician of the Department, 55—Table showing losses in States most damaged, 56— Summary of counties in these States, 57-Food-plants, 57-Cultivated grains and cultivated and wild grasses, 57-Rice, 58-Polygonum, 58-Stages of growth-Descriptive, 59-Egg, 59-Larval stages, 59-Pupa, 59-Imago, 59-Original description, 59-Le Baron's description, 59-Fitch's varieties, 60-Riley's variety melanosus, 60-A new variety, 60-Number of broods and hibernation, 60-First accurate statement, 60-Hibernation, 60-Influence of severe cold, 61-Odor reveals hiding places, 62-Habits, 62-Flight, 62-Oviposition, 62-Young larvæ, 63-Growth, 63-Migration, 63—Habits on corn, 64—Preparations for hibernation, 64-Erroneous statement as to oviposition, 65-Exceptional habits, 65-Natural enemies and diseases, 65-Insect enemies, 65-No true internal insect parasite yet known, 65-A possible hair__

48

THE CHINCH-BUG-Continued.

worm parasite, 65-Lady-bird enemies, 65-Weeping lacewinged fly, 66-True bugs which prey upon it, 66-Inefficacy of lady-bugs when plant-lice are present, 66-Vertebrate enemies, 67—Common quail, 67—Other birds, 67—Quail laws, 67—Diseases, 68-Dr. Shimer's account of his observations on the epidemic in 1865, 68-Professor Forbes' investigations, 69-His studies of bacterial diseases of other insects, 70-Professor Riley's comments, 70-Wet weather and chinch-bug, 71-Professor Forbes' experiments, 71-Comments, 71-Wet weather and the disease, 72-Dr. Thomas' theory, 72-Professor Riley's comments, 73-An anonymous prediction, 73-Table of temperature and rain-fall in North Carolina, 74-Official records of precipitation in chinch-bug States for 1885, 1886, and 1887, 74-Remedies and preventives, 75-Earlier recommendations, 75—Preventions, 75—Clean cultivation, 75—Diversified farming, 75-Rotation of crops, 76-Early sowing and manuring, 76-Rolling, 76-Sowing, an unattractive crop with wheat, 76-Direct winter remedies, 77-Burning, 77-Fall plowing and harrowing, 77-Gas lime, 77-Trapping, 77-Trampling, 77-Direct summer remedies before migration, 77-Irrigation, 78—Burning, 78—Prevention of migration—Direct remedies during and after migration, 79-Ditching, 79-Tarred boards or tar alone, 80-Sowing strips of plants distasteful to the bugs around the fields to be protected, 80-Sowing strips of favored food around the fields to be protected, 80-Hot water and soap-suds, 80-Kerosene emulsion, 80-Professor Forbes' experiments, 81-Mr. Hubbard's formula, 81-Professor Atkinson's test, 81-Professor Osborn's experiments, 82-Mr. Warren's letter, 83-Bogus chinch-bugs, 83-False chinch-bug, 83-Insidious flower-bug, 84-Ash-gray leaf-bug, 84-Flea-like negro-bug, 84-Bibliographical list. 84.

THE CODLING-MOTH, Carpocapea pomonella. By L. O. Howard...... Introductory, 88—Remarks on the bibliography, 88—Geographical distribution, 89-Date of introduction into America, 89-The insect popularly described, 89-Larva, 89-Cocoon, 89-Moth, 90-Habits and natural history, 90—Broods, 90—Its round of life, 90— Irregularity of development, 91-Number of larvæ in a single apple, 91-Larvæ of the second generation, 91-Hibernation in the larval state, 92—Apparent exception to this rule, 93—Cocoons found in apple barrels, 92-Habits of the insect in the North of Germany, 92-Food-plants, 92-Stone fruits of the Rosaces less infested, 92-European records of its occurrence in walnuts and oak-galls, 92-A closely allied species known to feed upon walnuts, 94-Natural enemies, 94-Birds, 94-A hair-worm parasite, 94—True hymenopterous parasites, 94—Predaceous insects which feed upon the larvæ and pupæ, 95-Remedies, 95-The destruction of windfalls-feeding and trampling-the use of sheep and hogs, 96-Jarring or picking infested fruit from the trees. 96-Killing the moth, 97—Miss Walton's observations, 97—Mr. De Long's observations, 97—Capture of the moth with baits, 98— Insect-catching flowers, 98-Summary of the question of attracting the moth, 98-Quotations from Professor Riley's Fourth Missouri Report, 99-Attraction of moth impractical, 99-Trapping the worm-bandages, shingle traps, etc., 100-"Trimble hay-

88

THE CODLING-MOTH-Continued.

band system," 100—Professor Riley's rules concerning this system, 100—The Wier shingle trap, 101—Cloth bands, 101—Dr. Chapin's work in California in 18-2, 102—Spraying the trees with arsenical mixtures, 103—Professor Cook's statements, 103—Professor Forbes' conclusions from his experiments in 1885, 104—Mr. Goff's experiments at Geneva, N. Y., during 1885, 106—Supplementary conclusions by Professor Forbes, 106—Mr. Alwood's experiments in Ohio the past season, 109—Results, 110—Mr. Cushman's letter, 111—The consideration of these poisons in California, 111—Time of day for spraying, 112—Notice of Mr. Dixon's paper, 113—Apparatus for applying the arsenical mixtures, 114.

115

123

123

143

147

154

SILK-CULTURE—REPORT OF THE YEAR'S OPERATIONS. By Philip Walker Distribution of silk-worm eggs, 115—European egg-producers, 115—The establishment of Signor Susani, of Milan, 115—Deydier establishment at Aubenas, France, 117—Mulberry trees, 117—The Cattaneo nurseries, 117—Experimental stations, 118—The Royal Sericultural Experimental Station at Padua, 118—Experimental silk filature at Washington, 119—Production and purchase of cocoons, 120—Co-operating organizations, 122.

Coquillett

Letter of transmittal, 123—The gas treatment for scale-insects, 123—Extract from Mr. Craw's paper, 124—Results of Dr. Dimmock's experiments on various insects with pure gases, 125—The tent, 126—Apparatus for operating the tent, 126—The McMullen tent, 126—The Wolfskill fumigator, 127—The Titus fumigator, 128—The Culver fumigator, 129—The gas, 129—The dry cyanide process, 129—The dry gas process, 130—The cyanide and soda process, 131—Remarks, 132—Agitating the air in the tent, 133—

Experiments, 134—Index to experiments, 142.

REPORT ON EXPERIMENTS AGAINST SCALE-INSECTS. By Albert Koebele

Letter of submittal, 143—The value of arsenic as an addition to
the kerosene emulsion, 143—Results of various experiments with
resin compound, 143—Experiments with resin compound, 146—
Experiments on Aphidides with resin compound, 146.


REPORT ON THE SEASON'S OBSERVATIONS IN INDIANA, AND ESPECIALLY UPON CORN INSECTS By F. M. Webster......

Letter of transmittal, 147—The twelve-spotted Diabrotica, 148—The corn plant-louse, 148—Corn bill-bug, 149—Myochrous denticollis, 150—Flea-beetles, 150—The greasy cut-worm, 150—Ants, 150—Drasterius dorsalis (f), 151—Chinch bug, 151—Corticaria pumila, 151—Calathus gregarius (Say) versus the Colorado potato-beetle, 151—A new enemy to the bean and cow-pea, 152—The straw-

berry saw-fly, 152—Wheat wire-worm, 153,

Letter of transmittal, 154—The turf web-worm or sod-worm, 154—Extracts from crop report, 155—Habits and life-history, 156—Description of different stages, 158—Remedies, 159—Natural enemies, 160—Other mention of the species and related forms, 160—The wheat-head army-worm, 160—Extracts from Iowa crop report, 160—Blister beetles, 161—The false chinch-bug, 169—Notes on miscellaneous insects, 162.

Report on the season's observations in Nebraska. By Law-	
rence Bruner	164
Letter of submittal, 164-Introduction, 164-Colorado potato-	
beetle, 164—Chinch-bug, 165—Causes of increase, 165—Locusts,	
167—Cabbage insects, 168—The codling-moth, 168—Forest-tree	
insects, 168—The American Cimbex, 169—Other insects, 170.	
REPORT ON EXPERIMENTS IN APICULTURE. By N. W. McLain	170
Letter of submittal, 170 — Diseases of bees, 171 — Bacillus alvei	
(Cheshire), 174—Quotations from Professor Cheshire, 172—Treat-	
ment, 174—Starved brood, 174—Symptoms, 175—Remedy, 175—	
The control of reproduction, 175.	



U. S. DEPARTMENT OF AGRICULTURE. DIVISION OF ENTOMOLOGY.

BIBLIOGRAPHY

of

THE MORE IMPORTANT CONTRIBUTIONS

TO

AMERICAN ECONOMIC ENTOMOLOGY.

PREPARED, BY AUTHORITY OF THE SECRETARY OF AGRICULTURE,

BY


SAMUEL HENSHAW.

INDEX

 \mathbf{TO}

Parts I, II, and III.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1890.



INDICES TO PARTS I, II, AND III.

SYSTEMATIC INDEX OF THE NEW NAMES PROPOSED BY R. D. WALSH AND BY WALSH AND RILEY.

[The Walsh and Riley names are followed by W. & R. The first number following the name refers to the number of the paper in the list, the second to the page where the species is first described. As is well known, the Walsh collection was destroyed in the Chicago fire in October, 1871. A few of the types sent to Drs. Hagen and Riley are in the collections of the Museum of Comparative Zoology at Cambridge and the National Museum at Washington. Those at Cambridge are indicated by a *, those at Washington by **.]

ARACHNIDA.

ACARDIA. **∡oarida**.

> Acarus pruni-crumena, 373; 43. s.-enigma (Cecidomyia), 46; 608. s.-semen (Cecidomyia), 46; 606.

MYRIAPODA.

CHILOGRAPHA.

Julida.

Cambala multistriatus (Iulus), 198; 84 = annulata, Say.

HEXAPODA.

PSEUDONEUBOPTERA. Procides.

> Paocus amabilis, 24; 362.* bifaccietus, 30; 188.* confluens, 20; 185. conterminus, 39; 185.* geologus, 24; 362.*1

> > lichematus, 30: 188.* madoscens, 30: 186.*

permedidus, 39: 185,* perplexus, 24; 361.*

pellutus, 24; 361.* purus, 24; 361.* rafus, 99; 185.

semistriatus, 24; 361.*

Perlida.

Acroneura rupinsulencia, 24 ; 363.* Chloroperla brunnipennic, 24; 367.* fumipennie (Perla), 24; 366.* nana, 24; 367.*

Perlides-Continued.

Perla decipiens, 24; 364.*

elongata, 24; 366.*1 flavescens, 24; 363.*1

producta, 24; 365.* varians, 24; 364.*

Ephemerida.

Bestis sicca, 24; 371.*1 Bestisca, 24; 378.

Cloe dubia, 24: 880.*1

ferruginea, 24; 379.* fluctuans, 24: 879.*

mendax, 24; 881.*

Ephemera flaveola, 24; 377.* myope, 30; 207.*

Ephemerella, 24; 877.

consimilia, 24; 878.

excrucians, 24; \$77.**

Hoptagenia, 39; 197.

cruentata, 80; 205.* maculipennis, 39; 396. simplex, 20; 204.*

Hexagenia, 80; 197.

Palingenia flavescens, 24; 878.** pulchella, 34; 575.41

terminata, 24; 878.42

vittigera, 24; 378.*

Pentagenia, 30; 196.

quadripunctata, 39; 198. Potamanthus f odonatus, 24; 372.

Siphlurus interlinesta (Betis), 30: 190= femorata, Say.

Odomata

Agrion binetatum, 24; 867.* dentiferum, 39; 236.*

hageni, 24; 386. 39; 234.*1

434.

```
Jassida-Continued.
Odonata-Continued.
   Cordulia i molesta, 39; 254.
                                                        Typhlocyba aurea, 22; 3.
   Gomphus amnicola, 24; 896.*1
                                                                     binotata, 22; 3.
                                                                     pallidula, 22; 3.
              consobrinus, 89: 242.*
                                                    Tingilidæ.
              fluvialis, 24; $94.*1
                                                        Gargaphia amorphæ (Tingis), 45; 469.**
              grasilinellus, 24; 394.*1
                                                                    tilize (Tingis), 45; 408.44
              quadricolor, 39; 246.
                                                    COLBOPTERA.
              vastus, 24; 391.*1
              ventricosus, 39; 249.
                                                     Scolytidæ.
    Herpetogomphus f rupinsulensis, 24; 388.
                                                        Scolytus fagi, 220; 58.
    Hetserina pseudamericana, 39; 223.*
                                                     Calandridæ.
             rupamnensis, 39; 230.*)
                                                        Sphenophorus zee, 337; 117** = sculptilis
             rupinsulensis, 24; 383.
                                                           Uhler.
             scelerata, 39; 267.
                                                     Ouroulionida.
             texana, 39; 227.
                                                         Anthonomus cratægi, 197; 266.
    Lestes inæqualis, 24: 385.
                                                                      syncophanta, 197; 265.
    Macrogomphus f spiniceps, 24; 389.
                                                                      tessellata, 197; 267.
                                                         Apion languinosum, 197; 209 = walshii, Smith.
    Macromia flavipennis, 24; 398.
              illinoienais, 24; 397.
                                                         Coccotorus prunicida (Anthonomus), 33; 373
    Ophiogomphus mainensis, 39; 255.
                                                           = scutellaris, Lec.
ORTHOPTERA.
                                                         Conotrachelus cratægi, 35; 37.
Phaemida.
                                                                        puncticollis, 34: 21.
    Diapheromera velii, 45; 410.*
                                                     Chrysomelidæ.
HEMIPTERA.
                                                         Fidia viticida, 272; 87.**
                                                         Physonota 5-punctata (Cassida) W. & R.
Coccidæ.
    Chionaspis harrisii (Aspidiotus), 2; 308**=
                                                           767; 4**= unipunotata, Say.
                  furfurus. Fitch.
                                                     Trypetidas.
                salicis-nigræ (Aspidiotus), 373;
                  40 = salicis, Linn.
                                                         Trypeta pomonella, 373; 33.**
    Pulvinaria acericola (Lecanium) W. & R., 389;
                                                     Anthomyida.
                 14'* = innumerabilis, Rath.
                                                         Homalomyia leidyi, 382; 138.
               maclure (Lecanium) W. & R., 389;
                                                                       prunivora, 382; 138.
                 14 = innumerabilis, Rath.
                                                                       wilsoni, 382; 138.
Aphidida.
                                                     Tachinida.
    Aphis bella, 27; 299.*
                                                         Nemoræa militaris (Senometopia) 6; 367**=
          carduella, 27; 300.
                                                          leucaniss Kirkp.
          quercifolize, 27; 298.
                                                     Syrphida.
    Byrsocrypta pseudobyrsa, 27; 306.
                                                         Pipisa radicum W. & R., 495; 83**= 1 femo-
                 vagabunda, 27; 306.*1
                                                           ralis Loew.
    Calaphia, 27; 301.
                                                     Midaida.
                                                         Midas fulvipes, 46; 306.
              betulella, 27; 301.*
   Pemphigus formicarius, 27; 808.
                                                     Cecidomyida.
                                                         Cecidomyia albovittata, 40; 621.*
                formicetorum, 27; 308.*
               ulmi-fusus W. & R., 518; 109.*
                                                                     cornuta, 46; 625.
    Phylloxera caryæ-globuli, 27; 309.**
                                                                     crategi-bedeguar, 876; 79.
                carym-semen (Dactylosphæra)
                                                                     orbitalis, 46; 623.
                 373; 23.*
                                                                     q-pilulæ (Cynips), 41; 481.
                                                                    .s.-batatas, 46; 601.**1
    Schizoneura cornicola (Erigeoma f), 27; 304.
                                                                     s.-brassicoides, 46; 577.**1
                 fungicola (Eriosoma?), 27; 304.
                                                                     a.-cornu, 46; 576. 197; 224.
    Chloroneura, 22; 42 = Empossos, Walsh.
                                                                     s.-coryloides, 46; 588.*
    Empos albicans, 22; 4.
                                                                     s.-gnaphaliodes, 46; 583.*
    Empoasca, 22; 3.
                                                                     s.-hordeoides, 46; 500.
               abnormis (Chloroneura), 22; 4.
                                                                     s.-nodulus; 46; 599,*
               consobrina, 22; 4.
                                                                     s.-rhodoides, 46; 596.*
               malefica (Chloroneura), 22; 4=
                                                                     s.-siliqua, 46; 591.*1
                 viridescens, Walsh.
                                                                     s.-strobilisous, 46; 582. 197; 223.
               maligna (Chloroneura), 22; 4 = ob-
                                                                     a.-strobiloides, 46: 580.
                 tusa, Walsh.
                                                                     s.-triticoides, 46; 596.*1
               obtusa, 22; 4.
                                                                     s.-verruca, 46; 606.
               viridescens, 22; 3.
                                                                     v.-coryloides W. & R., 518; 107.**
    Erythroneura australis, 22; 4.
                                                                     v.-pomum W. & R., 518; 106.**
                  octonotata, 22; 4.
                                                         Diplosis annulipes, 46; 629.
                  siczac, 22: 4.
                                                                 atricornia, 46; 628.
```

¹ Also in the National Museum.

² The pages refer to the separate.

Ichneumonidæ.

Accenitus rupinsulensis, 385; 144.

Midaida-Continued.

atrocularis, 46; 626.

```
Bassus bicapillaris, 385; 88.
            decem-maculata, 46; 631.
            helianthi-bulla, 197; 228.
                                                                 ruficrus, 385; 86.
            septem-maculata, 46; 631.
                                                                 semifasciatus, 385; 86.
                                                                 tripicticrus, 385; 85.
LEPIDOPTERA.
                                                         Catocentrus, 385; 98.
Tortrinida.
                                                         Chorinæus cariniger (Polyrhabdus), 385; 98.
    (Frapholitha prunivora (Semasia), 373; 80.**
                                                         Cryptocentrus, 385; 156 = Mesoleius, Holmg.
    Phoxopteria fragarize (Anchylopera), W. & R.,
                                                         Cryptus albicaligatus; 385; 82.
      409: 90.**
                                                                  albisoleatus, 385; 80.
Pyralidæ.
                                                                  atricollaris, 385; 72. **
    Acrobasis nebulo (Phycita), 2; 308 = indigi-
                                                                  cinctipes, 385; 74.
      nella, Zeller.
                                                                  nigricalceatus, 385; 77.
Geometridæ.
                                                                  picticoxus, 385; 82.
    Aplodes venustus (Hipparchiscus), 40; 81**=
                                                                  rhomboidalis, 385; 74
      mimosaria, Guen.
                                                                  rufifrons, 385; 75.
    Hipparchiscus, 40; 300 = Aplodes, Guen.
                                                          Cteniscus albilineatus, 385; 107.
                                                                    ornatus (Exenteron), 385; 105.
Bombycida.
    Halesidota antiphola, 40; 288 = tessellata, S. & A
                                                          Echthrus annulicornis, 385; 159.
                harrisii,45; 430**= tessellata, S. & A.
                                                          Ephialtes gigas 385; 110.
    Limacodes! tetradactylus, 40; 300.
                                                                    pusio, 385; 111.
                                                                    pygmæus, 385; 111.**
    Phobetron hyalinum (Limacodes !), 40; 200.**
                                                         Exetastes illinoiensis (Leptobatus), 385; 148.
    Sphingicampa, 40; 290.
                                                                    anaveolens, 3/5; 146.
                    distigma, 40; 290 = bicolor,
                                                         Exochiscus, 385; 96 = Orthocentrus, Grav.
                      Harris.
                                                          Exochus albiceps, 385; 96.
Ægeriadæ.
                                                                   annulicrus, 385; 95.
    Ægeria hospes (Trochilium), 197; 270.
                                                                   atriceps, 385; 95.
                                                          Glypta alboscutellaris, 385; 127.
HYMENOPTERA.
Tenthredinidæ.
                                                                  diversipes, 385; 125.
    Euura perturbans, 197; 254.
                                                                  ruficornia, 385; 129.
           s.-gemma; 197; 250 = orbitalis, Nort.
                                                                  rufipleuralis, 385; 127.
                                                          Hemiletes fuscatus, $80; 12 = var. of nema-
           s.-nodus, 197; 253.**
           s.-ovum, 197; 251.
                                                                        tivorus, Walsh.
    Nematus fur, 197; 263.
                                                                      nemativorus, 380; 11.
              hospes, 197; 261.
                                                          Joppidium, 385; 67.
               inquilinus, 197; 260.
                                                                      ruficeps, 385; 70.
                                                          Lampronota amphimilæna, 385; 117.
               mendicus, 197; 261.
               s.-desmodioides, 197; 257.**
                                                                       breviventris, 385; 120.
                                                                        imitatrix, 385; 121.
               s.-pisum, 197; 258.**
               a.-pomum, 197; 255.**
                                                                        interpellata, 885; 118.
     Pristiphora grossulariæ, 140; 123.**
                                                                        pictiventria, 885; 119.
                                                          Mesochorus vitreus, 6; 368.**
                 sycophanta, 197; 263.
                                                          Orthocentrus pusilius (Exochiscus), 385; 97.
 Cynipida.
                                                                         stigmaticus, 385; 101.
     Acraspis q.-erinacei (Cynips), 41; 483.
                                                                         trifasciatus, 385; 100.
     Amphibolips prunus (Cynips) W. & R., 518;
                                                          Pararhyssa, 385; 109 = Rhyssa Grav.
       104.**
                                                           Pesomachus minimus, 6; 868.**
     Ancistrophus, 821; 74.
                                                           Pimpla coaleba, 885; 141.
                    l.-pisum, 821; 74.**
                                                                   investigatrix, 385; 142.
     Andricus q.-flocci (Cynips), 41; 482.**
                                                                   pictipes, 385; 135.
               q.-podagras (Cynips), 41; 491.*1
                                                                   vidua, 385; 140.
     Biorhisa q.-forticornis (Cynips), 41; 490.*
                                                           Polyrhabdus, 385; 98 = Chorinaus Holmgr.
     Ceroptres ensiger (Amblynotus), 41; 496*=
                                                           Polysphinota nigriceps, 385; 144.
                   petiolicola, O. S.
                                                                         nigrita, 885; 144.
                 inermis (Amblynotus), 41; 498.
                                                                         pimploides, 385; 144.
      Holcaspis mamma (Cynips), 518; 102.**
                                                           Tryphon atricoxus, 885; 104.
      Synergus albipes (Synophrus), 41; 496* = lans,
                                                       Braconida.
                  Fitch.
                                                           Apanteles militaris, 6; 369.**
                mendax, 41; 498.
                rhoditiformis, 41; 499* = lignicols,
                                                        Chalcidida.
                                                           Antigaster, 384; 368 = Eupelmus, Dalm.
                  0.8.
                                                           Decatoma dubia, 384; 300 = var. of varians,
      Tribalia, 41; 470.
                                                                       Walsh.
                batatorum, 41; 471.
                                      <sup>1</sup>Also in the National Museum.
```

Chalcidida-Continued.

excrucians, 384; 300 = var. of nigriceps, Walsh. hyalipennis, 384; 300.

nigriceps, 384°, 300. nubilistigma, 384°, 300. simplicistigma, 384°, 300.*** varians, 384°, 300.***

Eupelmus mirabilis (Antigaster), 384; 369.** Eurytoma abnormicornis, 384; 299.

a abnormicornis, 384; 299.
auricepe, 384; 299.**
bicolor, 384; 298.
diastrophi, 384; 299.
gigantes, 384; 300.
globulicola, 384; 299 = var of prunicola, Walsh.
prunicola, 384; 298.**
punctiventris, 384; 299.

Chalcidida-Continued.

seminatrix, 884; 299 = var. of suriceps, Walsh.

Glyphe viridascens, 6; \$70.**

Haltichella perpulchra (Hockeria), 11; 258. Hockeria, 11; 258 = Haltichella, Spin.

Semiotellus chalcidephagus, 384; 369.** Smicra albifrons (Chalcis), 6; 369.**

Formicidæ.

Formica aphidicola, 27; 310. Lesius latipes (Formica), 27; 311.

Pompilidos.

Agenia subcorticalis, 375; 162.

Ceropales rufiventris, 375; 163. Sphecidæ.

Ammophila pictipennia, 375; 164.**

Bembecidæ.

Megastizus brevipennis (Stizus), 875; 162.**

SYSTEMATIC INDEX OF THE NEW NAMES PROPOSED BY C. V. RILEY.

[The types, without exception, are preserved in the collection of the National Museum, Washington, D. C.]

ARACHNIDA Psyllides. Blastophysa, 2272; 75. ACARINA. celtidis-gemma, 2272; 74. Trombidiidas. Tetranychus ! americanus (Leptus), 1326 : 17. Calophya nigripennis, 2272; 69. Ceropsylla, 2272; 76. ! irritans (Leptus), 1326; 18. sideroxyli, 2272; 76. Trombidium giganteum, 1632; 143. Pachypsylla, 2208; 157: 2272; 71. locustarum, 1632; 142. celtidis-mamma, 2208; 157. 2272; muscarum, 1632; 144. Hydrachnida. 73. Hydrachna belostomæ, 1632; 146. Rhinopsylla, 2272; 78. Gamasida. schwarzii, 2272; 78. Urepoda americana, 1626; 275. Jassida. Izodida. Diedrocephala flaviceps, 1767; 78. Ixodes bovis 1309a; 118. Lygarida. Oribatida. Nysius destructor, 1829; 113. = angustatus. Hoplophora arctata, 1363; 216. Uhler. Tyroglyphida. COLEOPTERA Tyroglyphus phylloxerse, 1368; 215. Soolytida. Scolytus carys, 938; 69. = 4-spinosus, Say. Acarus aceris-crumena, 1265; 339. Ourculionidas. HEXAPODA. Ampeloglypter vitis (Madarus), 1059; 132. = secostria, Lec. ORTHOPTERA. ∡eridiida. Podapion, 2231; 62. gallicola, 2231; 62. Caloptenus atlanis, 1423; 169. Tyloderma fragarise (Analcia), 1801; 44. Dendrotettix, 2410; 86. Meloida. quercus, 2410; 86. Gryllida. Hornia 1601. (Ecanthus latipennis, 2026; 61. minutipennis, 1601. HEMIPTERA. Bruchida Coccidas. Bruchus fabes, 1801; 55. Kermes galliformis, 1972; 482. DIPTERA. Mytilaspis pomicorticis, 1329; 96. = pomorum, Occinidas. Bouché. Oscinis brassices, 2291; 322. Aphidida. Anthomyida. Lachnus platanicola, 2188; 196. Anthomyia ver. calopteni, 1557. = angue. Pemphigus acerifolii, 1678; 16. tifrons, Meig. fraximifolii, 1678; 17. zeas, 1050 ; 155. populi-monilie, 1678; 13. Saroophagida. populi-ramulorum, 1678; 16. Sarcophaga sarraceniss, 1390; 238. populi-transversus, 1678; 15. Tackinida. Phylloxera caryse-avellana, 1901; 230.

caryso-fallax, 1423; 118.

carys-scissa, 1901; 230.

Schisoneura americana, 1678; 4.

carys-gummosa, 1423; 118. caryse-ren, 1423; 118.

ulmi (Eriocoma), 1059; 124.

Exorista cecropise, 1112; 101.

Tachina aletiæ, 1712; 162.

flavicauda, 1127; 51.

anonyma, 1811; 129.

Masicera archippivora, 1801; 150.

doryphors (Lydella), 1059; 111.

```
Arilida.
```

Proctacanthus missouriensis (Asilus), 1127; 122 = milberti, Macq.

Hmulida.

Simulium meridionale, 2394; 512. pecuarum, 2394; 512. piscicidium, 1174; 228

Myoetophilidæ.

Mycetophila persica, 963; 397.

Cesidomyidas.

Cecidomyia cupressi-ananassa, 1180; 244. Diplosis pyrivora, 2392.

LEPIDOPTERA.

Pterophoridæ.

Platyptilus carduidactylus (Pterophorus), 1059; 180.

Timeide.

Blastobasis iceryseella, 2394; 485. Coleophora malivorella, 1721; 254. Gelechia gallæsolidaginis, 1059; 175. Holocera glandulella, 1810; 18.

Prodoxus, 1830; 155.

ønescens, 2000; 636. einereus, 2000; 636. decipiens, 1830; 155. intermedids, 2000; 635. marginatus, 2000; 635. Pronubs, 1829; 150: 1836; 55.

maculata, 2000 ; 633.

yuccasella, 1329 ; 151 : 1836 ; 56.

Tortricidas.

Conchylis erigeronana, 1968; 316. œnotherana, 1968; 316. Eccopsis ferrugineana (Exartema), 1968; 317.

monetiferana (Exartema), 1968; 317. Grapholitha gallæ-saliciana, 1968; 320.

ninana, 2176; 661. olivaceana, 1968; 320.

Mellisopus, 1969; 322.

aurichalceana 1969; 323.

Pædisca celtisana, 1968; 319. giganteana, 1968; 318.

Penthina fullerea.* = hebesana, Walk.

Phoxopteris cornifoliana, 1968; 324. murtfeldtiana, 1968; 323.

Proteoteras, 1968; 321.

sesculana, 1968; 321.

Semasia helianthana, 1968; 319.

Steganoptycha claypoleana (Sericoria), 2112;

Teras cinderella (Tortrix), 1311; 47.

Pyralidæ.

Acrobasis nebulella (Phycita), 1311; 42. =var. of indiginella Zeller.

vaccinii, 2291.

Chilo orysællus, 2119; 188. Clydonopteron, 1929; 287.

tecomæ, 1929; 288.

Pempelia hammondi, 1311; 46.

Geometridæ.

Paleacrita, 1438; 278. Synchlora rubivora (Aplodes), 1059; 140.

Noctuida.

Acronycta betulæ, 2280; 2.

populi, 1127; 120. Agrotis cochranii, 964; 414.

morrisoniana, 1409; 286.

scandens, 1059; 78.

Anomis texana, 2343; \$50.

Exyra ridingsii (Xanthoptera), 1411; 348. Laphygma autumnalis (Prodenia), 1301; 116.

> fulvosa (Prodenia), 1301; 117. obscura (Prodenia), 1301; 117.

Lithophane cinerea (Xylina), 1301; 135.

Plusia brassice, 1127; 111.

Pyrophila conspersa (Amphipyra), 1301; 74. Bombyoidæ.

Cerura multiscripta, 1411; 241.

Nola sorghiella, 2119; 187.

\pmb{E} geriadæ.

Bembecia rubi (Ægeria), 1263; 113. = marginata, Harris.

HYMENOPTERA.

Cynipidæ.

Cynips q.-glandulus, 1606; 578. q.-mellaria, 1942; 298.

Didactyum, 1749; 52 — Hexaplasta, Forst. Hexaplasta zigzag (Didictyum), 1749; 52.

Ichneumonidæ.

Hemiteles cressonii, 1059; 177.

thyridopterygia, 1059; 150.

Ichneumon obsoletus, 1570; 55. = var. of brevipennis, Cress.

Limneria lophyri, 1570; 32.

Thersilochus conotracheli (Porizon), 1301; 28. Braconidæ.

Apanteles acronyctæ, 1960; 312.

aletiæ, 1960; 306. cacœcia, 1960; 305. cassianus, 1960; 307.

flaviconchæ, 1960; 308. hemileucæ, 1960; 309.

hyphantrise, 2394; 533.

limenitidis (Microgaster), 1361; 158.

megathymi, 1960 ; 304. paleacritæ, 1960 ; 313.

pieridivora, 2097; 679. = var. ef congregatus, Say.

politus, 1960; 307. rufocoxatis, 1960; 310.

ruiccoxana, 1960; 310 scitulus, 1960; 310.

smerinthi, 1960; 311. thecise, 1960; 308.

Bracon charus, 1428; 75.

Exothecus prodoxi, 1831; 156.

Meteorus hyphantriæ, 2394; 532. Microgaster gelechiæ, 1059; 178.

Microplitis ceratominæ, 1960; 303.

gortynæ, 1960 ; 304. Perilitus indagator, 1311 ; 43.

Sigalphus rufus, 1301; 27. = var. of cureslionis, Fitch.

Spathius trifasciatus, 1329; 106.

^{*}Described and figured in A. S. Fuller's "Injurious insects." <Tilton's Journ. of Hortic. and Florist's Companion, October, 1868, v. 4, pp. 207-209.

Chalcidida.

Cirrospilus flavicinctus.1

Eurytoma bolteri, 1069; 177. = var. of dias-

trophi, Walsh.

Isosoma grande, 2288; 111. 2291; 358.

tritici, 2060; 247.

Merisus subapterus, 2332; 416.

Pentarthron 2 = Trichogramma.

Pleurotropis phyllotretæ, 2291; 308.

Spilochalcis mariæ (Chalcis), 1112; 101.

Stictonotus isosomatis, 2119; 186.

Chalcididæ-Continued.

Tetrastichus esurus (Cirrospilus), 1712; 162. productus, 2332; 419.

Trichogramma minutum, 1301; 158. 2026; 68. pretiosa, 1712; 161.


Proctotrupidas.

Caloptenobia, 1643; 306 = Scelio, Latr. Scelio ovivora (Caloptenobia), 1643; 306 = famelicus, Say.

Telenomus bifidus, 2378; 48. 2394; 531.

¹ Described and figured in Lintner's First Report N. Y. State Entomologist, p. 159.

Proposed (Record of Amer. Ent., 1871, p. 8) for Trichogramma minutum. Pentarthron has, however, been used for a genus of beeties.



INDEX.

About Spains, 113/.	Acrobasia comsociella, 1311.
larva, 1248, 1277.	hammondi == Pempelia hammondi.
Parasites on its larva, 1277.	indiginella, 2, 841, 378, 511, 574, 810, 949,
Abia caprifolia, 2360.	975, 998, 1012, 1069, 1151, 1282, 1311,
Abnormal coccon, 1797.	1590, 1962.
Abroad. Colorado potato-beetle, 1864.	juglandis, 1311.
Abutilon. Aletia feeding on, 2343.	nebulella z. v., 1811 = A. indiginella.
Acacia filiaria. Galle on, 2176.	pebulo = A. indiginella.
ACACIA. INSECTS APPECTING.	vaccinii n. sp., 2391, 2849.
	• • •
Dorthesia characias, 1730.	Acroneura rufinsulensis n. sp., 24.
Grapholitha ninana, 2176.	Acronycta acericola = A. americana.
Lytta cinerea — Macrobasis unicolor.	americana, 571, 841.
Macrobasis unicolor, 1290.	betulæ n. sp., 2280.
Acacias. Large white scale on, 1780.	lepusculina, 1127.
Academy of Science. Address of president St.	oblinita, 1208, 1276, 1801.
Louis, 1564, 1629.	populi <i>n. sp.</i> ., 1127.
Acenitus rupinsulensis n. sp., 395.	xylinoides, 1329.
Acanthia lectularia, 497, 1502, 1741.	Acrosoma spines, 813.
Acaridous gall on Crategus, 376.	stellata, 1869.
Acerina, 412, 1444, 1639, 1710, 1739, 1992.	Acrydii, 1969, 2267.
Acarus aceria-crumena n. sp., 1265.	Actias luna, 776, 1811.
pruni-crumena n. sp., 373.	selene, 1311.
salicis-snigma, 46.	Ad interim committees, 394.
salicis-semen, 46.	Adalia bipunctata, 2452.
scabiei, 497.	Address at Atlanta Cotton Convention, 2038.
Accusation. Unjust, 1587.	on entomology, 1454.
Acer dasycarpum. Scale on, 1575.	
	St. Louis Academy Science. President's,
sp. Supposed tineid larva on leaf of fossil, 2084.	1564, 1629.
	Washington Entomological Society.
Achatodes see, 811.	President's, 2355.
Achemon Sphinx, 1127.	Adelops, 2083.
Acheta abbreviatus — Gryllus abbreviatus.	Adimonia rufosanguinea, 156.
Achetida Gryllida.	Æcidium berberidis, 1605.
Achorutes nivicola, 608.	Ægeria acerni, 743, 1068, 1360, 1368.
Acoloithus falsarius, 134.	albicornis, 2410.
Acorn cups. New oak gall on, 1606.	caudatum = Alcathes caudatum.
gall, 1944.	exitiosa == Sannina exitiosa.
moth, 1310, 1311.	hospes, 197.
Acraspia qerinacei, 41.	impropria, 2265, 2410.
Acridides, 391, 431, 433, 929, 1446, 1452, 1548, 1645,	polistiformis = Sciapteron polistiformis.
1947, 1959, 2002, 2118, 2267, 2811, 2331,	ругі, 2410.
2363.	rubi n. sp., 1963 = Bembecia marginata.
Dimorphism in, 1889.	sp., 802.
Effects of winter on the eggs of, 1550.	syrings = Podosesia syrings.
Egg parasites of, 2356.	tipuiiformis, 55, 840, 481, 974, 1642, 2288.
Acridium americanum, 1418, 1549, 1570.	Ægeriidæ, 302, 576, 2841.
eaten out by ants, 1418.	, life habita, 2410.
• •	Ægerian maple borer. A new, 1360.
obecurum, 2119.	
perigrinum, 1959,	Agiale pofaqui = Megathymus yucces.
	381

Zechna constricta, 759. Alabama. May beetles swarming in, 1812. Agassis, L. Criticism of entomological state. ments, 44. Agave. New use of American, 1498. Agenia architecta, 375. bombycina, 375, 548. cupida, 375. subcorticalis n. sp., 375. Agnomia anilia, 785. Agonoderus comma = A. pallipes. pallipes, 3, 1825, 2817. Agraulis vanille, 2410. Agricultural advancement in the United States, 1750. editorial excursion, 1349. Agriculture and its advancement. Entomology and its relation to, 1829. American writers on insects in relation to, 2288. Entomological work at the Department of, 1892. Insects in relation to, 2238. Legislation in regard to insects injurious to, 1468. New insects injurious to, 2055. Agrilus ruficollis, 821, 1124, 1771. Agrion binotatum n. sp., 24. dentiferarum n. sp., 39. hageni n. ep., 39. Agrotidæ, 8, 868, 901, 964, 1504, 1522, 2310. Agrotia, 2288. annexa, 2291. clandestina, 901, 2391. cochranii n. sp., 964 = A. messoria, Harris. Description of a new species of, 1409. devastatrix = Hadena devastatrix. fennica, 2355. herilis, 964, 1059. inermis = A. saucia. lycarum, 2156. malefida, 2291, 2322. messoria, 901, 964, 1059, 1873, 2156, 2291, 2331, 2368. Harr. vs. Agrotis scandens Riley, 2156. morrisoniana n. sp., 1469. repentis = A. cochranil. saucia, 607, 1059, 1695, 1775, 1826, 1941, 2291. scandens n. sp., 1069, 1408, 2156. Riley vs. Agrotis messoria Harr., 2156. sp., 281. subgothics, 964, 1069. telifera = A. ypsilon. tricosa, 964, 2324. ypsilon, 626, 658, 964, 1059, 1504, 2291. AILANTHUS. INSECTS AFFECTING, 1848. Attacus cynthia, 899, 1179, 1343, 1460, 1718. Œta compta = Œ. punctella. punctella, 1059, 1343. Samia cythia = Attacus cynthia.

Ailanthus ailk-worm, 899, 1311, 1718.

Air-holes not needed in sending insects, 655,

in Missouri, 1460.

naturalized, 1179.

Spread of Pieris rape into, 1720. Alaria florida = Rhodophora florida. Alcatho caudatum, 481. Alcohol. Bugs in, 605. Alcochara anthomyia = A. nitida. nitida, 2253. Aleocharini, 1729. Aletia argillacea = A. xylina. Aletia xylina vs., 2339. Migrations and hibernation of 1000 chryselides. Not, 1826. Chrysalides supposed to be those of, 1775. Moths mistaken for, 1976. xylina, 328, 636, 1127, 1338, 1353, 1363, 1360, 1649, 1657, 1689, 1702, 1712, 1719, 1721, 1722, 1728, 1736, 1749, 1758, 1769, 1814, 1836, 1821, 1845, 1882, 1882, 1888, 1908, 1914, 1982, 1958, 1976, 1997, 2088, 2064, 2069, 2102, 2119, 2130, 2141, 2155, 2164, 2295, 2348. Aletia xylina. (See also Cotton-worm.) Anatomy, 2343. Annual generations, 1736. Bibliography, 2343. Bulletin No. 3, U.S. Ent. Com., 1762. Characters, 1786, 2343. Classification, 2343. Destruction of the moth, 1762. Destructiveness, 1763, 2343. Devices for the destruction of 2343. Flight (powers of), 1763. Habita, 2343. Hibernation, 1763. History of literature of, 2343. Insects liable to be mistaken for, 2348 Invertebrate enemies, 1763. Machinery for the destruction of, Mechanical means of killing, 1763. Meteorological influence upon, 2343 Migrations, 1763. Natural enemies, 1763, 2343. Natural history, 2343. Nomenclature, 2343. in other countries, 2343. Past history in the United States, 2843. Poisoning the worm, 1763. Preventive measures, 1763, 2343. Remedies, 1768, 2343. Stages, 1763. Terrestrial influences, 2343. in the United States. Hibernation of. 2141. Vertebrate enemies of, 1763. vs. Aletia argillacea, 2329. Wet weather (effects of), 1763. Aleurodes, 221. on Oxalis, 1791. Allorhina nitida, 54, 718.

Moths caught in, 1774.

Alpine flowers. Fertilizers of, 1838. Anarsia lineatella, 2235. Alwood, W. B. Report on Ohio insects, 2382. Anasa tristis, 409, 825, 867, 1858, 1948. Tests with insecticides on garden Anatis 15-punctata, 959, 1274, 1450, 1672. insects, 2382. Anax junius, 1851, 2455. Alypia octomaculata, 343, 1059,1127, 1130, 1208, 1363. Anchylopera fragariæ n. sp., 499 = PhoxopterisAmara, 1643. fragarise. Ambiguous Hippodamia, 2119. Ancistrophus n. g., 821. Amblycorypha oblongifolia, 569, 1060, 1157, 1329, l.-pisum n. sp., 821. Anderson, E. H. Cotton-worm in south Texas in Amblynotus ensiger n. sp., 41 = Ceroptres petioli-1883, 2253. cola. Report on cotton insects, 2348. inermis n. sp., 41 = Coroptres iner-Andrena, 800. mis. Andricus q.-californica, 1967. Amelanchier. Lixus parvus on, 2404. q.-flocci, 41. America. Problem of the hop-plant louse in Euq.-operatola, 1352. rope and, 2400. q.-operator, 1352. American Agave. New use of, 1498. q.-podagræ, 41, 45. A. A. S. Entomology at the 1880 meetq.-punctata, 45, 1822. ing of, 1893. seminator, 739, 821, 1037. Permanent subsection of ento-Angoumois grain-moth, 2291. mology, 2027. Angular winged katydid, 1363. bean Bruchus, 1301. Eggs of, 1518. Cimbex, 2291, 2418. Animal kingdom, 393. Entomological Society, 621. Anisced and grain-weevil, 1742. Entomologist, 1309. Anisodactylus baltimorensis, 627. confusus, 2042. Salutatory, 386. insects. Imported and native, 1115, 1127. Anisoplia austriaca, 1935. Anisopteryx, 91, 96, 274, 717, 902, 910, 1891. lackey moth, 1301. Meromyza, 1058, 1059, 2291, 2394. sescularia, 1604. Procris, 1127. Differences between Ani-Staphylinidæ wanted, 1786. sopteryx pometaria tent caterpillar. Eggs of the, 1329. and, 1604. Ames, Iowa, Osborn, H. Experiments at, 2844. pometaria, 1127, 1363, 1423, 1488, 1482, Ammophila pictipennis n. sp., 375, 543. 1539, 1540, 1604, 2149, 2288, 2267. Amorbia. Pine, 2363. Variation in imagos, 1540. Amorpha fruticosa. Walshia amorphella on, 1127, Venation of wings, 1540. 2856 vernata = Paleacrita vernata. Ampeloglypter ater, 821. Anisota = Dryocampa. sesostris, 821, 1059. rubicunda = Dryocampa rubicunda. vitis = A. sesostris. Anomala lucicola, 118. Ampelophaga myron, 837, 1056, 1086, 1127, 1247, Varying, 2291. Anomia erosa, 2848. 1290, 1960. Ampelopsis. Swellings in roots of, 1428. exacta, 1651, 2343. Amphibolipe aciculata = A. q.-spongifica. texana n. sp., 2343. q.-inanis, 45, 518, 1448. urena, 2119. q.-prunus, 518, 1606. xylina = Aletia xylina. Anonymous Tachina-fly, 1423, 1625. q.-spongifica, 41, 45, 518, 1448. Anophthalmus, 2038. Amphicorus, 1747. Ant-hill. Hemispherical larva at bottom of, 1789, bicaudatus, 2, 55, 249, 485, 517,633, 889, 898, 1163, 1185, 1311, 1329, 1747, 1776, Antherma pernyi, 1311. vama-mai, 1311. as a silk-producer, 1346. Amphipyra conspersa n. sp., 1301 = Pyrophila con-Anthicus, 2105. spersa. Anthocoris insidiosus = Triphleps insidiosus. pyramidea = Pyrophila pyramidea. Anthomyia, 1648. pyramidoides = Pyrophila pyramidoiangustifrons, 1541, 1557. des. Hibernation of, 1471. brassics, 2291. Cabbage, 2291. Amphiscepa bivittata, 2116. Amputating insects, 1556. calopteni n. sp., 1541, 1557 = angusti-Anabrus, 2267. frons, Meig. ceparum, 225. purpurascens, 1959. Egg parasite, 1570, 1625, 1643. simplex, 1959. zeas n. sp., 657, 1059, 1065. Analcis fragarise n. sp., 1801 = Tyloderma fraga-

Anametis grisea, 2117.

Anaphora, 2636.

Anthomyids. North American, 1966.

Anthonomus crategi n. sp., 197, 376, 1244.

musculus, 2322, 2331, 2362,

Anthonomus prunicida n. sp., 38 = Coccotorus Aphis, Bad work of the grain, 1688. scutellaris. bella n. sp., 27. quadrigibbus, 42, 254, 1088, 1178, 1176, brassics, 1865, 2291. 1228, 1301, 1358, 2232. carduella n. sp., 27. cerasi == Myzus cerasi. suturalis, 2363. sycophanta n. sp., 197, 821. Cherry, 960. tessellata n. sp., 197. Grain, 2394. Grape-vine, 170. Anthophora abrupta, 372, 1601, 1651. Hop, 1001, 2291, 2394, 2418. sponsa = A. abrupta. Anthreni, 483. maidis, 23, 27. mali, 59, 198, 201, 270, 291, 325, 412, 507, 589, Anthrenus mussorum, 492, 572. scrophularise, 1659, 1663, 1699, 1752. 550, 579, 591, 883, 952, 996, 1056, 1189, 1429, 1899. varius, 5772. prunifoliæ, 678. Antigaster n.g., 384 = Eupelmus. mirabilis n. sp., 884 = Eupelmus miraquercifolis n. sp., 27. ribis = Mysus ribis. vs. Eupelmus, 1932, 1978. Root, 990, 1038. Antiopa butterfly, 1234. rudbeckise = Nectarophora rudbeckisa. Antistrophus n. g., 821, 1131. solani, 1679. lygodesmise-pisum n. sp., 821. vitia, 27, 102, 170. Anta, 1278, 1786, 2418. vs. rust. Grain, 1806. Acridium eaten out by, 1413. Aphodius inquinatus, 1445. and aphides, 520, 2300. Aphorista vittata, 2406. in Arisona. Leaf-stripping, 2532. Aphrophora, 93. gardens. Destroying black, 438. quadrangularis, 601. horticulture. Utilisation of, 2089, 2187. Apiaries. Floating, 1785. injurious in Arisona, 1987. Apiculture. McLain, N. W. Experiments in, 2394. nests in gardens, 496. Report on experi-Red, 2309. ments in, 2363, 2419. Apanteles acronyctes n. sp., 1960. Apidse, 2098. aletim n. sp., 1960, 2343. Apion languinosum n. sp., 197 = A. walshii. cacosciso n. sp., 1960. Apionine. Gall-making genus of, 2231. Apis mellifica, 13, 372, 441, 782, 833, 1455, 1774. cassianus n. sp., 1960. congregatus, 1947. Aplodes, 40. faviconche n. v., 1960. mimosaria, 40. rubivora n. sp., 1959 = Synchlora rubiglomeratus, 2291. hemileucæ n. sp., 2378, 2394. VOPA. limenitidis n. sp., 1301. venustus = A. mimosaria. megathymi n. sp., 1960. Apple bark-lice on pears, 973. militaris n. sp., 6, 12, 1059, 1857, 1442, louse, 1552. Scurfy, 2305. borer and root Aphis, 1038. paleacrite n. sp., 1960. pieridivora n. v., 2097. remedy, 1028. borers, 870, 1583. politus n. ep., 1960. rufocoxalis n. v., 1969. Coleophora, 172L scitulus n. v., 1960, Curculio, 1176, 1301, 1441, 2338. smerinthi n. sp., 1960. go under ground to transform ! Does the, 1228. theclm n. sp., 1960. Apatura alicia, 1363, 1977. gall. Grape-vine, 1329, 1898. celtis, 1356, 1363. growing on a grape-vine, 408, 436. APPLE. INSECTS AFFECTING. clyton, 1856, 1363. here = A. clyton. Acrobasis hammondi == Pempelia hammondi. Amphicerus bicaudatus, 249, 485, 517, 633, 839, lycaon = A. celtis. Apoeliaus mytilaspis, 1329. 893, 1163, 1311, 1329, 1747, 1776, 1820. Aphid. Bitten by an, 2205. Anametis grisea, 2117. Aphides, 952, 1786. Anthonomus cratægi, 1244. Ants and, 520, 2300. quadrigibbus, 42, 1088, 1176, 1228, Aphididæ, 58, 198, 265, 518, 520, 1888, 2017, 2072, 2119, 1301, 2288. Aphis mali, 201, 270, 335, 507, 589, 550, 579, 591, 2230, 2350, 2407. Means against, 80, 102, 198, 508, 644, 678, 883, 996, 1189, 1429. 1148, 1184. Arctia virginica - Spilosoma virginica. of the United States. Genera of the, 27. Aspidiotus conchiformis - Mytilaspis pomo-Notes on the. 1678. harrisii - Chionaspis furfurus, Aphis. Apple borer and root, 1088. rapax, 2285.

Attaons cecropia, 139, 1028, 1076, 1215.

avens = Nectarophora granaria,

APPLE. INSECTS AFFECTING-Continued. APPLE. INSECTS AFFECTING-Continued. Bostrichus bicaudatus = Amphicerus bicaudatus. Brochymena annulata, 684. Bucculatrix pomifoliella, 92, 1811, 1783. Buprestis femorata = Chrysobothris femorata. Capsus oblineatus — Lygus pratensis. Carpocapsa pomonella, 330, 378, 467, 523, 565, 769, 1010, 1022, 1059, 1311, 1320, 1334, 1373, 1398, 1666, 1677, 1682, 1721, 2037. Caterpillars, 1054. Catydid, 29. Cecidomyia sp., 963. Ceresa bubalus, 2195. Chatochilus pomonellus, 874. Chilocorus bivulnerus, 731. Chionaspis furfurus, 69, 345, 373, 579, 951, 1061, 1152, 2305. Chloroneura malefica — Empoasca viridescens. maligna = Empoasca obtusa Chrysobothris femorata, 20, 379, 537, 555, 634, **670, 1059, 1423, 1475, 1513, 1640, 2238, 2267.** Clisiocampa americana, 66, 77, 336, 1301, 2288. Coccus harrisii — Chionaspis furfurus. Coolodasys unicornia, 749. Coleophora malivorella, 1721. Conotrachelus nenuphar, 1214. Corycia vestaliata, 744. Cybocephalus californicus, 2079. Datana ministra, 467, 922, 2222. Diaspis harrisii - Chionaspis furfurus. Dipteron, 548. Elaphidion parallelum = E. villosum. villosum, 839. Emposeca malefica = E. viridescens. maligna = E. obtusa. obtusa, 22. viridescens, 22, 30. Epicærus imbricatus, 296. Eriosoma pyri = Schisoneura lanigera. Euphoria melancholica, 842. Euryomia melancholica = Euphoria melancholica. Galls, 552. Gastropachna americana, 812. velleda = Tolype velleda. Haltica punctipennis, 2383. Helops micans, 963. 2238. Hemipteron, 243. Hyphantria cunea, 112, 2238. textor = H. cunes. Ithycerus noveboracensis, 869, 1033, 1085, 1188. Iulus marginatus == Spirobolus marginatus. Ixodes unipunctata, 1133. Lagos opercularis, 145, 796. Larva, 758, 1054, Lepidopteron, 243, 602. Limacodes pithecium = Phobetron pithecium. sp., 847. Limenitis ursula, 145. Lygus pratensis, 2. Lytta cinerea = Macrobasia unicolor. Macrobasis unicolor, 38. Macrodactylus subspinosus, 748.

Minings, 532.

Mycetophila persion, 963.

25 ent

Mytilaspis pomicorticis = M. pomorum. pomorum, 69, 201, 345, 378, 423, 565, 951, 973, 982, 1059, 1061, 1829, 1552, 1772, 2238, Notodonta concinna = Œdemasia concinna. unicornis = Colodasys unicornis. Oberea sp., 1850. Œcanthus niveus, 1691. Œdemasia concinna, 922. Orgyia leucostigma, 1000, 1151, 2288. Papilio glaucus, 1048. turnus, 1048. Pempelia hammondi, 810. Pemphigus pyri = Schizoneura lanigera. Phobetron pithecium, 1272. Phoxopteris nubeculana, 1721. Phycis indiginella, 511, 574, 810, 949, 975, 998, 1069, 1151, Phycita nebulo = Phycis indiginella. Polycaon confertus, 2103. Prionus laticollis, 2287. Psenocerus supernotatus, 1458. Psocus venosus, 1263, 1526. Reduvius raptatorius - Sinea diadema. Saperda bivittata = S. candida. calcarata, 963. candida, 20, 67, 379, 421, 870, 962, 1023, 1038, 1059, 1218, 1332, 1441, 1513, 1608, 1630, 2238, 2266. Schisoneura lanigera, 373, 467, 495, 963, 997, 1938, 1059, 1187. Sciara mali, 211. Sinea diadema, 638, 781. Solenobia, 38. Spilonota oculana ... Tmetocera ocellana. Spilosoma virginica, 296. Spirobolus marginatus, 963. Teras cinderella, 1311. malivorana, 1311. Tettigonia sp., 951. Tineid, 77. Tmetocera ocellana, 747. Tolype velleda, 972. Tortricid, 847. Tremex columbs, 1126. Trypeta pomonella, 177, 373, 1320, 1553, 2119, Urocerid, 893. Xyleborus obesus, 1583. Apple. Lappet caterpillars on the, 972. Large green caterpillars on the, 1076. leaf Bucculatrix, 1311. crumpler, 574, 949, 975, 998. mistaken for Curculio, 1069. folder. The lesser, 1311. sewer. Chapin's, 1721. skeletonizer, 1311, 1322. tyer. The green, 1311. worm, 1322. Small, 747. maggot, 1553, 2119, 2238. fly, 373, 1320.

and mountain ash. Plant-louse eggs on,

579.

Apple. Oak, 1448. Apple trees. New York weevil on, 1085. Oyster-shell bark-louse of the, 1329, 2238. Scarred, 951. and peach borers, 1475, 1518. Seed ticks under bark of, 1133. pear. Two new foes of the, 22. South. Muscle-shaped bark losse pest. New, 2383. on, 1774. Pigeon Tremex in, 1126. Borer in, 839. twig. borers, 249, 485, 517, 633, 1163, 1311, plant-louse, 883. Plum Curculio breeds in, 1214. 1329, 1747, 1776, 1820. root-blight, 997. Flat-headed, 565. borer, 2287. Small borer in, 1458. plant-louse, 373, 495, 1059. twigs. Catydid eggs on, 29. Eggs on grape canes and, 1323. scions. Swellings on, 548. Larva boring along the axis of, 1850. Small reddish snout-beetle in, 1244. Small galls and minings in, 552. snout-beetle or four-humped Curculio, 1088. vs. apple-tree plant lice. Scab in, 1139. tree bark-lice, 345, 1059. Woolly, slug-like worm on, 796. Native, 1152. worm, 211, 330, 376, 1022, 1059, 1666. louse, 255. in, 177. borer, 1832, 1640. natural history, remedies, 1373. Flat-headed, 1059, 1423, 2238, Notes on, 1682, 1721. 2267 trap. Thomas Wier's, 1312. worms, 523, 1010. horse chestnut. Flatheaded, 1316. by machinery. Killing, 769. Severe cold and hibernating, 769. Insect foes of the, 421. Apples and codling-moth. Michigan, 1677. New facts about the roundheaded, 1630. Melancholy chafer iu, 842. Oviposition of the round-Rose bug on, 748. headed, 2266. Appliances. Remedies and, 2375. Applied entomology. General truths in, 2291, Round-headed, 1059. 1608. Variations in the two-striped Apply soap, 1367. Appropriations for United States Entomological borers, 20, 67, 379, 962, 1059. Commission, 1843. on south side of trees, 634. Aquatic larvæ, 1851. bugs, 685. Sphinx larva, 1951. caterpillar. Unicorn, 749. Arachnida, 1965. caterpillars, 66, 874, 922, 1054. Characters, 59. cocoons, 139. Poisonous, 2399. Egg-masses and cocoons on, 1000. Aragnomus griseus, 2428. Insect foes of the, 467. Aramigus fulleri, 1708, 1721, 1740, 2117. insects, 77, 145, 201, 276, 296, 514, 731, Habits of, 1708. 1056, 1151, 1733. Archippus butterfly, 1301. Lappet caterpillar on, 812. Pupa of, 1235. Large moth on, 1028. Architects. Galls and their, 518, 821. Arotia isabella — Pyrrharctia isabella. worm on, 1048. leaf crumpler, 574, 1311. virginica == Spilosoma virginica. virgo, 674. plant lice, 270, 335, 591, 996, 1429. in Oregon, 1899. Arctic insects, 2347. Scab in apple vs., 1139. Arctiidæ, 2242. louse, 1127. Argiope riparia, 830, 1299. Argynnis, 1301. again. Eggs of, 550. Eggs of the, 507, 539. bellona, 1410. roots caused by root-lice. Knots in, Argyramæba, 1959. 1187. Arhopalus = Cyllene. infaustus = Cyllene decorus. Insects affecting, 963. scales, 69. pictus = Cyllene pictus. Tent caterpillars of the, 336, 1301, robinis = Cyllene robinis. 2238, 2291. speciosus = Plagionotus speciosus. tineid, 92. Aristolochia. Papilio philenor feeding on, 1774. web-caterpillar, 112. Arisona. Ante injurious in, 1987. worms, 511, 744, 810. Arma spinosa = Podisus spinosus. Army-worm, 11, 17, 120, 670, 876, 906, 1127, 1442, 1483, trees. Bark lice on, 423. Eggs on, 638. 1551, 1570, 2119, 2239, 2267, 2269. Injured, 537. Additional notes, 2001. Insects clustered on, 1263. and canker-worm, 910. Native bark-lice on, 1061. Complete life-history of the, 1670.

Army-worm. Eggs of the, 1484. Ash root-borer, 1876. and its enemies, 10. saw-fly, 2291. The fall, 1256, 1267, 1282, 1301, 1482. trees in the nursery. Holes around the roots Further notes on the, 2253. of, 471. Further notes and observations Ashmead, W. H. Report on insects injurious to on the, 1856. garden crops in Florida, 2388. Asby-gray lady-bird, 2119. Hibernation of, 7, 8, 2086. How it comes and goes, 1442. Important point yet to ascertain eggs in, 2118. Asilid larva, 1643. in its history, 1442. Ichneumon fly, 1127. Asilidæ, 1543. and its insect foes, 12. Asilus, 660, 668, 1127. Natural enemies, 1442. flies, 1127, 1543, 1558, 1643, 1959. history complete, 1551. fly. Large, 668, 1269. in New York and other Eastern States. Supposed, 1990. milberti. Northern, 1835. sericeus, 668, 1543. notes and inquiries; its work in clover, 1877. farinalia, 980, 2337. in 1882, 2122. olinalia, 1363. Parasites, 6, 10, 1670. Asparagus beetle, 524. Prospects. Chinch-bug and, 2087. ASPARAGUS. INSECTS AFFECTING. Crioceris asparagi, 524, 2135. question, 8, 9. 12-punctata, 2135. Recent outbreaks, 2262. remedies, 2090. Aspidiotus, 1, 389. Supplementary notes, 1482. aurantii, 2394. True, 647. vs. the clover hay worm, 2000. in the West. Genuine, 2009. Wheat-head, 1570, 2418. rapax, 2235. Canker worms not, 1304. worms. Prof. Riley on, 1800. Three so-called, 328, Aspidoglossa subangulata, 372. What are, 1400. Aspila virescens, 1689, 2343. Arrow-weed. Carpocapsa saltitans on, 2173. ASTER. INSECTS AFFECTING. ARTEMESIA. INSECTS AFFECTING. Diabrotica vittata, 148. Cochylis hilarana, 1059. Epicauta pennsylvanica, 38. Galls, 1347. Gortyna nitela, 75, 940. Articulates, Yersin. Function of the nervous system of, 1959. Aster stalk borer. Dahlia and, 940. Arzama densa, 2357. obliquata, 2223. Asaphes memnonius, 1874. Athysanus maritima, 2116. Asclepias. Larva on, 1535. Asemum mæstum, 2267. Asexual females. Lichtenstein's theory as to di-Atomizers. Sprinklers and, 1857. morphic, 2072. Atlantic migratory locust, 1423, 1625. Ash. Borers in black, 1733. Atta fervens, 1964, 2532. Cocoons on the flowering, 950. Attacus angulifera, 1811. colored locust, 2363. destroyers. Linden and, 1847. gray blister beetle, 1209, 1558. 1112, 1215, 1249, 1311, 1583. leaf-bug, 1127, 1423, 2418, pinion, 1301. ASH. INSECTS AFFECTING. luna - Actias luna. Attacus promethea, 950. Fatus denudatum, 1783. Hyphantria cunes, 1733. 1797, 2204. textor = H. cunes. ricini, 1718. Attagenus injuring feathers, 56. Neoclytus capræa † 1849. Parandra brunnea, 1733, 1876. megatoma, 2113. Trochilium denudatum = Fatua denudatum.

Xyloryotes satyrus, 471.

and mountain ash, 705.

Plant-louse eggs on apple and mountain, 579.

Ash larva. Prickly, 1914.

Asia Minor. Bombylid larvæ destroying locust missouriensis n. sp., 1127 =: ProctacanthusAsopia costalia, 83, 675, 948, 993, 1132, 1363, 2090. conchiformis = Mytilaspis pomorum. barrisii n. sp., 2=Chionaspis furfurus. pinifolii — Chienaspis pinifolii. salicis-nigræ n. sp. = Chionaspis sali-Lytta atrata = Epicauta pennsylvanica. Astoma gryllarium = Trombidium locustarum. locustarum = Trombidium locustarum. Trombidium developed from, 2071. Atlanta cotton convention address, 2038. cecropia, 189, 209, 238, 277, 287, 297, 542, 629. 635, 786, 832, 841, 911, 1028, 1029, 1076, cynthia, 542, 899, 1179, 1311, 1343, 1460, 1718, polyphemus = Teles polyphemus. promethea, 259, 268, 604, 629, 950, 1311, 1486, Attractive but untrue, 1644. Attus, 2302. Aulax, 318. Austrian pines. False caterpillars on Scotch and,

BARLEY. INSECTS AFFECTING. Azalea. Beetles on, 156. Lachnosterna fusca, 2394. Prasocuris varipes on, 156. Bacillus alvei, 2418. Nectarophora granaria, 2394. Back-rolling wonder, 1363. Schizoneura ap., 2394. Bacterial disease of the imported cabbage-worm Siphonophora avenæ = Nectarophora granaria Barley-root louse, 2394. 2251. Bacterium, 2251. . F. M. Webster. Insects affecting, 2394. Barnard, W. S. Machinery for destroying Aletia, Bad packing, 837. Bætis femorata = Silphlurus femoratus. 2343. interlineata n. sp., 39 = Silphiurus femora-Tests of machinery for destroying tus. the cotton-worm, 2253. Basket-worm, 271, 1059, 1090, 1650. sicca n. sp., 24. Bætisca n. g., 24. Bass. Insects from stomach of rock, 1793. obesa. Pupa of, 43. Larvæ in stomach of black, 1792. Bag-worm, 1059, 1090, 1189, 1650, 2378. Bassus bicapillaris n. sp., 385. ruficrus n. sp., 385. New, 1441. at South Pass, Ill., 1156. semifasciatus n. sp., 385. Thyridopteryx ephemeræformis. Retripicticrus n. sp , 385. marks on, 2272. Bass wood. Dipteron on. 548. worms, 182, 266, 538, 641, 738, 1036. Batrachedra salicipomonella, 821. again, 815. Bazille, Louis. Obituary, 2370. and borers, 1472. Be on the guard, 708. Ways of, 1424. BEAR. INSECTS AFFECTING. Bahia, Brazil. Cotton culture and the insects af-Bruchus fabæ, 1001, 1631, 2127, 2238. fecting the plant at, 1788a. granarius, 1120. ebsoletus, 1155, 1232, 1296. Bailey, J. S. Tree borers of the family Cossidæ, 2253. Epilachna corrupta, 2235. Lytta cinerea == Macrobasia unicolor. Balaninus rectus, 1810, 1311. Balsam fir. Lachnus strebi on, 1039. Macrobasis unicolor, 38. Bamboo Sinoxylon, 2454. Bean. A new enemy to, 2418. Banded Ips in calyx of pear, 1239. weevil, 1155, 1296, 2127, 2238. Baridius sesestris = Ampeloglypter sesestris. American, 1301. trinotata = Trichobaris trinotata. Beans. Buggy, 1631. Bark beetles, 295. Beautiful wood nymph, 1127, 1363. borer. Hickory, 938, 1329. Bed bugs, 1502. Peach-tree, 2047. . Lepidium vs., 1741. Bedeguar of the rose, 1939. borers, 220. boring beetles. Packard on the develop-Bee bread devoured by worms, 1293. ment of, 2267. Bug preying on honey, 13. lice, 1, 264, 935, 944, 989, 994, 1007. disease. Possible cause of the, 704. again, 1040. enemy, 1186. on apple-trees, 423, 1059. fly, 1959. Native, 1061. killer. Nebraska, 1127, 1311. Curculio and, 487. killers, 1543. eggs in Missouri. Supposed, 1084. Beetle named, 627. on grape-vine, 1212. Remarkable felting caused by a, 2113. Maple, 1004. - in Russia. Excessive injury by, 1935. Means against, 104. on sugar maple, 1014. Native apple-tree, 1152. in thistle. 169. on the pear, 982. Beetles, 1736. pears. Apple, 973. around peach-trees, 751. Food-habits of ground, 1760. Preventive of, 1012. remedy, 896. in dried English currants, 1260. on rose bushes, 1303. in flouring-mills, 1261. louse, 987, 1397, 1583. injuring cabbages and fuchsias, 1868. in stomach of meadow lark, 1013. Apple, 1552. apple - trees South. in the corn-field, 2817. Muscleshaped, 1774. Mites parasitic on, 315. enemy, 154. named, 732, 802, 806, 1172, 1187. Imported apple-tree, 255. on buttercup and asales, 156. Plant-feeding habits of predaceous, 1956. Insect foes of the, 417. in Missouri. Oyster-shell, 1169. supposed to be feeding on wheat, 1746. on Osage orange, 389. swarming about the lawn, 718. Oyster-shell apple, 1127, 1329, 2238. under dead fish, 817. Scurfy apple, 2305. working in wheat, oats, and rye, 1259. Belated individuals of the periodical Cicada, 618. Tulip-tree, 271.

Belfrage, G. W. Obituary, 2161. Belostoma americanum, 473, 534, 738, 1570, 1584. grande = B. americanum. Belvoisia bifasciata, 1329, 2343, Bembecia marginata, 576, 1363, 1454. rubi = B. marginata. Bembex faeciata, 543. marginata, 2230. Bembidium 4-maculatum, 3. Bennett, T. Report of experiments at Trenton, N. J., 2344. Berberry. Aphidida on, 644. Berne. Phylloxera convention at, 2147. Berry and cherry twigs, 1503. Beschreibung einer den Birnen schädlichen Gallmücke, 2392. Best's fruit-tree invigorator again, 530. invigorator once more, 545. Betula nigra. Acconycta on, 2280. Bibio albipennis, 257, 683. Bibionidæ, 2076, Bibliography of gall literature, 2059. Bidens chrysanthemoides. Syrphus sp. caught by, 1761. Bill bug. Corn, 2291, 2418. Biorhiza forticornia, 41, 745. Birch. Clytus sp. ? on, 2267. Bird question dispassionately considered, 1768. Birds. Professor Riley in favor of the, 1675. Relation of Carabidæ and Coccinellidæ to. vs. insects, 199, 369. Bittacus apterus. Habits of, 2092. Bitten by an aphid, 2205. Bitter-sweet. Enchenopa binotata on, 1913. White waxy secretion on stems of, 1913. Black ants in gardens. Destroying, 438. ash. Borers in, 1733. bass. Larva in stomach of, 1792. blister beetle, 1059, 1558. beetles on potatoes, 1881. breeze fly, 1127. bug, 314. current. Current-worms and, 1079. grape-vine caterpillars, 1018. knot once more, 930. legged tortoise beetle, 1127. potato beetles. Large, 1206. rat blister beetle, 1059, 1558. walnut. Insects feeding on sap of, 1195. and willow. Enemies of the, 2333. Blackberry borers, 302. cane borers, 576. gall insects, 318. Pithy, 1131.

Galls on, 263.

Ægeria rubi = Bembecia marginata.

Bembecia marginata, 576, 1364, 1454.

nebulosus, 263, 610, 1131.

Diastrophus cuscutæformis, 609.

sp., 318.

BLACKBERRY. INSECTS AFFECTING.

sp., 302.

Agrilus ruficollis, 1771.

Eudemis botrana, 1232.

Lobesia botrana = Eudemis botrana. Oberea bimaculata, 302, 783, 1863. perspicillata = 0. bimaculata. tripunctata, 1503. Œcanthus níveus, 1503. Psylla rubi = Trioza tripunctata. Trioza tripunctata, 669. Blackberry. New insect foe of the, 669. and raspberry canes. Gouty galls on, 1771. scale, 117. twig. Fuzzy galls on, 609. ·Pithy galls on, 610. Blastobasis icerywella n. sp., 2394. Blastophysa n. sg., 2272. celtidis-gemma n. sp., 2272. Blatta, 2267. germanica = Ectobia germanica. Blattidæ, 573, 1858. Blennocampa pygmæa, 1056, 1252. Blepharida rhois, 849, 1863. Blepharoceridæ, 1981, 2004. Blight. Apple-root, 907. Blind-eyed Smerinthus, 1912. Going it, 588. as those who shut their eyes. None so, 275. Blissus leucopterus, 6, 90, 180, 398, 548, 886, 888, 890, 891, 894, 893, 1068, 1127, 1357, 1402, 1428, 1432, 1453, 1508, 1538, 1681, 2018, 2089, 2048, 2067, 2119, 2143, 2237, 2252, 2271, 2418. Blister beetle. Ash-gray, 1059, 1209, 1558. Black, 1059, 1556. rat, 1059, 1558. Margined, 1059, 1558. on potatoes. Black, 1881. Retarded development in, 1800. Striped, 1059, 1568. beetles, 134, 289, 301, 362, 1059, 1558, 2238, 2418. Larval characters and habits of, 1600. from New Mexico, 1796. Notes on life-history of, 1651. Pear leaf, 1739. Blcod-red lady-bird, 2119. sucker and pear slug, 1253. sucking cone-nose, 795, 854. Conorhinus, 2291. Blow-fly. Cheese-fly and, 1266. Bluebird. Ichneumonid from stomach of, 1878. Larvæ from stomach of, 1871. Bluebirds feeding on parasitic and predaceous insects, 1885. Blue caterpillars of the vine, 1127, 1368. spangled peach-worm, 1301. Bogus chinch-bug, 1127, 1423. Colorado potato-beetle, 1558. bug, 519. Thrips. True Thrips and, 203. Yucca moth. True and, 1804. Boll-worm, 821, 1801, 1353, 2119, 2164, 2848. Report on, 2164. worms feeding on oabbage, 1915. Bombardier beetle, Kansas, 1311.

BLACKBERRY. INSECTS AFFECTING-Continued.

Lecanium sp., 117.

Bombus, 800, 1032, 1226.

Brimstone for borers, 981.

British pest, 2413.

Broad-necked Prionus. Eggs of, 1329.

Hibernation of the queens of, 1032. winged katydid, 1329, 1363. Brochymens annulata, 684. pennsylvanicus, 800. , virginicus, 1774. Brown mantispian, 1243. Brown's answer to inquiries about Colorado pe-Bombycidse, 1718. Bombylid larvæ destroying locust eggs in Asia tato-beetle. Mayor, 1491. Bruchus of Europe just imported. Grain, 1120. Minor, 2118. fabæ n. sp., 1301, 1631, 2127, 2238. Bombylli, 1959. Bombyliidæ, 2118. Grain, 1301. Larval habits, 1947, 2169, granarius, 1120, 1301. Life-habits, 2002. obsoletus, 1155, 1282, 1296. piei, 434, 1120, 1301, 1357, 1916, 2238. Natural history of certain, 1928. Bombylius major, 1947, 1959. Bruner, L. Additions to the chronology of locust ravages in 1880-'81, 2267. Bombyx arrindria, 899. mori = Sericaria mori. List of North American locusts, 2267. Bordered soldier bug, 1059. Notes on other locusts and the west-Borer, 47. ern cricket, 2267. Borers, 55, 141, 283, 967, 1083. from Nebraska, 2291. Bag-worms and, 1472. Observations in the Northwest on the Rocky Mountain lo-Brimstone for, 931. cust, 2165. Canker-worms and, 941. Tansy for, 1016. on the Rocky Mountain Boston. Entomologists at, 1894. locust and other in-Bostrichus = Amphicerus. sects in the Northbicaudatus = Amphicerus bicaudatus. west during the summer of 1863, 2277. Botis. Cauliflower, 2232. repetitalis, 2232. Report on the abundance of the Bot flies, 114. Rocky Mountain locust fly. Cattle, 2238. in 1885, 2363. locusts in Texas in Horse, 2238. Sheep, 2238. spring of 1886, 2882. Squirrel, 526. Nebraska insects, 2382. Botrytis rileyi, 2232. of the season's observations Bots, 1563. in Nebraska, 2418. Bowels. Larvæ in human, 382. on the Rocky Mountain lo-Box Pavlla, 2291. cust in Montana in 1880, Brachinus, 802. 2267. americanus, 802. the Rocky Mountain lo-Brachyrhynchus, 553. cust in Wyoming and Brachytarsus variegatus, 18, 1461, 1479. Montana, etc., in 1881, Bracon charus n. sp., 1423. 2267. Cherished, 1423. Bucculatrix. Apple-leaf, 1311. scolytivorus n. sp., 1329. pomifoliella, 92, 1311, 1733. Bradynotes opimus n. sp., 1959. thuiella, 1311. Brandt on nervous system of insects, 2225. Buck fly, 595. Branner, J. C. Cotton-caterpillars in Brazil, 2343. moth, 1329. Preliminary report of observations Buckeye leaf. Sericoris claypoleans on, 2114. upon insects injurious to cotton, stem-borer, 2114. orange, and sugar-cane in Brazil, Buckwheat to destroy insects. Use of, 1744. 2277. Webster on insects affecting, 2394. Brauer on Hirmoneura. Review of, 2275. Bud-eating insects, 1408. Braula cœca not particularly injurious to the Buffalo. Entomology in, 2035. honey-bee, 1982. gnat problem in the Lower Mississippi Brazil. Branner, J. C., on cotton-caterpillars in, Valley, 2416. 2348. Southern, 2291, 2394, 2418. insects injurious to gnate, 2394. cotton, orange, and Webster on, 2388. sugar-cane in, 2277. tree hopper, 415, 1329. Lignified snake of, 2136. injurious to potatoes, 2167. Breakfast. Entomological, 2311. Bug. A most precious, 575. Breeding-cages, 695. Buggy beans, 1631. silk-worms, 1648. peas, 1916. Breeze-fly. Black, 1127. Bug-hunter in Egypt, 872. Brenthian. Northern, 1363. hunters. Cruel, 440.

Bugs, 354.

in alcohol, 605.

Butterfly larva injurious to cotton squares, 1872. Bugs. Hogs ve., 387. Our, 2374, pine-trees, 2121. Toads vs., 1108. larvæ. Summer dormancy of, 1410. Buhach. Coquillett on the production and manunamed, 666. facture of, 2364. tongue, 1815. Bulletin Div. Ent. U. S. Dept. Agric., No. 1, 2164. trees of Monterey again, 2052. wings. Chemical change in color of, 1935. 2, 2165, 8, 2253. Byrsocrypta pseudobyrsa n. sp., 27. 4. 2277. vagabunda n. sp., 27. 6, 2304. Cabbage Anthomyia, 2291. 8, 2315. bug. Harlequin, 1099, 1311, 1534, 2238, 2263, ° 9, 172√. 2291. 10, 2378. butterfly. Imported, 1127, 2291. 11, 2344. Larger, 2232. 12, 2864. Southern, 1127, 2232, 2238. 13, 2382. cut-worms, 2291. 14, 2388. Remedies for, 2291. 15, 2389. flea-beetle. Colorado, 2291. Michigan Agric. College, No. 39 (Reinsects, 1297, 2291, 2418. view), 2501. CABBAGE. INSECTS AFFECTING. United States Entomological Commis-Agrotis annexa, 2291. sion, No. 1, 1577. clandestina, 2291. 2, 1578. malefida, 2291. 8, 1736. messoria, 2291. 6, 2026. saucia, 2291. Bumble-bees vs. red-clover, 2376. telifera = A. ypsilon. What becomes of, 1032. ypsilon, 626, 2291. Bundle of entomological queries, 556. Anthomyia brassics, 2291. Buprestidse, 821, 2267. Aphis brassics, 1365, 2291. Buprestis divaricata = Dicerca divaricata. Botis repetitalis, 2232. femorata = Chrysobothris femorata. Caloptenus spretus, 2291. Burr-oak. Insects stripping the, 966. Doryphora 10-lineata, 2291. Elaterid, 740. Burgess, E. Anatomy of Aletia, 2348. Burying beetles, 815, 845. Epicærus imbricatus, 1968. Bushels of butterflies, 238. Hadena devastatrix, 2291. Butalis cerealella, 1232. Heliothis armigera, 1915. Buteo pennsylvanicus, 643. Lygus lineolaris = L. pratensis. Buthus carolinianus, 458, 1119. pratensis, 2291. Butler, A. W. The periodical Cicada in south-Mamestra chenopodi = M. trifolii. eastern Indiana, 2364. picta, 1127, 2232. Buttercup. Beetles on, 156. subjuncta, 2291. . INSECTS AFFECTING. trifolii, 2232. Adimonia rufosanguinea, 156. Murgantia histrionica, 1097, 1297, 1311, 1584, Galeruca rufosanguinea = Adimonia rufosan-2238, 2263, 2291. guines. Nysius angustatus, 2291. Butterflies. Bushels of, 288. Oscinia, 2291. caught by the tongue, 1761. Phyllotreta albionica, 2291. Controlling sex in, 1852. vittata, 2291. Effects of cold applied to the chryssimmermani, 2291. alids of, 1778. Pieris monuste, 2232. oleracea, 835, 1127, 2232, 2238. Hackberry, 1856, 1368. Imitative, 613. protodice, 1127, 2232, 2238. rapes, 1127, 1257, 1279, 1294, 1694, 1721, Migrations, 1770, 1991. 2221, 2232, 2238, 2251, 2298. Migratory, 1622, 1635. named, 461. Pionea rimosalis, 1915, 2232. of New England. Notice of Scud-Plusia brassicse, 1127, 2232, 2238. der's, 2464. Plutella cruciferarum, 1291, 1297, 2232. North America. Review of Ed-Strachia histrionica - Murgantia histrionica. ward's, 396, 479, 623, 827. Cabbage-lice, 1365. Odor in, 2025. Mamestra, 2282. Philosophy of the pupation of, 1711. Oscinia, 2291. Swarm of, 406, 1050. pest. Our new, 1257. peats, 740. Butterfly chrysalis, 1585. Goat-weed, 1117, 1127, 1329. Piones, 2232. larva. Carnivorous, 2359. plant-louse, 2291.

Cabbage plants. Cut-worms severing, 626.

Plusia, 1127, 2232, 2238.

Plutella, 1291, 2232.

Callosamia promethea = Attacus prometh

Callostoma fascipennia, 2075, 2118.

Calocoria chenopodi, 2458.

worm. Bacterial disease of the imported, rapidus, 2363. Calomicrus. Characters of, 1799. Imported, 2282, 2288, 2291. Calophya nigripennia n. sp., 2272. A new remedy for the imported, Caloptenobia n.g., 1648 = Scelio. 220R ovivora n. sp., 1643 = Scelio fameliparasite, 2221. cus. in the South. The imported, 1714. Caloptenus, 2267. Successful introduction of a paraatlanis n. sp., 1423, 1549, 1643, 2198, 2232, site of the imported, 2291. The westward progress of the im-Packard on the development of, ported, 1694, 1721. 2267. Zebra, 2232. hivittatus, 1959. worms, 835, 1137, 1279, 1915, 2282. devastator, 2323, 2377. upon gillyflowers, 614. differentialis, 1423, 1549, 2323. Remedy for, 1920, 2232. Epidemic disease of 2340 Safe remedies for, 1948. Dimorphism in, 1889. Cabbages and fuchias. Beetles injuring, 1868. femur-rubrum, 147, 475, 1423, 1437, 1451, Cabinet pests. Means against, 1057. 1549, 1508, 1643, 2471. Caccecia fractivittana, 2475. Mite infesting, 146. rileyana, 1059, 1192. spretus, 147, 368, 873, 475, 536, 926, 929, rosaceana, 735. 1423, 1433, 1484, 1487, 1451, v-marked, 2363. 1452, 1456, 1473, 1481, 1482, Cactus lady-bird, 2119. 1488, 1492, 1493, 1494, 1495, Caddis-worms. Carnivorous habits of, 1839. 1501, 1521, 1527, 1532, 1538, Calandra gransria, 510, 586, 1259, 1673, 1683, 1742. 1541, 1549, 1555, 1557, 1562, · 1564, 1565, 1566, 1567, 1570, oryze, 510, 1673, 1683. remotepunctata, 15. 1571, 1572, 1573, 1575, 1576, Calaphis n. g., 27. 1577, 1578, 1582, 1585, 1588, betulella n. sp., 27. 1596, 1598, 1612, 1613, 1614, Calathus gregarius, 2418. 1615, 1617, 1643, 1669, 1674, California. Chinch-bug, 2431. 1693, 1959, 1994, 1996, 2003, Fruit insects in, 2245. 2005, 2007, 2040, 2044, 2057, Fuller's rose-beetle in, 1740. 2267, 2276, 2291, 2335, 2366. Grape Phylloxera in, 1727. (See also Locust; Rocky Grasshopper ravages in, 2323. Mountain locust.) Agricultural bearings of 1643. Injurious insects in. 2074. Aid of animals against, 1643. Interest in economic entomology in, 2053. Air-sacks of, 1950. Anatomy of, 1643. Introduction of Lestophonus iceryse into, 2522. Animals as foes to, 1643. Autumn flights in 1877, 1643. Locust ravages in, 1959. Birds that feed on, 1643. migratory locust, 2863. Brain of, 1959. Overflow bug in, 2099. Phylloxers in, 2140. Breeding-grounds of, 1643. Promotion of silk-culture in, 2006. British-American data, 1643. red scale, 2394. Characters, 1625. Winged Phylloxers in, 1895. Chronological history, 1625. Chronology of ravages, 1643, Californian locust, 1959. orange insects, 2373. 1950. Callidium antennatum, 50, 319. Classification, 1643. janthinum, 50, 319. Colorado data for 1877, 1643. Crops as affected by, 1643. Callidryas philes, 1271. Secondary sexual characters of, 1784. Dakota data, 1643. Callimorpha fulvicosta = C. lecontei. Destruction (devices for), 1643. lecontei, 1801. (means of), 1643. Development of, 2267. sp., 1871. as influenced vestalis = C. lecontei. Calliphora, 1390. by meteorovomitoria, 1266. logical condi-Callipterus, 1678. tions, 1969. Callochlora viridis = Parrasa chloris. Devices for destruction of Callocamia angulifera — Attacus angulifera. 1642.

Caloptenus spretus. Distribution of, 1635, 1643, | Caloptenus spretus. Remedies, 1643. 1959. (southern limita of), 1969. Effects that follow severe injury, 1643. Eggs, 1643. in same ground for two consecutive years, 1643. Enemies (invertebrate), 1643. (natural), 1625, 1969. (vertebrate), 1643. Fires (influence of prairie), 1648. Flight, 1939. (powers of), 1959. Flightain 1877 (autumn), 1643. from 1877 to 1879, 1959. General considerations, 1625. Habita, 1625, 1643, (at night), 1643. Histology, 1643, 1959. Injury (effects that follow severe), 1643. Invertebrate enemies of, 1643. Iowa data, 1643. for 1877, 1643. Kansas data for 1877, 1643. Losses caused by, 1643. Means against, 1959. Metamorphoses, 1643. Meteorological conditions (influence of, on migrations), 1960. 'Migrations, 1643, 1959. (influence of meteorological conditions on), 1959. in 1877, 1643, Minnecota data, 1643. for 1877, 1643. Missouri data for 1877, 1643. Native home, 1625. Natural enemies, 1625, 1959. Natural history, 1625, 1643. Nebraska data, 1643. for 1877, 1643. New Mexico data, 1643. Night babits of, 1643. Nomenclature, 1648. Parasites on eggs, 1612. Permanent breeding-grounds of, 1648. Powers of injury, 1625. Practical considerations, 1625. Prairie fires and, 1643. Previous visitations of, 1643. Ravages of (chronology of), 1643, 1959.

(east of the Mis-

sissippi), 1625. (relative to agri-

culture), 1959.

Statistics of losses caused by, 1643. in the temporary region after the departure of the return swarms, 1648. Texas data, 1967. for 1877, 1648. Transformations, 1625. travel. Direction in which young, 1643. Uses of, 1643. Utah data, 1643. Vertebrate enemies of, 1648. Visitation of, 1648. Weather (effects of), 1643. Wings (time of year when acquired), 1643. Young in country where they hatch, 1625. Calopteron and Photinus. Food of, 2358, reticulatum, 806. terminale, 806, 2358. Calosoma calidum, 721, 1274, 1314, 1857, 1672. scrutator, 1226, 1898. Calves. Lice on 889. Cambala annulata, 193, 286. multistriatus = C. annulata. Camel cricket, 1059. Camnula atrox, 1959. pellucida, 1959, 2044, 2813, 2363. Campoplex fugitiva ... Limneria fugitiva. Camptonotus scudderi, 2276. Campyloneura vitripennia, 1801. Canadian Entomologist (Review), 895, 479, 827. Canker-worm, 86, 172, 1021, 1066, 1127, 1863, 2012. again, 173. Army-worm and, 910. Cure for, 1426. Fall, 1423, 1482, 2238. Howell, A. J. Experience with the Spring, 2267. once more, 274. parasite, 717. Spring, 1423, 1482, 2238. worms, 91, 96, 202, 1482, 2267, 2270. not army-worms, 1304. Borers and, 941. Concerning, 2149. Remarks on, 1438. at the West, 1539. Canna flavida. Pamphila ethlius feeding on, 1897. Hesperid larva feeding on, 1897. Cannibal mites, 624. Cantharia cinerea = Macrobasia unicolor. vesicatoria, 2118. Canthon chalcites, 1172. hudsonias == C. lævis. lævis, 1172, 1386. Cape. Grape Phylloxers not at the, 1841. Capitalizing specific names, 2170, 2257. Capnia minima, 1141. Capsus oblineatus - Lygus pratensis. 4-vittatus - Pecilocapeus lineatus Carabidæ, 1537, 1726, 1788, 1760, 1956.

```
Carabidæ and Coccinellidæ to birds. Relations
                                                        Cecidomyia cupressi-ananassa n. sp., 1180.
                                 of, 2152.
                                                                      destructor, 6, 62, 122, 150, 157, 250, 426,
                               Food relations of,
                                                                        494, 711, 1506, 1581, 1587, 1685, 1690,
                                 2151.
                                                                        1721, 1787, 2008, 2031, 2235, 2267, 2311,
            Curculio-feeding habits of, 372.
                                                                        2365, 2396, 2398.
            Horn's classification of, 2051.
                                                                      impatientis, 852.
            Mode of oviposition of, 2341, 2355.
                                                                      leguminicola, 1690, 1721, 2238.
Care of entomological specimens. On the, 2290.
                                                                      orbitalis n. sp., 46.
Carnivorous. Are honey-bees, 2098.
                                                                      oryzæ, 1949.
                butterfly larva, 2359, 2361.
                                                                      q.-pilulæ n. sp., 41, 799.
                habits of caddis-worms, 1839.
                                                                      q.-symetrica, 799.
                propensities of plant-feeders, 1867.
                                                                      s. . wnigma n. sp., 46 = A carus s. - suig-
Carolina sphinx, 1262.
Carpet-beetle. The imported, 1699.
                                                                      s.-batatas n. sp., 46.
                 Trapping the, 1752.
                                                                      s.-brassicoides n. sp., 46, 518.
        bug. 1659.
                                                                      s.-cornu n. sp., 46, 197.
        pests, 1663.
                                                                      s.-coryloides n. sp., 46, 518.
Carpets in Texas. A tineid on, 2531.
                                                                      s.-gnaphalioides n. sp., 46.
Carpocapsa, 934.
                                                                      s.-hordeoides n. sp., 46.
             pomonella 57, 307, 330, 367, 373, 377.
                                                                      s. nodulus n. sp., 46.
               387, 454, 467, 523, 565, 769, 977, 1010,
                                                                      s.-rhodoides n. sp., 46.
                1022, 1056, 1059, 1254, 1301, 1311, 1320,
                                                                      s.-semen n. sp., 46 = A carus s.-semes.
                1329, 1334, 1308, 1378, 1394, 1398, 1476,
                                                                      s.-siliqua n. sp., 46, 1170.
                1496, 1609, 1666, 1695, 1721, 1887, 2037,
                                                                      s.-strobiliscus n. sp., 46, 197.
               2230, 2261, 2293, 2418.
                                                                      s.-strobiloides n. sp., 46, 518.
             saltitans, 1496, 1609, 2163, 2173.
                                                                      s.-triticoides n. sp., 46.
Carrying out the law, 1993.
                                                                      s. verruca n. sp., 46.
Carteria lacce, 2119.
                                                                      secalina, 2267.
         larrese, 2119.
                                                                      solidaginis, 798.
          mexicana, 2119.
                                                                      sp., 799, 963.
 Case-bearer. Dominican, 1363.
                                                                      tritici - Diplosis tritici.
                Lepidopterous, 260.
                                                                      tubicola, 1232.
                Walnut, 1311.
                                                                      v.-coryloides n. sp., 518, 1329.
Cassida, 94, 1238.
                                                                      v.-lituus = C. v.-viticola.
         aurichalcea = Coptocycla aurichalcea.
                                                                      v.-pomum n. sp., 403, 518, 1281, 1229,
         bivittata, 681, 993, 1127, 1510.
                                                                        1898, 2136.
         clavata = Coptocycla clavata.
                                                                      v.-viticola, 791, 1077, 1116, 1829.
         nigripes, 1127, 1510, 1697.
                                                         Cecidomyid gall on Crategus, 376.
         pallida = Coptocycla aurichalcea.
                                                                      in wheat, 18
         5-punctata n. sp., 767 = Physonota uni-
                                                         Cecidomyidæ, 294, 518, 799.
            punctata.
                                                                        on grape. Galls of, 111.
                                                                           salix. Galls of, 46, 197.
          texana, 2096, 2215, 2235.
 Cassididæ, 105, 767, 973, 1056, 1082, 1510.
                                                                        new species, 46.
 Castor bean. Locusts eat the, 1645.
                                                         Cecil's books of natural history (Review), 506.
 Catalogues and monographs, 1765.
                                                         Cecropia Cryptus, 1301.
 Catalogus coleopterorum, 1866.
                                                                   moth, 1112.
 Catalpa sphinx, 2119, 2291.
                                                                          caterpillars, 786.
 Cat-bird. Tipula eggs in stomach of the, 1735,
                                                                          cocoon, 635,
                                                                   silk-worm, 1311.
 Catcher. Dr. Hull's Curculio, 651.
                                                                               Remarkable abundance
 Caterpillar nuisance, 1995.
 Caterpillars named, 857, 1276.
                                                                   Tachina fly, 1311.
             Stinging, 1748.
                                                                   worm, 1215, 1249.
. Catocentrus n. g., 385.
                                                                          on elder, 1533.
 Cattle bot-fly, 2238.
                                                         CEDAR. INSECTS AFFECTING.
 Catydid eggs, 29, 247.
                                                             Callidium antennatum, 319.
          as an enemy to Doryphora 10-lineata, 313.
                                                                       janthinum, 319.
 Catydids. Food of, 143, 241.
                                                             Cerambycid, 1324.
 Cauliflower Botis, 2232.
                                                             Thyridopteryx ephemerseformis, 427.
 Cause of variation in organic forms, 2460.
                                                         Cedar longicorns, 319.
 Cave fauna of Kentucky. Coleopterous, 2033.
                                                         Celæna renigera = Hadena renigera.
 Cecidomyia, 1077, 1461.
                                                         CELERY. INSECTS AFFECTING.
              albovittata n. sp., 46.
                                                             Buprestis divaricata = Dicerca divaricata
              carbonifera, 1924.
                                                             Diceroa divaricata, 1063.
                                                             Tettigonid, 452.
              cornuta n. sp., 46.
              Cranberry, 194.
                                                             Trochilium sp., 1063.
              cratægi-bedeguar n. sp., 376.
                                                        Celery. Leaf-hoppers on, 452.
```

Celtis, 2208. Chelifer cancroides, 1884. Centennial insects, 1611. Chelymorpha argus, 767. Those, 1511. cribraria = C. argus. - Central America. Prevalence of screw-worm in, Chemical changes in the color of butterfly wings, Report on cotton crop and its Chenopodium hybridum. Lixus macer bred from, 2404 enemies in, 2343. Cephalomyia ovis = Œstrus ovis. Cherished bracon, 1423. Cerambycidse, 1324, 2267. Chernetidse, 1884. Food habits of 1902. Cherry aphis, 960. Cerastophorus cinctus = Chion cinctus. borer. Flat-headed, 2267. Ceratina dupla, 1171. Caterpillars nest on wild, 242. Ceratocampa imperialis = Eacles imperialis. Cocoon on wild, 259. regalis - Citheronia regalis. CHERRY. INSECTS AFFECTING. Ceratomia catalpse, 2119. Acarid, 1265. hageni, 2119. Aphis cerasi = Myzus cerasi. Cercopidida, 882. Attacus polyphemus, 1164. Cercopsylla n. g., 2272. promethes, 257. sideroxyli n. sp., 2272. Dicerca divaricata, 2267. Cereals. Insects injurious to, 2238. Eriocampa cerasi, 1222. Ceresa bubalus, 415, 1323, 1329, 2107, 2195, 2235. Hyphantria cunea, 242. Cermatia forceps, 752, 764, 1154. textor = H. cunes. Cerococcus quercus, 2119. Mysus cerasi, 960, 1067, 1207, 1251. Ceropales rufiventris n. sp., 375, 543. Œcanthus niveus, 1503. Ceroplastes cirripediformis, 2450. Scolytus rugulosus, 1940. Selandria cerasi = Eriocampa cerasi. n. sp., 1964. Ceropsylla n. g., 2272. Telea polyphemus = Attacus polyphemus. Cherry plant-lice. Destroying, 1207. sideroxyli n. sp., 2272. Ceroptres ensiger, 41. and their foes, 1251. tree borers, 1063. ficus, 745. inormis, 41. plant lice, 1067. petiolicola, 41. trees. Minute borers in, 1940. Cerotoma caminea, 2418. Slug on pear and, 1222. Cerura multiscripta n. sp., 1411. twigs. Berry and, 1503. Cetonia, 2105. Chester onion pest, 2319. Chestnut. Gall on, 353. inda = Euphoria inda. Cherocampa pampinatrix = Ampelophaga myron. Gregarious worms on horse, 1192. Chætochilus pometellus, 874. Phylloxera sp. on, 353. Chatophorus, 1678. Chicken mite, 1308. Chickwood. Geometer, 1059, 1154. Chætoptria hypericana, 2076. Chafer. Melancholy, 522. Hæmatopis grataria on, 1059, 1154. Rose, 1329. Chilo orysmellus n sp., 2119, 2120. Chalcid egg parasites. Efficacy of, 2115. Chilocorus bivulnerus, 38, 188, 301, 417, 730, 731, fly, 1643, 1959. 1534, 1798, 1883, Chalcididso, 128, 384, 2115, 2343. CHINA-TREE. INSECTS AFFECTING, 1964. Joint-worm not one of the, 49. Atta fervens, 1964 Chalcis albifrons n. sp., 6 = Smicra albifrons.Ceroplastes sp., 1964. Lecanium sp., 1964. fly, 1059. Chinch-bug, 584, 886, 891, 894, 1127, 1423, 1506, 1681. Inflating, 1059. marise n. sp., 1112 = Spilochalcis marise. 2018, 2039, 2048, 2271, 2368, 2418, Mary, 1811. and army-worm prospects, 2087. ovata, 2343. Bogus, 1127, 1423. Chalcophora virginiensis, 2267. Bug allied to, 90. Chalepus trachypygus, 278, 1911, 1949, 1973, 2119. Bugs on grape-vines mistaken for, Chambers, V. T. Obituary, 2218. 398. Change of habit, 2096. Diseases of, 2143. False, 1329, 1423, 2291, 2418. in vegetation caused by locusts, 1495. in New York, 2237, 2252, 227L Chapin's apple-leaf sewer, 1721. Chariesterus antennator, 847. not in seed grain, 888. Chauliodes. Larval characteristics of, 1652. notes, 2119, 2291. pectinicornis, 712. once more, 890, 894. bugs, S. A. Forbes. Experiments on, 2165. rastricornis, 712. Genuine vs. bogus, 1402. Chauliognathus pennsylvanious, 372, 374, 1959. Locusts vs., 1482. Cheese-fly and blow-fly, 1266. Other insects affecting, 1703, Chion cinctus, 55. skipper, 1700. Chienaspis euronymi, 2403.

Chionapsis furfurus, 2, 188, 264, 345, 372, 878, 579, Cicada. Song notes of the periodical, 2334. in southeastern Indiana, Butler, A. W. 951, 1008, 1040, 1061, 1152, 1564, 1968, 2395. The periodical, 2394. pinifolii, 97, 1329, 2079. Massachusetta. Periodsalicis, 378. ical, 2216. Sting of the, 371, 872, 407. nigræ n. ep., 373 = C. salicis.Chipman, A. J. Locust notes in 1880, 2267. tibicen, 920, 1034, 1546. Chiro.omus, 167, 625. tredecim = Tibicen tredecim. Twigs punctured by periodical, 1055. Chlenius impunctifrons, 2340. Chlamys plicata, 1363. years, 51: Cicadas, 26, 2311. Chlorion cæruleum, 543. Chloroneura n.g., 22 =Empossos. Cotton-worms and, 1997. abnormis n. sp., 22 = Empoasca ab-Destroying, 2305. normia. and walking-sticks, 920. malefica n. sp., 22 = Empoasca viri-Cicadidæ, 431. descens. Cicadula exitiosa, 1766. maligna n. sp., 22 = Emposeca obtusa. Cicindela repanda, 1763. Chloroperla brunnipeunis n. sp., 24. 6-guttata, 347. fumipennis, 24. vulgaria, 628. nana n. sp., 24. Cicindelidæ, 719. Chlorops, 1461, 2060. Cimbex americana, 699, 1380, 1514, 2291, 2333, 2418. Chorinsous cariniger, 385. laportei = C. americana. Chortophaga viridifasciata, 1566. Cioidse, 1937. Chronology of locust ravages in 1880-'81, 2267. Circotettix maculatus n. sp., 1959. Chrysididse, 375. Circular No. 1, United States Entomological Com-Chrysobothris, 934. mission, 1575. femorata, 2, 6, 55, 352, 377, 379, 537, 2, United States Entomological Com-555, 578, 634, 870, 1059, 1227, 1250, mission, 1576. 1816, 1367, 1423, 1454, 1472, 1475. 9, U.S. Dept. Agric., Div. Ent., 2303. 1513, 1640, 2267. Cirrhophanus triangulifer, 2189. Chrysochus auratus, 735, 790. Cirrospilus esurus n. sp., 1712 = Tetrastichus esu-Chrysomela bigsbyana, 323. rus. cyanea == Gastroidea cyanea flavicinctus n. sp., in Lintner's first philadelphica, 323. report. (See foot-note 1, p. 379.) scalaris, 45. Cis fuscipes, 1737. Chrysomelidse, 323, 2229. Citheronia regalia, 425, 629, 702, 775, 803, 869, 1087, Chrysopa plorabunda, 584, 1127. 1275. sp., 372, 538. Clemens, B. Obituary, 223. Cicada, 189, 213, 372, 431. Cleonus sp., 1755. alias the seventeen-year locust. Period-Cleora. Evergreen, 2363. ical, 1159, 1971, 1979. Cleridæ, 1363. Belated individuals of the periodical, 648. Clerus nigripes = C. 4-guttatus. canicularis = C. tibicen. 4-guttatus, 200. cassini = Tibicen cassini. Click-beetle larvæ, 1643. Fungus in, 1809. beetles, 84, 351, 1625. in Iowa. Seventeen-year, 1737. Climate on Cicada septendecim. The influence of Little, 1242. 2329. in Massachusetts. Periodical, 2321. for Doryphora, 1412. Note on the periodical, 2318. on prolongation of life of insects. Effects our first brood established. Periodical, of. 1129. 707. Climbing cut-worms, 1069. parvula = Melampsalta parvula. Clisiocampa americana, 66, 77, 112, 240, 277, 336, 350, Periodical, 474, 527, 619, 1059, 1159, 1489, 363, 642, 819, 1181, 1301, 1352, 2238. 1624, 1836, 2014, 2216, 2312, 2314, 2315, 2318, disstria, 240, 291, 328, 363, 645, 646, 688, 2320, 2363. 1127, 1181, 1200, 1301, 1331. Premature appearance of the periodical, sylvatica = C. disstria. 2330, 2326. Cloantha derupta, 2410. pruinosa = C. tibicen. Clœ dubia n. sp., 24. in savin-twigs. Eggs of the periodical, ferruginea n. sp., 24. fluctuans n. ep., 24. septendecim = Tibicen septendecim. mendax n. sp., 24. Influence of climate on 2329. Close of the first volume, 701. Instinct of, 2144. Clostera americana = Ichthyura inclusa. Some popular fallacies re-Clothes moths, 70, 1633.

observed in the United States,

garding, 2367.

Seventeen-year, 1034, 1098, 1698, 2312, 2814.

Clover. Army-worm notes and inquiries, its | Codling moth, jarring down infested fruit, 1318. work on, 1877. beetle. Flavescent, 2894. Bumble-bees vs. red, 2376. enemy. New imported, 2028. hay-worm, 1363, Army-worm vs., 2090. CLOVER. INSECTS AFFECTING, 1690, 2050. Asopia costalia, 83, 675, 948, 1132, 1363, 2090. Cecidomyia leguminicola, 1690, 1721, 2238. Drasteria erechtea, 1459. Hylesinus trifolii, 1690, 1721, 1777, 1846. Leucania unipuncta, 1877. Oscinis sp., 2394, Phytonomus punctatus, 2010, 2028, 2061, 2119. Pyralis olinelis = Asopia costalis. Sitones flavescens, 2394. Clover-leaf beetle, 2119. weevil. Further notes on, 2061. New imported enemy to, 2010. root borer, 1721, 1846, 2512. seed midge, 1721, 2238, 2291. stem maggot, 2394. Webster, F. M. Insects affecting white, weevil, 1777. worms, 83, 675, 948, 1132. Clubbed tortoise beetle, 1127, 1558. Larva of, 1238. Cluster flies. Notes on, 2174. fly, 2128. Clydonopteron n. g., 1929. tecom# n. sp., 1929. Clytus capræa = Neoclytus capræa. Oak, 2267. pictus = Cyllene pictus. robinise = Cyllene robinise. speciosus = Plagionotus speciosus. Coocid mistaken for a gall, 1972. Coccidso, 1329, 1883, 1968, 2088, 2137, 2235, 2520. Means against, 104, 218, 264, 301, 487, 508, 944, 2372, 2373, 2403. Monograph of, 1828. Coccinella munda = C. sanguinea. sanguinea, 6. Coccinellidæ, 69, 822, 323, 639, 894, 1883, 1956, 2119. to birds. Relations of Carabidæ and, 2152. Food relations of Carabids and, 2151. Habits of, 125. Coccotorus scutellaris, 33, 84, 42, 254, 878, 377, 1301, 1378. Coccus, 159. harrisii = Chionaspis furfurus. Cochylis hilarana, 1059. Cocklebur. Rhodobænus 13-punctatus on, 1301. Sphenophorus, 1301. Cockroach eggs, 573. Cockroaches. Means against, 1197. Cockscomb elm gall, 1884. Codling moth, 57, 307, 378, 1022, 1059, 1127, 1254,

1329, 1334, 1368, 1476, 1677, 2119.

2293, 2418.

heresies, 1398.

again, 1301, 1311.

Michigan apples and, 1677. Wier's trap, 1334. Cœliodes inæqualis = Oraponius inæqualis. Cœlodasys unicornis, 749, 880, 1280. Coffee-borer, 498. Cohn, F. The Hessian fly in Silesla in 1869, 2267. Colaspis, 872. barbara, 1301. flavida, 129, 133, 135, 231, 993, 1301, 1311, Grape-vine, 231, 1301. Cold applied to chrysalids of butterflies. Effects of. 1778. on Curculio. Influence of extreme, 1885. and hibernating apple-worms. Severe, 2037. on insects. Effects of severe, 1818. the scale insects of the orange in Florida. Voyle, J. Effects of, 2277. Coleophora, Apple, 1721. malivorella n. sp., 1721. Coleoptera, 1329, 1904, 2267. Classification of North American, 2025 Myrmicophilous, 2105. of North America. The new classification of, 2175. Parasitic, 2353. Willow-galls made by, 46. Coleopterorum. Catalogus, 1866. Coleopterous cave fauna of Kentucky, 2083. larvæ. Remarkable, 40. Colina. The genus, 2259. philodice, 461. Collecting and preserving insects, 881, 1067. Colonies. Silk-culture in the, 2278. Colopha ulmicola, 27, 187, 151, 518, 1384, 1678. Colorado beetle poisonous f Is the, 1405. cabbage flea-beetle, 2291. grasshopper, 502. potato-beetle, 21, 991, 1069, 1311, 1839, 1863, 1428, 1482, 1558, 1570, 2291, 2294, 2418. abroad, 1864, 1407, 1481. again, 1301. Bogus, 1558. in Europe, 1597. Experience with the, Inquiries concerning the, 1491. Insect enemies of the, Mite parasite of the, 1505. Native home of the, 1462. in New Hampshire, 1859. New York, 1379. Onward march of the, 1218. poisonous? Is the, 1483. Poisonous qualities of the, 1425. Specific for, 1490. potato-bug, 293, 334, 866.

Bogus, 519.

Colorado potato-bug. Poisonous qualities of the, Cordulia! molesta n. sp., 39. 1101. Cordyceps, 332, 478. bugs. To destroy, 650. ravenelii, 594, 640. Coloration in insects. Unity of, 50. Corous tristis = Anasa tristis. Come. Still they, 908. Corimelana pulicaria, 354, 637, 741, 1046. Common Curculio and its allies, 254. Corn bill-bug, 2119, 2291, 2418. flesh-fly, 1423, 1570, 1625, 1643. borer, 754. longicorn pine-borer, 2291. Confessing the, 256. May beetle. Eggs of, 1829. Curculio, 1042. Commoner insects. Notes on our, 1802. cut-worm, 1059. Companion wheat-fly, 2394. ear-worm, 2238. Complimentary, 504. feeding Syrphus-fly, 2422. Composite. Antistrophus confined to the, 1131. fields. Beetles in, 2317. CORN. INSECTS APPECTING, 2119. Compsomyia macellaria, 2256. Agonoderus pallipes, 2317. Comstock on the classification of insects, 2390. Diaspinse, 2227. Anthomyia zeas, 657, 1059, 1066. J. H. Report on miscellaneous insects, Aphis maidis, 23. 2119. Butalis cerealella, 1232. Conchylis erigeronana n. sp., 1968. Calandra granaria, 1683. oryzse, 1683. conotherana n. sp., 1968. Cetonia inda = Euphoria inda. Conclusions. Jumping at, 253. Cone-nose. Blood-sucking, 795, 854. Cut-worm, 1073. Diabrotica longicornia, 1905, 2187. Confessing the corn, 256. Empretia stimulea, 754. Confounding friend with foe, 1378. Conocephains, 2267. Ruphoria inda, 1655. Conorhinus. Blood-sucking, 2291. Gortyna nitela, 754. sanguisugus, 497, 795, 854. sp., 138. Conotrachelus, 984. Hadena sp., 1047. anaglypticus, 372. Heliothia armigera, 192, 1301, 1664, 2119, 2238. crategi n. sp., 35, 358, 1301, 1358. Ipe fasciatus, 4. geminatus, 84. 4-signatus = I. fasciatus. nenuphar, 6, 83, 42, 50, 98, 188, 237, 254, Lepidopteron, 162. 290, 329, 367, 372, 373, 377, 387, 439, Mesograpta polita, 2422. 487, 501, 566, 589, 620, 649, 680, 703, Noctuid, 181. Pempelia lignosella, 2119. 873, 930, 958, 978, 1056, 1059, 1062, 1069, 1071, 1129, 1185, 1173, 1177, Sphenophori, 2030. 1201, 1214, 1258, 1801, 1325, 1857, Sphenophorus robustus, 2119. 1358, 1372, 1477, 1685, 2230, 2296. sculptilis, 337, 1042, 1301. puncticollis n. sp., 34 = C. geminazeas = 8. sculptilis. tus. Corn insects. Potato and, 138. Controlling sex in butterflies, 1352. Larval habite of Sphenophori that attack, Convolulacem, Insecta, 1238. 2030. Cook on insecticides. Review of, 2501. A new enemy to, 1905. Cooke's injurious insects of the orchard. Review foe of the, 337. of, 2245. insect foe of green, 1655. Copidryas gloveri, 2486. pest. Unknown, 1973. Copper underwing. Spattered, 1301. plant louse, 2418. Copris carolina, 416. root Diabrotica, 2187. Coptocycla, 1238. louse, a new enemy to the corn, 23. aurichalcea, 681, 786, 802, 1127, 1510. rustic, 1059. bivittata = Cassida bivittata. Serious injury to stored and cribbed, 1682. clavata, 693, 742, 767, 1238, 1558. Sphenophorus, 1301. guttata, 1127, 1510. stalk borer. Smaller, 2119. nigripes = Cassida nigripes. Webster, F. M. Insects affecting, 2418. Coquillett, D. W. Remedies for cottony cushionworm, 192, 820, 1301. scale, 2394. Worm in 181. Report on the gas treatment for worms, 162, 1047, 1664, 2119. scale insects, The war on, 1522. 2418. Corrected. A few errors, 977. locusts of the San Correction, 939. Joaquin Val-Entomological, 1328. ley, California, Phylloxera, 1855. 2363. Corrodentia, 2267. production and

Corticaria pumila, 2418.

Corycia vestaliata, 744.

manufacture of buhach, 2364. Corydalus corautus, 130, 308, 473, 1329, 1570, 1584, COTTONWOOD. INSECTS AFFECTING-Continued. 1627, 1652, Pemphigus vagabundus, 446. Larval characters of, 1652. Plectrodera scalator, 1525. Corynetes rufipes == Necrobia rufipes. Saperda calcarata, 71. Corythuca ciliata, 200, 400. Cottonwood killer, 1654. Coscinoptera dominica, 1363, 2093. leaf beetle. Streaked, 2291, Cossidse, Bailey, J. S. The tree borers of the galls, 713. family, 2253. Unsightly galls on, 446. Cossus robiniæ, 37, 245, 967, 1003, 1121. Worms on, 1459. Cotton-worm, 1127, 1363, 1649, 1702, 1719, 2077, 2119. Cotalpa lanigera, 40. Cotton belt. Smith, E. A., 2343. 2130, 2164, 2295. boll-worm, 2238. articles in the News, 1719. bolls. Euryomia melancholica vs., 2451. Barnard, W. S. Tests of machinery in Brazil, Branner, J. C. Insects injurifor destroying, 2258. ous to, 2277. Bulletin on the, 1786. caterpillar, 1353, 1801. Chrysalids dug up in corn-fields miscaterpillars in Brazil, Branner, J. C., 2343. taken for those of the, 1790. convention. Address at Atlanta, 2038. damage in 1881, 2119. crop and its enemies in Mexico, Central To destroy the, 1338. and South America, and the West Indestroyer. 'Best, 1882. dies. Report on the, 2343. Early appearance of, 184. culture and the insects affecting the plant Facts about, 1657. at Bahia, Brazil, 1788a. How to manage the, 1831. destroyers, 1886. use Paris green for the, 1544. field mistaken for those of the cottoninjury in regions overflowed by Misworm. Chrysalids dug up in, 1790. sissippi. Protection from, 2119. insects, 636. investigation, 1833. COTTON. INSECTS AFFECTING. Jones, R. W. Observations and ex-Aletia argillacea = A. xvlina. periments on, 2164. xylina, 646, 1127, 1338, 1353, 1363, 1369, moth hibernation, 1953. 1649, 1657, 1702, 1719, 1721, 1722, 1736, Not the, 1778. notes from Vera Cruz, Mexico. In-1758, 1769, 1814, 1845, 1852, 1882, 1886, 1997, 2088, 2064, 2067, 2119, 2180, 2155, teresting, 1845. 2164, 2295, 2343. Orange insects and, 2067. Anomia xylina = Aletia xylina. parasites, 1712. Euphoria melancholica, 2451. Parasites bred from, 1722. Euryomia melancholica = Euphoria melan-Poisoning devices for the, 2119. cholics. Possible food-plants for, 2064, 2119, Gryllus sp., 2384. Heliothia armigera, 646, 1353, 2288, 2343 in prairie belt. First appearance of, Hyperchiria io, 2343. Saturnia io = Hyperchiria io. question of hibernation settled, 2119. Spilosoma acræa, 2343. A remedy for the, 1369. Theclapmas † 1872. Report on, 2164. Cotton insects, Jones, R. W. Report on, 2343. in south Texas in 1883. Anderson, W. J. Report on, 2343. E. H., 2253. Sowing cotton-seeds in hot-beds as a Kerosene as a means means against the, 1772. against, 2164. moth. Habits of, 1758. in the United States, 1769. Hibernation, 1728. Cotton-worms, 2291. plant. Insects affecting, 1721. and Cicadas, 1997. Cottony cushion-scale, 2394, 2418. planters. Suggestions to, 1832. Coquillett, D. W. Reme-Plums and, 1477. dies for, 2394. seeds in hot-beds. Sowing, 1772, -Koebele, A. Remedies for, squares. Butterfly larvæ injurious to, 1872 2394. maple-scale, 1816, 2291. Cottonwood borers, 1525. scale insect on maples, 1515. dagger, 1127. Couper's thorn leaf gall. Mr., 376. A foe to, 1834. COTTONWOOD. INSECTS AFFECTING. Crab-apple borers, 602. Acronycta lepusculina, 1127. Crack-jaw names, 592. Aphid, 495. Crambus. Vagabond, 2119. vulgivagellus, 1990, 2029, 2119. Drasteria erechtea, 1459.

Lina scripta, 1654, 1834, 2291. Pemphigus populicaulis, 446, 713. Cranberry fruit worm, 2291, 2849.

galls, 194.

CRANBERRY. INSECTS AFFECTING. Ctenucha latreilliana = C. virginica. Acrobasis vaccinii, 2291. virginica, 1241. Cecidomyia sp., 194. Cucumber beetle Striped, 773, 1127, 2258. Phytopus sp., 2493. Cranberry leaf-folder, 2291. Smith, J. B. Report on insects affecting the hop and the, 2277, 2291. Crane-flies, 1278. fly larva, 512. Craponius insequalis, 367, 378, 1059. Cratagus. Insects on, 376. Cratonychus incertus == Melanotus incertus. Cratypedes putnami, 1969. Culex, 625. Cremastochilus, 2105. mosquito, 2185. Creophilus villosus, 805. pipiens, 2071. Crepidodera brevis, 1782. Culicidæ, 603, 2162. cucumeria, 857, 401, 1056, 1059, 1558, 1782. Distribution of species of, 1782. helxines, 356. parvula, 1782. Crescent. The little Turk and his, 329. Cribbed corn. Serious injury to, 1683. Cricket. Grapes out by a tree, 414. in Louisiana. Destructive, 2884. in raspberry canes. Eggs of tree, 2195. 1088. Tree, 251, 953, 961, 999. Snowy tree, 2291. Western, 1959, 2267. Bruner, L. Report on, 2267. catcher, 875. Histology of, 1959. Crickets, 143, 1363. Corp. 1042. Mole, 1270. Crioceris asparagi, 524, 2185. erv. 1178. 12-punctata, 2185. Crippled moths, 729. Critic oriticised, 222, 942. Criticism. Friendly, 1307. Crop gall-gnat, Wagner, B. Observations on the new, 2267. Croton-bug as a library pest, 1717. apple, 1228. Cruel bug-hunters, 440. Crumpler. Apple-leaf, 949. in Georgia, Rascal, 1962. mistaken for Curculio. Apple-loaf Rascal leaf, 311, 373, 1811, 1580 Cryptocentrus n.g., \$85 = Mesoleius. Plum, 1372. Cryptorhopalum, 1746. Cryptus albicaligatus n. sp., 385. albisoleatus n. sp., 385. annulicornis = Pimpla annulicornia. occropia, 1811. cinctipes n. sp., 385. conquisitor = Pimpla conquisitor. Quince, 1801. inquisitor = Pimpla inquisitor. remedies, 566. junceus = Linoceras junceus. nigricalceatus n. sp., 885. nuncius, 1112. picticoxus, 385. pleurivinctus = Pimpla conquisitor. rhomboidalis n. sp., 885. rufifrons n. sp., 385. samis, 2343. Curculios on pears, 1858. Cteniscus albilineatus n. sp., 885.

ornatus, 385.

bug. Striped, 148. fica-beetle, 1050, 1558. CUCUMBER. INSECTS AFFECTING. Diabrotica vittata, 148, 773. Eudioptis nitidalis, 808, 843. Phakellura nitidalis = Eudioptis nitidalis. Cucumber. Worm boring in, 808, 842. Cucurbitaceous vines. Insects affecting, 1137. Cupuliferse. Cynips confined to, 1131. Curculio, 958, 978, 1063, 1080, 1127. again. Comparative scarcity of the, 782. and its allies. Common, 254. Apple, 1176, 1361. leaf crumpler mistaken for the, anout-beetle or four-humped, and bark-lice, 487. breeds in apple. Plum, 1214. Carabida feeding on, 372. Hull's, 372, 651, 875. extermination possible. Great discov-Fighting the, 237. Flea-beetles and, 873. flice by night. How the, 706. Four-humped, 1068. Gas-waste vs., 549. go underground to transform! Does the humbug. Another new, 589. A new, 680. Influence of extreme cold on the, 1385. injurious to grape, 267. Means against the, 290. natural history and how to catch him. parasite. Porison, 1391. Sigalphus, 1801. Parasites of the plum, 1751, 1796. Paris green for the, 1258. Plum, 1059, 1129, 1301, 2296. To protect plums from the, 1971. remedy. The new, 1177. Ransom, 1201. scarcer than last year, 620. Is the, 640. trap. Smith's patent, 969. Use of the word, 1069. Curculionide. Monograph of, 1828. Currant borers, 481. bush borer, 974.

Current, Current-worms and black, 1979. Four-lined leaf-bug on, 715. and gooseberry worms, 1570. CURRANT. INSECTS AFFECTING. Ægeria tipuliformis, 340, 481, 974, 2238.

Alcatho caudatum, 481.

Amblycorypha oblongifolia, 1157.

Aphis ribis = Mysus ribis.

Capsus 4-vittatus = Pocilocapsus lineatus.

Eliopia ribearia - Eufitchia ribearia.

Eufitchia ribearia, 176, 364, 772, 877, 1234.

Mytilaspis pomicorticis = M. pomorum. pomorum, 973.

Mysus ribis, 304.

Nematus ribesii, 78, 840, 364, 380, 381, 772, 1081,

1224, 1570, 1696, 2238.

ribia - N. ribeaii.

.ventricosus= N. ribesii.

Phylloptera oblongifolia - Amblycorypha oblongifolia.

Poscilocapsus lineatus, 715.

Pristiphora grossulariss, 176, 364, 772, 1570,

Procopis affinis, 1171,

Psenocerus supernotatus, 481.

Silvanus surinamensis, 1260.

Tenthredinid, 136.

Trochilium tipuliformis = Egeria tipuliformis.

caudatum = Alcatho caudatum.

Current plant-lice, 304.

louse, 322

stalk-borer, 2238.

worm, 877, 1204.

Imported, 1127, 1570, 2238.

Native, 1570, 2238.

Pear-slug and, 1031.

worms, 364, 882, 1696.

Destructive, 78.

Gooseberry and, 772.

Currants. Beetle in dried English, 1260.

Green worms on gooseberries and,

136.

Crahion-scale. Cottony, 2394, 2418.

Coquillett, D.W. Remedies for, 2394.

Koebele, A. Remedies

for, 2394.

Cushions. Ravages of moths in, 1731.

Cussed red-leg, 1598.

Cuterebra buccata, 526.

Cut-worm, 868,992.

Climbing, 1059.

Corn, 1059.

Dark-sided, 281, 1059, 2291, 2968.

Dingy, 1050.

eggs, 1095.

Glassy, 1059, 2291, 2394.

Granulated, 2291.

Greasy, 1059, 2291, 2418.

lion, 1059, 1314.

moth. Eggs of, 607.

Shagreened, 2291.

Small white bristly, 1059.

Speckled, 1059, 2291,

Cut-worm. Variegated, 1059, 2291.

W-marked, 1059, 2291.

Western striped, 1059.

Wheat, 1059.

Yellow-headed, 1059.

Cut-worms, 8, 85, 321, 658, 914, 1059, 1315, 2288, 2291.

Cabbage, 2291.

A chapter on, 964.

Climbing, 321, 359.

destroying recently sown wheat, 455.

Means against, 226, 312, 1059, 2387.

Novel facts about, 901.

originate! How, 787.

and other insects? Can land be in-

sured against, 801.

parasites, 849.

Remedy for cabbage, 2291.

Ridding the ground of, 2810.

severing cabbage-plants, 626.

Smilex injured by, 1941.

from the stomach of a robin, 1873.

Tree, 228, 281. Cybister fimbriolatus, 750, 816.

Cybocephalus californicus, 2079.

Habits of, 2079.

nigritulus, 2079.

Cycloosphala immaculata, 1236.

Cyclone nossle, 2327. Cylindrical Orthosoma, 1059.

Cyllene, 560.

decorps, 50.

pictus, 45, 50, 55, 89, 269, 308, 560, 828, 941,

robinise, 37, 45, 50, 55, 141, 308, 472, 560, 828,

921, 941, 967, 1196, 1250, 1288, 1516.

Cynipid gall on oak twigs, 1822.

Cynipida, 518, 799, 821, 2072, 2348.

Dimorphism in, 41, 1980.

Illinois, 41.

Important work on, 2125.

List of, 41.

New classification, 41.

Synoptic table of genera, 821.

Cynipa, 1181, 1559.

aciculata = Amphibolips spongifica.

bicolor = Rhodites bicolor.

confined to Cupuliferse, 1131.

Dimorphism in, 41. feoundatrix, 1606.

gallæ-tinctoriæ, 1560.

n. sp., 1822.

q.-californica = Andricus q.-californica.

a.-decidus, 1925.

q.-erinacei n. sp., 41 = Acraspis q.-eriracei.

q.-flocci n. sp., 41 = Andricus q.-flocci.

q.-forticornis n. sp., 41 = Biorhisa forticornis.

q.-frondoss, 821, 774, 1666.

q.-glandulus n. sp., 1606.

q.-inanis = Amphibolips q.-inanis.

q.-mamma n. sp., 518 = Holcaspis mamma.

q.-mellaria n. sp., 1942.

q.-operator = Andricus operator.

q.-operatola, 1352.

Deloyala = Coptocycla.

clavata = Coptocycla clavata.

Cynips, q. pilulæ n. sp., 41 = Cecidomyia q.-pilulæ. Deltocephalus xanthoneurus, 2114. Delusion. A popular, 525. q. podagræ n. sp., 41 = Andricus q. poda Dendrotettix n. g., 2410. q.-prunus n. sp., 518 = Amphibolips q.-pruquercus n. sp., 2410. Deræocoris rapidus = Calocoris rapidus. nna. Dermanyssus avium, 1308. q.-punctata = Andricus q.-punctata. q.-saltatorius=Neuroterus q.-saltatorius. gallinæ, 1308. q.-spongifica-Amphibilops q.-spongifica. Dermatobia hominis, 497. seminator == Andricus seminator. Dermatoptera, 2267. Dermostes lardarius, 72, 726, 1190, 1240, 1703. sp., 799. terminalis, 1448. Toothed, 2363. Cynthia atalanta = Pyrameis atalanta. vulpinus, 2368. Dermestidæ, 1746, 2133, 2180. cardui - Pyrameis cardui. Cypress. Cecidomyia cupressi-ananassa on, 1180. Means against, 56, 246, 462. Deserved honor, 2124. galls, 1180. Cyrtoneura a parasite or a scavenger! Is, 2102. Desmia maculalis, 468, 993, 1056, 1158, 1301, 1569, . stabulans, 2102, 2343. Cytoleichus sarcoptoides, 2135, 2157. Desmocerus palliatus, 358. Destructive larvæ, 722. Cytophyllus concavus, 516, 569, 1005, 1157, 1363, locusts. Literature of, 1969. Dactylosphæra caryæ-semen n. sp., 373 = Phyllox-Devastating locust, 2963. era caryæ-semen. Devil's darning-needle, 1709. vitifoliæ = Phylloxera vastatrix. riding-borse, 457, 1059. Dactylosphæridæ, 1301. Dexia rustica, 2260. Dagger. Cottonwood, 1127. Dexides. Larval habits of, 2260. Smeared, 13-1. Diabrotica. Corn-root, 2187. Dahlia and aster stalk-borer, 949. 12-punctata, 123, 687, 2418. Gortyna nitela injuring, 862, 940, 1009. Long-horned, 1905. stalk-borer, 862, 1009. longicornis, 169, 1905, 2187. Daisy as an insecticide. Ox-eye, 1861. Twelve-spotted, 687, 1127, 2418. Dakota. Locust flights in, 2007. vittata, 123, 148, 175, 255, 772, 887, Dakruma convolutella, 1059. 2236, 2238, 2260. Dana, J. D. Criticism of the entomological views Diapheromera, 2267. of, 44. femorata, 144, 448, 920, 1111, 1395, and his entomological speculations. Pro-1701, 1709, 1721, 2276. fessor, 82. velii n. sp., 45. Danaidæ, 613, 1301. Diaspinse. Comstock on the, 2227. Danais archippus, 406, 461, 613, 1285, 1301, 1340, 1535, Diaspis harrisii — Chionaspis furfurus. 1622, 1635, 1711, 1756, 1770, 2052. rosæ, 1303. Dangerous looking, 814. Diastrophus, 318, 1131. Dapsilia rutilana, 1713, 1721. cuscutæformis, 609. Darapsa myron = Ampelophaga myron. nebulosus, 263, 610, 1121. Dardanelles. Sareophaga lineata destructive to Dicerca divaricata, 871, 1063, 2267. locusts in, 2075. Didictyum n.g., 1749 — Hexaplasta. Dark grape-worm, 763. zigzag n. sp. = Hexaplasta zigzag. sided cut-worm, 1059, 2291, 2363. Diedrocephala flavipes n. sp., 1766, 1767, 2363. Darning-needle. Devil's, 1709. Differential locust, 1423, 2363. Darwin's work in entomology, 2129. Digger wasps, 309, 1643, 1959. Dascyllid in stomach of black bass, 1792. Dilar in North America, 2023. Dasyllis thoracica, 1226. Dilophus, 2076. Datana contracta, 50. Diminished Pezomachus, 1127. ministra, 50, 182, 151, 165, 467, 922, 1045, Dimmock's inaugural dissertation, 2082. 1229, 2222, 2333. Dimorphic asexual females. Lichtenstein's theory as to, 2072. Decatoma dubia n.sp., 384 = var. of varians. excrucians n. sp., 384 = var. of nigri-Dimorphism in Cynipidse, 1980. Cynips, 41. hyalipennis n. sp., 384. Dytiscidæ, 28. nigriceps n. sp., 384. locusts, 1889. nubilistigua n. sp., 384. Dingy cut-worm, 1059. simplicistigms n. sp., 384. Dinoderus pusillus, 2104. sp., 536. as a museum pest, 2104. varians n. sp., 384. Diones, 1499, 1688. Defense. Inexpert, 2476. Diphryx prolatella, 2120. Delicate long-sting, 1329. Diplax rubicundula, 2007.

Diplodus luridus, 872.

Diplosis annulipes n. sp., 46.

Diplosis atricornis n. sp., 46. Doryphora, 10-lineata, 1412, 1423, 1425, 1431, 1447, atrocularis n. sp., 46. 1450, 1462, 1469, 1480, 1482, decem-maculata n. sp., 46. 1490, 1491, 1505, 1519, 1524, found in phylloxera galls, 1901. 1528, 1558, 1564, 1570, 1593, grassator, 2150. 1597, 1610, 1639, 1672, 1714, helianthi-bulla n. sp., 197. 1859, 1948, 1993, 2081, 2235, nigra, 2363, 2393. 2291, 2294. Pear, 2363. · Fire cure for, 121. pyrivora n. sp., 2393. juncta, 48, 50, 372, 401, 519, 1056, 1059, septem-maculata n. sp., 46. 1431, 1450, 1558, 1672, 2096, 2215. tritici, 109, 110, 142, 216, 280, 292, 372, 428, Dotted-legged plant bug 1311. 711, 1512, 1690, 1721, 2238, Double cocoons. The shedding of trachese, 1715. Dragon-flies, 1707. Diplotaxis not a vesicant, 1796. sordide, 1926. Dragon-fly. Large, 759. Diptera, 1329, 1736, 2267. Drasteria erechtea, 631, 1459, 2343. Parasitic, 2076. Drasterius amabilis, 1798, 1905. Preparation of, 2041. dorsalis, 2418. Swarming of, 2128, 2174. Drepanosiphum, 1678. Willow-galls made by, 46. Dried leaves as food for lepidopterous larvæ, 2159. Dipterous enemies of the honey-bee, 704. Driving nails into fruit-trees, 87. Phylloxera vastatrix, 2150. potato-beetles, 1044. galls of the willow, 46. Drones to death? Do worker bees sting, 1145. larva on swallow, 153. Drop of gold, 775. worm, 1059, 1090, 1650. larvæ in human body, 2256. Remarkable, 40. Drosera, 1499, 1688. Discontinuance of publication [of Am. Ent.], 2106. Drosophila amœna, 2119. Disease. Infesting Phylloxera with fungus, 1808. ampelophila, 2119. Diseases of beneficial insects. Fungus, 1813. nigricornis, 1607. chinch-bug, 2143. вр., 1607. Disippus butterfly, 1301. Drought. Connection of locust invasion with the Pupa of, 1193. occurrence of, 1422. on Hessian-fly. Effects of, 2031. Microgaster, 1301. Insects and, 2100. Disonycha alternata, 45. flaviventris, 1506. Drug-store pests, 510, 551. Dissosteria carolina, 667. Dryobius sexfasciatus, 583. Dryocampa, 1352. Ditching for young locusts, 1488. Dobson, 780. bicolor = Sphingicampa bicolor. Dock. Galls on supposed, 1165. pellucida = D. virginienais. DOCK. INSECTS AFFECTING. rubicunda, 244, 841, 915, 1329, 1352, Gastroidea cyanea, 1165. 1357, 2490, Gastrophysa cyanea = Gastroidea cyanea. senatoria, 151, 783. Doctoring fruit-trees again, 230. stigma, 488. Doctors differ, 98. virginiensis, 488. Dog-day harvest-fly, 1546. Dutchman's pipe. Papilio philenor on, 1321, 1774. Dolerus arvensis, 358. Worms on, 1321. unicolor, 1989. Dynastes tityrus, 580, 1216, 1292. Dollars. That hundred and fifty million, 1647. Dysderous suturellus, 2343. Domesticated katydids, 1536. Dytiscid in stomach of black bass, 1792. Dytiscide. Dimorphic, 28.

Dytiscus. Mode of feeding of the larva of, 2065. Dominican case-bearer, 1363. Dorthesia celastri, 1988. characias, 1686, 1730. Eacles imperialis, 1087, 1268. Covering of egg-puncture mistaken for, Earthquake commotion and locust multiplication and migration, Swinton, A. H., 2267. viburni, 1988. Eastern States. Supposed army-worm in New Doryphora, 1639. York and, 1990. Climate for, 1412. Eburia 4-geminata, 1014. 10-lineata, 21, 32, 48, 50, 88, 94, 119, 171, Eccopsis ferrugineana, 1968. 275, 276, 293, 318, 334, 347, Hickory, 2363. 365, 401, 411, 565, 650, 708, monetiferana, 1968. 709, 760, 866, 905, 908, 919, permundana, 2324. 925, 946, 968, 977, 983, 991, Variegated, 2363. Echthrus annulic rnis n. sp., 385. 995, 1019, 1027, 1044, 1056. 1059, 1068, 1096, 1101, 1218, Economic entomology, 1357, 1362. in California. Interest in, 1301, 1311, 1325, 1329, 1363, 1364, 1379, 1391, 1405, 1407, 2053.

Economic entomology in England, 1983.

ELM. INSECTS AFFECTING-Continued.

Dryobius sexfasciatus, 563. Fostering the study of, Galeruca calmariensis = G. xanthomelæ 2142. xanthomelæna, 1250, 1530, 1931, 2188, Importance of, 2238. Iowa, 2197. 2232, 2304, 2325, 2878, 2894. Recent advances in, 2281. Monocesta coryli, 1721. investigations in the South and West, Pemphiginæ, 1653. 1864. Physocnemum brevilineum, 696. notes, 2220, 2248. Saperda lateralis, 588. Ecpantheria scribonaria, 1140, 1153, 1311, 1802. Schizoneura rileyi, 1050. Ectobia germanica, 573, 1717, 2276. Tremex columba, 928. Edema albifrons, 423, 913. Elm-leaf beetle, 2188. Edinburgh International Forestry Exhibit. Re-Great, 1721. port of, 2342. Imported, 2232, 2304, 2325, 2378, 2394. Edwards' butterflies of North America (Review), 395, 479, 623, 827. Notes on the imported, 1931. Egg-feeding mites, 1637. and pear-tree borer, 928. parasite. Anthomyia, 1570, 1625, 1648. tree borer, 583, 696. parasites of Acridide, 2355. depredatore, 2414. Efficacy of chalcid, 2115. louse. Woolly, 1059. plant. Enemies of, 2096, 2215. saw-fly, 699. EGG-PLANT. INSECTS AFFECTING. Emmon's New York report, 84. Cassida texana, 2096, 2215, 2235. Emperor. Eyed, 1363. Doryphora 10-lineata, 21. Tawny, 1363. juneta, 2096, 2215, 2235. Emphytus maculatus = Harpiphorus maculatus. Egg puncture mistaken for Dorthesia. Covering Empos albicans n. sp., 22. Empoasca n. g., 22. punctures in raspberry and grape-vines, 2195. abnormis, 22. sack of some unknown spider, 1144. consobrina n. sp., 22. slits made by Homopters, 79. malefica = E. viridescens. Eggs in or on canes or twigs, 1829. maligna = E. obtusa. Experiments with locust, 1572. obtusa n. sp., 22. growth of insect, 2241. viridescens n. sp., 22, 30, 79. hatch? Will unimpregnated, 1029. Empretia stimulea, 60, 161, 424, 811, 829, 837, 1002. in maple twigs. Rows of, 665. Emulsions. Kerosene, 2391. Mass of, 65, 240. of petroleum as insecticides, 2134. of moth, 611. and their value as inbut para-itic cocoons. Not, 1290. socticides, 2126, 2200. in peach twigs, 554. Enchenopa binotata, 725, 1913, 1988. pear twigs. Rows of, 577. Enchophyllum binotatum = Enchenopa binotata. on sugar-maple, 350. Endrosa quercus — Lachnosterna quercua. in sumach, 18. England. Doryphora 10-lineata in, 2081. Egypt. Bug-hunter in, 372. Economic entomology in, 1983. Eight-spotted forester, 343, 1059, 1127, 1363. Hessian fly in, 2395. Eighth Report State Entomologist of Missouri, Introduction of the Hessian-fly into, 1482. 2398. Emige unserer schädlicherer Insekten, 1325. English currents. Beetles in, 1260. Elachistus euplectri, 2343. sparrow. Anent the, 1667. Elaphidion parallelum = E. villosum. Ennomos alniaria, 75. putator = E. villosum. magnaria = E. alniaria. tectum, 1602. subsignaria = Eudalimia subsignaria. villosum, 288, 793, 839, 1556, 2267. Entilia concava = Publilia concava. Elateridse, 3, 224, 430, 740, 846, 892, 932, 1030, 1504, sinuata, 163. Entomography of Hirmoneura, 2275. 1522, 1866, 2238. Entomological, 371, 869, 893, 1411. Monograph of, 1828. breakfast, 2311. Elder. Attacus cecropia on, 1533. cabinet. Walsh, 1107. Cecropia worm on, 1533. Elephantiasis or Filaria disease, 2185. circular. A recent British, 2509. Ellopia ribearia = Eufitchia ribearia. collection, 1328. Elm. Enemies of the, 1850. commission. Appropriation for, 1843. enemy, 1520. Bulletin No. 1, 1577. gall. Cockscomb, 1384. 2, 1578, ELM. INSECTS AFFECTING. 3, 1736. Cimbex americana, 699. 6, 2026. laportei = C. americana. Circular No. 1, 1575. 2, 15**76**. Colopha ulmicola, 1384.

Entomoligists in Europe. Number of, 1865. Entomological commission. Report. First, 1643. Second, 1959. Self-taught, 276. Address on, 1454. Third, 2267. Entomology. Fourth, 2343. all a humbug, 63. Reports of the U.S. Best works on, 418. Books useful for the study of, 418. 1757. in Buffalo, N. Y., 2035. correction, 1328. ignorance in the North, 431. California. Interest of economic, 2053. South, 390. Darwin's work in, 2129. information, 1351. Descriptive, 1199. notes, 2, 1396, 1554, 1609, 1610, 1681, 1958, 2015, 2058, 2066, 2135, Discussion on, 1391. 2147, 2168, 2184, 2192, 2219, Economic, 1362. 2228, 2247, 2261. in England. Economic, 1983. of the year, 2235. Fostering the study of economic, papers read at A. A. A. S., 1893. 2142. periodicals. New, 2056. General truths in applied, 2291, 2292. quackery, 439. Importance of economic, 2238. queries. A bundle of, 556. indeed run mad, 227, 1224. question, 1468. in Iowa. Economic, 2197. reports. Index to Ontario, 2228. Lecture on, 1894. Missouri, 1680. at Minneapolis, 2212. Society. American, 621. in Missouri, 1361. Address of president be-New York, 2244, 2254. fore Washington, 2835. Permanent subsection of, 2027. specimens. Care of, 2200. and the phonograph, 2481. speculations of the N. E. school of Professor Riley to Dr. Schaffer, 2360. naturalists, 44. in reality. Practical, 914. Professor Dana and Recent advances in economic, 2281. his, 82. horticultural, work at the Department of Agri-2230. culture, 1892 its relation to agriculture and its works, 831. advancement, 1329. wanted, 1529. Report Illinois State Horticultural Entomologist. American, 1309. Society on, 1056. Salutatory, 386. Missouri, 1113, 1127. Canadian (Review), 395, 479, 827. Study of, 2345. caught napping, 1110. Work in, 202. of Illinois. State, 327. Entomophthora, 1813. for Minnesota. A State, 1108. calopteni, 2240. of Missouri. First report of State, Epeira, 1847. 1059. riparia = Argiope riparia. Second, 1127. sp., 1144. Third, 1301. spinea, 813. Fourth, 131L Ephemera flaveola n. sp., 24. Fifth, 1329. flies. A hard story, 1043. Sixth, 1363. туера п. гр., 39. Seventh, 1428. Ephemerella n. g., 24. Eighth, 1482. consimilis n. sp., 24. Ninth, 1570. excrucians n. sp., 24. Index to reports of Ephemeridæ, 283, 525, 1043, 1205. State, 2026. Ephemerina, 2267. for New York. State, 1863. Ephemerinous genus Bætisca. Pupa of, 48. New York without an, 1330. Ephemeron, 283. One day's journal of a State, 383. Ephostia interpunctella, 325, 1293, 1896. for the Pacific coast, 2022. seas = E. interpunctella. Report U.S., 1878, 1721. Ephialtes gigas n. sp., 385. 1881-'82, 2119. pusio n. sp., 385. 1883, 2232, pygmæue n. sp., 385. 1884, 2291. Ephydra californica, 2203. 1885, 2363. gracilis, 2203. 1886, 2394. hians, 2203. 1887, 2418. Epicærus imbricatus, 35, 296, 1301, 1868, 2291. in the South. Field for the, 1109.

Entomologists at Boston, 1894.

Death of noted, 1285.

Epicanta, 1600, 1643, 1651, 2238, 2267.

atrata = E. pennsylvanica.

caneres, 124, 185, 401, 1056, 1558, 1600.

Epicauta corvina, 1206. Larval characters and habits of, 1600. habits, 1387. pennsylvanica, 38, 134, 185, 284, 362, 401, 732, 1056, 1558, 1600. puncticollia, 1381. vittata, 38, 134, 185, 401, 470, 794, 1066, 1230, 1558, 1600, 1860, 1969. Epidemic disease of Caloptenus differentialis, Epilachna borealis, 125, 417, 1289. corrupta, 2135, 2235. Epimeris. Grape-vine, 1301, 1363. Epipocus punctatus, 2406. Epitrix brevis = Crepidodera brevis. cucumeris = Crepidodera cucumeris. Distribution of species of, 1782. hirtipennis = Crepidodera parvula. Epizeuxis. Spruce, 2363. Erax, 1127. bastardi, 1127, 1269. Eriocampa cerasi, 1031, 1222, 1253, 1382, 2243. Eriosoma cornicola n. sp., 27 = Schizoneura cornicola. fungicola n. sp., 27 = Schizoneura fungicola. lanigera = Schizoneura lanigera. pyri = Schizoneura lanigera. tessellata = Pemphigus tessellata. ulmi n. sp., 1050 = Schizoneura ulmi. Eristalis, 986. Errors corrected. A few, 977. Erynnia alcee boring in stems of Malva sylvestris, 1602. Erythroneura, 164. australis n. sp., 22. octonotata n. sp., 22. tricincta = Typhlocyba tricincta. vitifex = Typhlocyba vitis. vitia = Typhlocyba vitis. siczac n. sp., 22. Eucalyptus. Galls on, 1965. Euchætes egle, 413. Euclemensia bassettella, 1972. Eudalimia subsignaria, 75. Eudamus tityrus, 787, 857. Eudemis botrana, 557, 585, 792, 1059, 1282. Eudioptis nitidalis, 808, 843, 1127. Eudryas grate, 1127, 1363. unio, 831, 1127, 1363. Eufitchia ribearia, 176, 364, 772, 877, 1068, 1224, 1570. Eumenes fraterna, 543, 1357, 1847. Eumenia atala, 2410. Eumenidæ, 770. Eunomia eagrus, 2242. Euonymus latifolia! Chionaspis enonymii on, 2403. Scale on, 2403. Euparia castanea, 2105. Eupelmus, 384, 1982, 2343. allynii, 2060, 2063, 2332. mirabilis, 384, 563, 1363, vs. Antigaster, 1978. Euphanessa mendica, 2340. Euphoria hirtipes, 2105. inda, 447, 1655.

melancholica, 372, 522, 842.

Euplectrus comstockii, 2848. platyhypenæ, 2343. Eupsalis minuta, 1363. Europe and America. Problem of the hop-plant louse in, 2400. The Colorado potato-beetle in, 1597. Grain Bruchus just imported from, 1120. Hagen, H. A. The Hessian-fly not imported from, 2267. Number of entomologists in, 1865. European natural enemies of the asparagus beetle. Pew, 2458. parasites. Importing, 208. Eurycreon rantalis, 2322, 2331, 2363. Euryomia melancholica = Euphoria melancholica. Euryptychia saligneana, 180, 1127, 2356. Eurytoma abnormicornis n. sp., 384. auriceps n. sp., 384. bicolor n. sp., 384. bolteri n. sp., 1059 = var. of E. diastrophi. diastrophi n. sp., 384, 1131. gigantea n. sp., 384. globulicola n. sp., 384 = var. of prunicola prunicida n. sp., 384. punctiventris n. sp., 384. seminatrix n. ep., 384 = var. of auriceps.sp., 62, 563. Eurytomid from stomach of Ohio shad, 1853. Eurytomides, 384, 1223. Euschistus fissilis, 2363. Euthyrhynchus floridanus, 2472. Euura orbitalia, 821. perturbans n. sp., 197. s.-gemms n. sp., 197, 821 = E. orbitalis. s.-nodus n. sp., 197. s.-ovum n. sp., 197, 821. Evagoras viridis = Diplodus luridus. Evarthrus orbatus, 872. Evergreen cleora, 2363. and other forest trees in New England and New York. Packard, A. S. Causes of destruction of, 2232, 2291, 2363, 2364, 2382, plant-lice, 1039. Evergreens. Borers in, 1324. Insects affecting, 271, 527, 1039, 1334, 2232, 2391, 2363. Exartema ferruginearum n. sp., 1968 = Eccopsis ferruginearum. montiferarum n. sp., 1968 = Eccopsis montiferarum. Agricultural editorial, 1849. Excursion. to Rocky Mountains, 988. Exenteron ornatus n. p., 385 =Cteniscus ornatur Exetastes illinoiensis, 385. suaveolens n. sp., 385. Exochiscus n.g., 385 = Orthocentrus. pusillus n. sp. = Orthocentrus pusillus. Exochus albiceps n. sp., 385. annulicrus n. sp., 385. atriceps n. sp., 385. Exorista cecropiso n. ep., 1112. doryphoree, 1059, 1431, 1450, 1672, flavicauda n. sp., 1127, 1442, 1670.

Exorista leucanise — Nemorsea leucanise. militaris = Nemoræa leucaniæ. osten-sackeni = Nemoræa leucaniæ. Exothecus prodoxi n. sp., 1831. Expected advent of the locust, 2307. Extensile penetrating organ in a gamasid mite, 1626. Exterminator. Fruit-pest, 2372. Extinguisher. Treat's insect. 531. Extreme cold on the Curculio. Influence of, 1335. Exuviation in flight, 1901. Exyra ridingsii, 1411. semicroces, 1385, 1390. Eyed emperor, 1363. Eyes and luminosity of fire-flies. Development of. 1840. None so blind as those who shut their, 275. Fagopyrum. Aerididæ that eat, 1645. Failure of tea-roses, 1708. Fall army-worm, 1256, 1267, 1282, 1301, 1482. canker-worm, 1423, 1482, 2238. web-worm, 1301, 1733, 2238, 2378, 2394. ou hickory, 460. Tent caterpillars and, 819. wheat, Webster, F. M. Insects affecting, 2291, 2363, 2394. Falling water. Moths attracted to, 2108. False caterpillars on the pine, 985. Scotch and Austrian pines, 1011. chinch-bug, 1829, 1423, 2291, 2418. indigo gall-moth, 1127. Walshia amorphella on, 1127. Fatherless and motherless race. That, 1650. race. Some further facts regarding that, 1658. Fatua denudata, 1733. Fear. A groundless, 284. Feathers. Attagenus injuring, 56. Felting caused by a beetle. Remarkable, 2113. Feniseca tarquinius, 2359, 2361. Food-habits of, 2407. Notes on, 2369. Plant-feeding habit of, 2361. Fertilization of Yucca, 1329, 1836, 2171. Fertilizers of alpine flowers, 1838. Fertilizing plants, 324. Few errors corrected, 977. Fidia. Grape-vine, 1231. longipes, 339. ap., 102. viticida n. sp., 272, 339, 1059, 1231. Field oricket, 2291. crops. Insects injurious to, 2238. for the entomologist in the South, 1109. Fiery ground-beetle, 1059. Fifteen-spotted lady-bird, 959, 1311, Fifth report State Entomologist of Missouri, 1329. Fifty million dollars. That hundred and, 1647. Fig insects, 2194.

Fighting the Curculio, 237.

Hessian-fly, 1587. Figurer's Insect World (review), 408.

Filaria disease. Elephantida's or, 2185.

sanguinis-hominis, 2185.

Filbert grape. Grape-vine, 1329. Fir Paraphia, 2363. Fire blight, 22, 30, 149, 285. cure for potato-beetles, 121. flica, 396, 1705. Development of eyes and luminosity of. Intermittence of phosphorescance, 1805. worm, 2291. First report noxious insects of the State of Illinois, 373. State entomologist of Missouri, 1059. U. S. Entomological Commission, 1643. volume. Close of the, 701. Fischer on naphthaline as an insecticide. Review of, 2274. Fish-fly. Large, 712, 903. Fitch. Dr. Asa, 1781. Flat-headed apple-tree borer, 1059, 2238, 2267. in horse-chestnut, 1316. twig borer, 555, 1423. borer in soft maples, 1250. cherry-borer, 2267. peach-borer, 2267. spruce-borer, 2267. Flavescent clover beetle, 2394. Flea-beetle, 1583. Colorado cabbage, 2291. Cucumber, 1558. Grape-vine, 298, 1252, 1301. Larvæ of grape-vine, 1041, 1074. Striped, 2291. Wavy-striped, 348, 564, 2291. in young tobacco-plants, 1782 Zimmerman's, 2291. beetles, 636, 2418. and Curculio, 873. Means against, 133. Satisfactory remedy for, 2236. like negro-bug, 1127, 1423, 2291, 2418. Fleas feeding on lepidopterous larvæ, 2110. Flesh-flies, 1643, 1736. fly. Common, 1423, 1570, 1625, 1643, Flesh-worms, 459. Flies in rooms. Swarms of minute, 506. Sucking organs of, 2182. Flight. Exuviation in, 1961. in insects is directed. How, 1891. Flights east of the Mississippi. Locust, 1549, 1613. Locust, 1590. Floating apiaries, 1785. Floods. One effect of the Mississippi, 2080. Florida, Ashmead, W. H. Report on insects injurious to garden crops in, 2388. Voyle, J. Effects of cold on the scale ininsects of the orange in, 2277. Flour. Poisonous, 586. and rye. Worms in, 72. Flouring-mills. Beetles in, 1261. Flower-bug. Insidious, 1127, 1423, 2418. Flower-garden. Insects in, 936. Flowering ash. Cocoons on, 950.

Flowers. Fertilizers of alpine, 1838.

Fluted-scale, 2389, 2418.

Recent California work against the,

Flying bug, 157.

locusta in Illinois, 1437, 1443.

Foe. Confounding friend with, 1378.

to corn. A new, 337.

cottonwood, 1834. green corn. A new insect, 1665.

Scavenger mistaken for a, 416.

Cherry plant-lice and their, 1251. Some friends and, 1274.

Folsom, Cal., Koebele, A. Notes on locusts about,

Food of Calopteron and Photinus, 2358.

habite of Feniseca tarquinius, 2407.

ground-beetles, 1769. longicorn beetles, 1902. Megilla maculata, 2145. thrushes, 1726.

of insectivorous plants, 1499.

for lepidopterous larvæ. Dried leaves as, 2159.

man. Locusts as, 1481.

Number of molts and length of larval life as influenced by, 2167.

plants for the cotton-worm. Possible, 2064, 2119, 2155.

of Samia cynthia, 2204.

relations of Carabide and Coccinellide, 2151. Salt-water insects used as, 2203.

for silk-worms, 1819.

trout, 1142.

Forage crops. Insects injurious to, 2238. Forbes, S. A. Experiments on chinch-bugs, 2165. Forbes' investigations on the food of fresh-water fishes. Professor, 2515.

Forest insects. Larvæ of injurious, 2267.

Tent caterpillar of the, 645, 688, 1181, 1200, 1301, 1331.

tree insects, 2418.

Packard, A. S. Notes on. 2253

trees in northern New England and New York, Packard, A. S. Causes of the destruction of evergreen and other, 2232, 2291, 2363, 2364, 2382. and weeds of Germany. Ratzeburg. Review, 1110.

Forester. Eight-spotted, 343, 1059, 1127, 1363. Forestry exhibition. Report on the Edinburgh international, 2342.

Forficula, 1798, 2267.

Formica aphidicola n. sp., 27.

fusca, 2105, 2226.

latipes n. sp., 27 =Lasius latipes.

rufa, 2105, 2114,

Formicidæ, 438, 1278, 1594, 1987.

Fossil tineids, 2084.

Four-humped Curculio, 1088.

Apple-snout beetle or, 1088.

lined leaf-bug on current, 715.

Fourth report State entomologist of Missouri, 1311.

Fourth report U. S. Entomological Commission, 2343.

Fowls. Internal mite on, 2157.

Parasites on, 1308.

France affected by Phylloxera. Half the vine area of, 2020.

Grape Phylloxera in, 2193.

Fraternal potter-wasp, 1127.

Friend. Another insect, 879.

with foe. Confounding, 1378. unmasked, 374.

Friendly criticism, 1307.

notes, 1806.

Friends and foes. Some, 1274.

Inquiring, 1642, 1656.

and insect foes. Insect, 28.

Frog-spittle insects, 691.

Frogs, 217.

Frosted lightning hopper, 1329.

Fruit culture in the South, 2265.

foe. A new, 35.

and fruit trees. Insects injurious to, 2238. Trimble's enemies of, 187.

growers. Insects of interest to, 367.

in northern Illinois. Six worst enemies of, 377.

insects in California, 2245. invigorator again. Best's, 530. Jarring down infested, 1318.

pest exterminators, 2372.

Snout-beetles injurious to, 1302. trees again. Doctoring, 230.

Driving nails into, 87.

Icerya purchasi, an insect injurious to, 2401.

FRUIT TREES. INSECTS AFFECTING.

Epicerus imbricatus, 35. Ithycerus noveboracensis, 16.

Lytta sence = Pomphopœa sence.

Oncideres cingulata 1 442, 443. Pomphopæa ænea, 1056.

Fruit trees. Insects injurious to, 16, 31.

Tarred paper for, 1568.

worm. Cranberry, 2291. Tomato, 1136.

Fruits and grains, Webster, F. M. Experiments on the effect of punctures of Hemipters en shrubs, 2382.

Fuchsia beetles, 133.

Fuchsias. Beetles injuring cabbages and, 1868. FUCHSIAS. INSECTS AFFECTING.

Haltica carinata, 183, 1868.

exapta = H. carinata.

Fulgoridæ, 787.

Fuller. Retirement of Mr., 1910.

Fuller's rose-beetle, 1708, 1721, 2291.

in California, 1749. Habits of, 1708. Not, 1825.

Fumago salicina, 2138.

Fumigation against orange scales, 2520.

Fungoid growths, 500.

Fungus in Cicada, 1809.

diseases of beneficial insects, 1813. Infecting Phylloxera with, 1863. Fungus foes, 1937. growths to destroy insects. Use of, 1808. Remarkable parasitic, 1313... White-grub, 594. Fuzzy galls on blackberry twigs, 609. Gad-fly. Sheep, 887. Galerita janus, 1824. lecontei, 1824. Galeruca calmariensis - G. xanthomelæna. rufosanguinea - Adimonia rufosan. guines. xanthomelæna, 1185, 1850, 1520, 1931, 2188, 2282, 2304, 2825, 2878, 2879, 2414. Gall on acorn-cups. A new oak, 1806. blackberry, 243. and raspberry canes. Gouty, chestnut, 858. Coccid mistaken for a, 1972. Cocksoomb elm, 1384. flies, 204. gnat, Loew. Description of the rye, 2267. Wagner. Observations on the new crop, **22**67.. on grape-vine. Large compound, 720. Grape-vine apple, 1329, 1898. leaf, 734, 840, 1716. insects, 1559. Galls and, 1967. Leafy oak, 774 literature. Bibliography of, 2059. louse. Hickory-stem, 685. making genus of Apionine, 2231. moths, 1127, 2356. Pemphiginse. Biological notes on, tortricid. A pretty and unique, 2176. Mossy-rose, 570, 1166. nute, 1560. Oak-fig, 745. leaf, 799. on oak twigs. Cynipid, 1822. Pelargonium, 1764. Pithy blackberry, 1131. Pod-like willow, 1170. Prickly ross, 1194, 1245. Raspberry gouty, 1124. root, 1149. Rose, 1235. on Solidago leaves, 1924. spotted touch-me-not, 852. Sugar-maple mite, 1265. Trumpet grape, 791, 1116.

Galleria cereana, 581, 716, 904, 1069, 1298, 1257, 1508, 1728. Gallmücke. Beschreibung einer den Birnen schädlichen, 2392. Galls, 137, 1561. on blackberry, 263. cottonwood leaf, 718. Unsightly, 446. Cypress, 1180. on Eucalyptus, 1965, and gall insects, 1967. Golden-rod, 798. Grape, 111, 160.

Galls on the grape-vine. Leaf, 724, 840, 1716. growing on wild sage, 1347. Hackberry psyllid, 220d. Jumping seeds and, 1496, 2163, 2173. on leaves of soft maple, 445. wild grape-vine. Conical, 1077. made by moths, 1127. Phylloxera. New hickory, 1901. and mining in apple-twigs. Small, 552. Oak and rose, 1037. Production of, 1056. on sugar-berry, 762. supposed dock, 1165. and their architects, 518, 821. on white oak. Woolly, 739. Gamasid mite. An extensile penetrating organ in a, 1626. Gamasus juloides infesting Iulus marginatus, DAR. Garden crops. Experiments with insecticides upon insects affecting, 2344. in Florida, Ashmead, W. H. Report on insects injurious to, 2388. insects, Alwood, W. B. Tests with insecticides on, 2888. Insects in the flower, 936. vegetables. Insects injurious to, 2238. web-worm, 2363. Gardens. Ants' nests in, 496. Destroying black ants in, 438 Toads in, 822. Gargaphia amorphæ, 45. tiliæ, 45, 400. Gas treatment for scale insects. Coquillett, D. W., 2418. waste vs. Curculio, 549. Gases against scale insects. Use of, 2389. Gasteracantha cancer, 2111. Gastroidea cyanea, 316, 1165. Gastropacha americana, 812. velleda = Tolype velleda. Gastrophilus equi, 1391, 1563, 2238. Gastrophysa cyanea = Gastroidea cyanea. Geelong, Victoria. Grape Phylloxera at, 1732. Gelechia abietisella, 2232. cerealella, 2291. gallæ-solidaginis n. sp., 1059, 1165, 2213. glandulella n. sp., 1306. Hemlock, 2232. White heart hickory, 2363. Gemminger and Harold's Catalogus Colcopterorum, 1866. General index and supplement to the Missour reports, 2026. truths in applied entomology, 2291, 2292. Genuine army-worm in the West, 2009. vs. bogus chinch-bugs, 1402. Genus grinding, 2120. Geographical distribution of the Rocky Mountain locust, 2267. range of species, 1614.

Geometer. Chickweed, 1059, 1154.

Knot-weed, 1059.

Geometra catenaria = Zerene catenaria.

Geometridæ. Larval characters_165.

Georgia. Rascal leaf-crumpler in, 1962.

GOOSEBERRY. INSECTS AFFECTING.

German asters. Diabrotica vittata on, 148. Dakruma convolutella, 1059. thoroughness, 1597. Ellopia ribearia = Eufitchia ribearia. Gigantic rhinoceros beetle, 580, 1216, 1292. Epicærus imbricatus, 85. water-bug, 534, 733. Eufitchia ribearia, 772, 1068, 1570. Gilly-flowers. Cabbage-worms upon, 614. Nematus ribesii, 140, 228, 333, 756, 772. Plutella cruciferarum injuring, ventricosus = N. ribesii. Pristiphora grossulariæ, 40, 189, 772. Gilt gold-beetle, 790. Tentbredinid, 136. Girdled pear-twigs, 848. Gooseberry saw-fly, 140. Girdler. Twig. 476, 1938. Imported, 228, 333, Girdling habits of Pædisca obfuscata, 2405. span-worms, 1068, 1570. Glassy cut-worm, 1059, 2291, 2394. worm, 189. Mesochorus, 1127. worms. Current and, 1570. winged soldier-bug, 1301. Imported, 756. Glaucopid caterpillar. Protective device em-Gordiacea, 612. Gordius, 861. ployed by a, 2242. Glorified squash-bug, 262. aquaticus, 447, 1148. Glossina morsitans, 1467. varius, 1148. Glover, Townend. Obituary, 2218, 2258. Gortyna nebris, 1646. Glow-worm, 1917. nitela, 331, 401, 632, 694, 734, 754, 820, 862, 940, 976, 993, 1009, 1056, 1059, 1210, 1558, Nature of the phosphorescence of, 1810. 1589, 1695, 1646, 1870, 2229. That, 1097. sp., 138. Glyphe viridascens n. sp., 12. Gouty gall on blackberry and raspberry canes, Glypta alboscutellaris n. sp., 385. 1771. diversipes n. sp., 385. Governor of Kansas on the grasshopper question. ruficornis n. sp., 385. Report to, 1573. rufipleuralis n. ep., 385. Governors of Western States on the Rocky Mount-Gnats, 625. ain locust, 1557. Goat-weed butterfly, 1117, 1127, 1829. Gracilaria. Purple willow, 2363. Paphia glycerium feeding on, 1329. Grain Aphis, 2394. Going it blind, 588. Bad work of, 1638. Gold-beetle. Gilt, 790. vs. rust, 1806. Drop of, 775. Bruchus, 1301. Golden-rod galls, 798. of Europe just imported, 1120. GOLDEN-ROD. INSECTS AFFECTING. Chinch-bug not in seed, 888. Cecidomyia carbonifera, 1924. GRAIN. INSECTS AFFECTING. solidaginis, 798. Aphid, 99. Euryptychia saligineana — Pædisca scudderi-Asopia farinalis, 2337. Bruchus granarius, 1301. Gelechia gallæ-solidaginis, 1059, 1165. Gelechia cerealella, 2291. Graphtholitha olivaceana, 2285. Nectarophora granaria, 126. Nectarophora rudbeckiæ, 2205. Grain moth, 2337. Pædisca scudderiana, 180. Angoumois, 2291. Siphonophora rudbeckiæ=: Nectarophora rud-A new leaf-hopper injurious to small, 1767. beckiss. plant louse, 99, 126, 1127. Trypeta solidaginia, 180, 798, 1059. Silvanus, 1259. Golden-rod tortoise-beetle, 1127. Sphenophorus, 2363, 2394. Gomphocerus shastanus n. sp., 1959. weevil, 15, 2291. Gomphus amnicola n.sp., 24. weevils. Aniseed vs., 1742. consobrinus n. sp., 39. Grains. Calandra granaria in, 1250. fluvialis n. sp., 24. and grasses, Webster, F. M. Insects afgrasilinellus n. sp., 24. fecting, 2394. quadricolor n. sp., 39. Webster, F. M. Experiments on the effect vastus n. sp., 24 of punctures of Hemipters on shrubs, ventricosus n. sp., 39. fruite, and, 2382. Gonatopus contortulus, 2116. Granddaddy long-legs, 838. pedestris, 2116. Granulated cut-worm, 2291. pilosus. Biology of, 2116. Grape-bark louse, 106. Goniocotes hologaster, 1094. berry moth, 557, 585, 792. Good words. More, 532. cane-gall Curculio, 1059. Gooseberries and currants. Green worms on, 136. canes and apple-twigs. Eggs in, 1223. Gooseberry and current worms, 772. Egg on, 855. fruit-worm, 1059. punctured, 513, 1233.

Grape Curculio, 373, 1059. GRAPE-VINE. INSECTS AFFECTING-Continued. . disease, 1311. Amphicerus bicaudatus, 1185. Anomala lucicola, 113. fruit worm, 1050. gall. Trumpet, 791, 1116. Aphis vitis, 102, 170. galls, 111, 160. Apis mellifica, 441. insects, 164, 267, 1056, 1569. Blennocampa pygmæa, 1252. Bostrichus bicaudatus - Amphicerus bicaudaleaf-folder, 1301, 1579. folders eaten by spiders, 468. Hairy, 664. Capsus oblineatus = Lygus pratensis. gall-loure, 373, 1301, 1311. Cecidomyia vitis-coryloides, 1329. galls, 127, 1295, 1435, 1531. lituus, 791, 1116. Leaf-hoppers of the, 399. pomum, 403, 1284, 1329, 1898. leaf-louse, 916. viticola, 791, 1077, 1116, 1329. Phylloxera enemy, 1530. Cecidomyid, 111. Phylloxera, 1329, 1363, 1373, 1482, 1721 Ceramby cid, 429. in California, 1727. Ceresa bubalus, 1323, 1329. France, 2193. Chærocampa pampinatrix = A m pelophaga Geelong, Victoria, 1732. myron. New biological facts regard-Cœliodes inæqualis = Craponius inæqualis. ing, 1421. Colaspis flavids, 129, 133, 135, 231, 1301, 1311. Newest facts of, 1404, Corimelæna pulicaria, 1046. not at the Cape, 1841. Craponius inæqualis, 373, 1059. permanently destructive, Curculio, 267. Dactylosphæra vitifoliæ = Phylloxera vasta-Notes on the, 1952, 2291. natural history Darapsa myron = Ampelophaga myron. of, 143**9**. Desmia macplalis, 468, 1158, 1301, 1569, 1579. Use of guano for, 1837. Drosophila ampelophila, 2119. root-borer, 373, 1301. Eggs, 855. New, 1059, 1127. Enchenopa binotata, 725. Enchophyllum binotatum = Enchenopa binorot. Phylloxers and, 1623, 1628. saw-fly, 2291. scale insect, 1706. Erythroneura sp., 164. seed Curculio, 1050. tricincta = Typhlocyba tri-Insect infeeting, 482. cincta. maggot, 1127. vitis = Typhlocyba vitis. vine Aphia, 170. Eudemis botrana, 557, 585, 792, 1059. apple-gall, 1329, 1898. Eudryas grata, 1127, 1368. An apple growing on a, 403, 436. unio, 1127, 1363. Bark-lice on, 1212. Fidia longipes, 839. beetles, 113, 129, 133, 339, 354. sp., 102. viticida, 272, 339, 1059, 1231. borer, 429. caterpillars, 124. Galls, 128, 791. caterpillars. Black, 1018. Graptodera chalybea = Haltica chalybea. Haltica chalybea, 208, 298, 1041, 1252, 1255, 1301, Colaspis, 231, 1301. again, 1311. 1821, 1855, Conical galls on leaves of wild, 1077. Harrisiana americana, 164, 218, 1127, 1136. Epimeris, 1301, 1363. Isosoma vitis, 482, 484, 1059, 1127. Fidia, 272, 1050, 1231. Lasioptera vitis, 720, 1829. filbert-gall, 1329. Leaf-hopper, 484. flea-beetle, 298, 1252, 1301, 1821, 1855. Lecanium vitis = Pulvinaria vitis. Larvæ of, 1041, 1074. Lygus pratensis, 682. and fuchsia beetles, 133. Macrodactylus subspinosus, 373, 565, 1375. a hickory. Hybrid between, 1284. Membracid, 286. Membracis ampelopsidis, 1183. hoppers, 1024. Œcanthus niveus, 286, 414, 723, 1059, 1323, 1329, Injured strawberry and, 682. insects, 128, 725. 1333, 1691, 2195, (FRAPE-VINE. INSECTS AFFECTING. Oncometopia undata, 36, 79, 164, 399. Acoloithus falsarius, 124. Orocharis saltator, 513, 1328, 1329. Acronycta oblinita, 1208, 1301. Orthosoma brunneum, 397. Ægeria polistiformis = Sciapteron polistiforcylindricum = 0. brunneum. Oxyptilus periscelidactylus, 664, 1059, 1175, Alypia octomaculata, 1059, 1127, 1130, 1208, 1363. Amblycorypha oblongifolia, 1329. Pelidnota punctata, 113, 129, 354, 725, 1221, 1301 Ampeloglypter secostris, 1059. Pemphigus vitifoliæ = Phylloxera vastatrix.

Ampelophaga myron, 1086, 1127, 1247.

Penthina vitivorana = Eudemis botrana.

GRAPE-VINE. INSECTS APPECTING-Continued. Grapes spoiled by something, 464. Grapholitha gallæ-saliciana n. sp., 1968. Philampelus achemon, 1001, 1127. pandorus, 1102. Grapholitha ninana n. sp., 2176. satellistia = P. pandorus. olivaceana n. sp., 1968. Phylloxera vastatrix, 127, 160, 373, 565, 724, 840, Habita of, 2285. 916, 1281, 1295, 1801, pruinivora, 367, 373, 1795. 1811, 1825, 1829, 1842, Graphops pubescens, 2229. 1363, 1374, 1376, 1421, Grapta, 1301. 1423, 1435, 1439, 1482, comma, 461, 1140. 1530, 1531, 1623, 1628, interrogationia, 420, 444, 1306. 1716, 1721, 1727, 1732, Graptodera carinata = Haltica carinata. chalybea = Haltica chalybea. 1837, 1841, 1895, 1900, 1907, 1952. punctipennis - Haltica puncti pennis. vitifoliæ = P. vastatrix. Grass-bug and its habits, 882. Poeciloptera pruinosa, 1329. insects, 541. Polycaon confertus, 2103. GRASS. INSECTS AFFECTING. Prionus imbricornis, 1127, 1274. Laphygma frugiperda, 1127. laticollis, 561, 1059, 1081, 1127. Leucania unipuncta, 1877. Proconia undata = Oncometopia undata. Grasses, Webster, F. M. Insects affecting small Procris americana = Harrisiana americana. grains and, 2394. falsarius - Acoloithus falsarius. Grasshopper. The, 1588. Psychomorpha epimeris, 1301, 1363. Colorado, 502. oggs, 536, 667. Pterophorus periscelidactylus = Oxyptilus periscelidactylus. Trombidium preying on, 624. Pulvinaria vitia, 106, 1212, 1706, 1716. Governors of Western States on Pyrophila pyramidoides, 671, 1301. the Rocky Mountain, 1557. Sciapteron polistiformis, 373, 1801, 1509. Hateful, 873, 475. injuries. New method of coun-Selandria vitis = Blennocampa pygmæa. Sinoxylon basilare, 1311. teracting, 2235. Smilia auriculata, 1183. injury in the near future. Prob-Spilosoma virginica, 1202, 1301. abilities of, 2335. Tettigonia coagulata, 1024. Outlook for locust or, 246L vitis - Typhlocyba vitis. machine. A satisfactory, 1592. Thyreus abbotii, 763, 1018, 1127, 1248. A new enemy of the, 1541. Parasite on hateful, 728. Typhlocyba tricincta, 25. pest of the West. Important obvitis, 203, 686. Grape-vine. Insects injurious to the, 267, 1059, servations on, 1571. 1081, 1096, 1091, 1102, 1118, 1130, 1137, prospect, 1698. 1158, 1175, 1202, 1221, 1255, 1281, 1301, question. Report to governor of 1311, 1329, 1363. Kansas, 1573. Large compound gall on, 720. ravages in California, 2323. leaf-galls, 724, 840, 1716. Rocky Mountain, 1587. hoppers, 484, 686. Western, 1998. pest. More about the, 1376. year! Is this a, 1565. plume, 1059, 1801. Grasshoppers, 368, 391, 402, 483, 1368, Procris, 213. Destructive, 2363. root-borer, 1509. and locusts, 26, 147, 929. roots. Wood-lice on, 1906. Mites on, 146. tomato-gall, 1329. Rear-horses vs., 590. trumpet-gall, 1329. Young, 2377. worm. Dark, 763. Grass-worm, 2119. Green, 671. Gray straight-horned anout-beetle. Large, 1033. Pyramidal, 1301. Gray. Personal reminiscences of Dr. Asa, 2419. vines. Bugs on, 1046. Greasy cut-worm, 1059, 2291, 2418. Caterpillars on, 1208. Great discovery. Curculio extermination possi-On the cause of deterioration in ble, 1178. some of our native, 1342. elm-leaf beetle, 1721. Egg-puncture in raspberry and, Lebis, 1301. leopard moth, 1311. Eggs of the tree-cricket on, 723. Green apple-leaf tyer, 1311. mistaken for chinch-bug. Bugs corn. A new insect foe to, 1665. on, 398. grape-vine worm, 671. to Phylloxera in sandy soil. Rehag moth, 1233. sistance of, 2250. striped maple-worm, 1329. worms on gooseberries and currents, 136.

Greenhouse pests, 508.

plants, 508.

Rose-chafers on, 1375.

Grapes cut off by tree-cricket, 414.

Honey-bees eating, 441.

Gregarious worm on horse-chestnut, 1192. walnut caterpillars, 1045. willow-worms, 856. Ground-beetle. Another herbivorous, 2042. Fiery, 1050. Murky, 486. Pennsylvania, 1059. Subangular, 372. beetles, 1127, 1558, 1625, 1643. Eggs of, 692. Food-habits of, 1760. Vegetal feeding, 1788. Groundless fear, 284. Growth of insect eggs, 2241. Grub fungus. White, 594, 646, 1064, 1430, 1436, 1599, 1803, 1823. information wanted. White, 1072. White, 68, 410, 1020, 1050, 2238, 2363, 2394. worm, 54. Grubs and guess-work, 1440. Gryllidæ, 929, 2267. Gryllotalpa, 2267. borealis, 562, 1270, 1798. columbia, 562. longipennis = G. columbia. Gryllus, 2267, 2384. abbreviatus, 143, 433. Guano for grape Phylloxera. Use of, 1837. Guard. Be on the, 708. Guess-work. Grubs and, 1440. Guide to the study of insects. Review of Packard's, 395, 479, 623, 827. Gymnetis nitida = Allorhina nitida. Gyrinus larva in stomach of shad, 1853. Habit. Change of, 2096. Hackberry butterflies, 1356, 1363. HACKBERRY. INSECTS AFFECTING. Apatura alicia, 1363. celtis, 1356, 1363. clyton, 1356, 1363. here = A. clyton. lycaon = A. celtis.Pachypsylla c.-mamma, 2208. c.-venusta, 2208. Hackberry payllid galls, 2208. Hadena, 2238. chenopodii = Mamestra trifolii. devastatrix, 1056, 1059, 2291, 2394, 2355. juncta, 1056. renigera, 964, 1059. sp., 1047. subjuncta, 901, 1059. Hæmatopis grataria, 1059, 1154. Hag-moth. Green, 1233. larva, 777, 1272. Hagen, H. A. The Hessian-fly not imported from Europe, 2267. Observations on certain North

American Neuroptera, 39.

Hagen's mystery. Dr., 1943.

worms, 861, 1643, 1959.

grape-leaf folders, 664. Haldeman, S. S. Death of, 1909.

Hair snakes, 612, 861, 1143.

Hairy caterpillar, 558.

Halictus sp., 800. Halisidota antiphola n. sp., 40 = H. tessellata. сагую, 45. harrisii n. sp., 45 = H. tessellata. tessellata, 40, 45, 50. Haltica alternata - Disonycha alternata. carinata, 133, 1868. chalybes, 203, 298, 1041, 1074, 1252, 1255, 1301, 1583, 1821, 1855. cucumeris = Crepidodera cucumeria. exapta = H. carinata. helxines = Crepidodera helxines. punctipennis, 2383, 2468. striolata = Phyllotreta vittata. Haltichella perpulchra, 11, 12. Halticidse, 873, 1858, 2236. Ham-beetle. Red-legged, 1363. Hams. Skippers injuring smoked, 1734. Hand-maid moth, 2222. Hard story. Ephemera flies, 1043. Harlequin cabbage-bug, 1099, 1311, 1534, 2238, 2268, Harmless insects, 1526, Harold's Catalogus Coleopterorum. Gemminger and, 1866. Harpactor cinctus = Milyas cinctus. Harpalus, 1643. caliginosus, 3, 486. erraticus, 1537. pennsylvanicus, 372. Harpiphorus maculatus, 499, 955, 965, 1056, 1264, 1570, 1586, 2324, Harris's bark-louse, 873. correspondence (review), 623. insects injurious to vegetation (review), 568. Harrisina americana, 164, 213, 789, 1056, 1127, 1136. Harvest bugs, 412. fly. Dog-day, 1546. mites, 1326. Hatch pupe. How to, 509. Will unimpregnated eggs, 1029. Hatching! Are the locusts, 1566. Hateful grasshopper, 378, 475. Parasites on, 728. locust, 1452. locusts. . Prairie fires and, 1453. Hawk's pellets, 601, 643. Hawthorn. Worms on, 1051. Hay. Worms under mulch, 1161. Head-maggot, 450, 1059. of winged insects, Packard, A. S. Number of segments in the, 2267. Hedge-hog caterpillar, 1153. Helia americalis, 2414. Helianthus. Lixus macer bred from, 2404. Heliothinse. Synopsis of (review), 2178. Heliothis armigera, 192, 636, 820, 993, 1136, 1256, 1301, 1353, 1664, 1695, 1886, 1915, 2119, 2238, 2343. marginidens = Pyrrhia exprimens. phlogophagus, 936, 945, 993, 1056. umbrosus, 2343. Hellgramite, 1570, 1584. fly, 473, 1329. Helops æreus, 751.

Helops micans, 963.

Hexaplasta, 1749, 1982.

pullus = H. æreus. zigzag n. sp., 1749, 1932, 2343. Hemaria thysbe, 879. Hexapoda, 1955, 2267. Poisonous, 2399. Hemerobidæ, 2267. Hemileuca maia, 595, 722, 785, 760, 1329, 1352. Hibernating Aletia chrysalids. Supposed, 1927. Hemiptera, 400, 1329, 2267. apple-worms. Severe cold and, 2637. upon shrubs, fruits, and grains. Web-Hibernation of Aletia xylina in the United States ster, F. M., 2382. a settled fact, 2141. Experiments of the ef-Amphipyra pyramidoides, 1471. fects of punctures of, army-worm, 2086. 2382. the cotton-moth, 1728. Hemipterological studies, 2034. worm moth, 1953. insects, 1129. Hemispherical larva at bottom of ant hill, 1789. Hemiteles, 77, Hickory-bark borer, 938. borer, 269, 308, 1401. cressoni n. sp., 1059. fuscatus n. sp., 380 = var. of nemati-Citheronia regalis on shell-bark, 775. Eccopsis, 2363. Fall web-worm on, 460. nemativorus n. sp., 380. galls, 360. thyridopterygis n. sp., 1059. Hemlock Gelechia, 2232. made by Phylloxera. New, 1901. Gelechia. White heart, 2363. HENLOCK. INSECTS AFFECTING. Buprestid, 2267. Hybrid between a grape and a, 1284. Cerambycid, 2267. HICKORY. INSECTS AFFECTING. Arhopalus pictus = Cyllene pictus. Gelechia abietisella, 2232. Cecidomyia tubicola, 1232. Hen. Death of a, 1094. Citheronia regalia, 775. Henderson's experiments. Mr., 1688. Clytus pictus = Cyllene pictus. Henous, 1643. Cyllene pictus, 89, 269, 308, 1516. confertus, 1600. larval habits, 1387. Datana ministra, 2222. Eccopsis, 2363. Heptagenia n. g., 39. cruentata n. sp., 39. Galls, 791. maculipennis n. sp., 39. Gelechia, 2363. simplex n. sp., 39. Hyphantria cunea, 460. Herbivorous ground-beetle. Another, 2042. textor = H. cunes. Herpetogomphus rupinsulensis n. sp., 24. Phylloxera caryse avellana, 1901. Hesperid larva feeding on Canna, 1897. caulis, 360, 685, Hessian-fly, 150, 250, 426, 1581, 1665, 2008, 2267, 2365. globuli, 360. Effects of drought on, 2031. scissa, 1901. in England, 2395. Scolytus carys = S. 4-spinosus. into England. Introduction of, 2398. 4-spinosus, 938, 1329, 1401, 1754. Fighting the, 1587. Hickory Scolytus, 938. half way around the world, 2440. stem gall-louse, 685. Koeppen's account of the, 2267. vs. locust-borer, 1516. in North America. Early references Hipparchiscus n.g., 40 = Aplodes. to the, 2267. venustus n. sp., 40 = Aplodes mi-Not the, 1445. mosaria. imported from Europe. Hagen, Hippodamra. Ambiguous, 2119. H. A., 2267. convergens, 639, 1251, 1431, 1672. Parasites of, 2332. glacialis 849. prior to the Revolution. Insects conmaculata = Megilla maculata. founded with the, 2470. Hirmoneura. Entomography of, 2275. Probable parthenogenesis of the, 1787. Larval stages and habits of, 2169. in seed wheat, 494. obscura, 2169, 2275. Silesia. Cohn, F., 2267. Hirundo americana, 1502. Wheat-rust and the, 1605. fulva, 1502. Hetserina pseudamericana n. sp., 39. Hispa scutellaris = Odontota dorsalis. rupamnensis n. sp., 39. Hitherto unknown life-habits of two genera of rupinsulensis n. sp., 24. bee flies, 2002. scelerata n. sp., 39. Hockeria n. g., 11 = Haltichella.texana n. sp., 39. perpulchra n. sp., 11 = Haltichella per. Heteronychus relictus = Ligyrus relictus. pulchra. Heteropelma datanæ n. sp., 2526. Hog caterpillar of the vine, 1127. Heteroptera, 1739.

Defensive odors of the, 354.

bilineata, 372, 1043, 1851.

Hexagenia n. g., 39.

pulchra.

Hog caterpillar of the vine, 1127.

infested with parasites, 1247.

Hogs vs. bugs, 387.

Holcaspis mamma, 518.

Holes around the roots of young ash-trees in the Horse-hair snakes, 612. Horses. Lice on, 258. nursery, 471. Holocera glanduella n. sp., 1310, 1311. Horticultural entomology. Recent advances in, Homalomyia leidyi n. sp., 382. 2230. pruinivora n. sp., 382. Horticulture. Utilization of ants in, 2089, 2137. wilsoni n. sp., 382. Houghton's insects in the orchard, 188. Home. Colorado potato-beetle's native, 1462. House-fly, 864, 2078. Homoptera. Egg-alite made by, 79. Proboscis, 1783. Honey-ant. Peculiarities of the Mexican, 1417. How great wits jump together, 567. bee. Braula cœca not particularly injuto hatch pupse, 509. rious to, 1982. Howard, L. O. Chinch-bug, 2418, Bug preying on, 18. Codling-moth, 2418. Dipterous enemies of 704. Howell, M. A. Experience with the spring Discussion on, 1455. canker-worm, 2267. Insect enemies of, 1059. Hubbard, H. G. Miscellaneous notes on orange bees carnivorous! Are, 2098. insects, 2164. eating grapes, 441. Report on cotton insects, 2343. locust seed weevil, 1026. Rust of the orange, 2291. weevil, 1474. Scale insects of the orange, 2119. producing oak gall, 1942. Hudson Bay Lepidoptera, 1985. Too fond of, 764. Hull's Curculio catcher, 872, 651, 875. Honor. Deserved, 2124. Hulst's observations on Pronuba yuccasella, Mr., Hop Aphis, 1001, 2291, 2394, 2418. 2371. and the cranberry, Smith, J. B. Insects Human animal. Parasites of the, 497. affecting the, 2291. body. Larvæ in, 382. growing in the West, 235, 279. lungs. On a larva of Scenopinus from, insects, 1001. 1348. plant louse in Europe and America. Prob-Humbert on Lucilia, 2255. lem of the, 2400. Humblebees, 800. fully solved. Problem of the, Humbug. Another, 195. 2308 new Curculio, 589. Life-history of the, 2398. Entomology all a, 63. Problem of the, 2400. New, 122. vine caterpillars, 444, 979. Curculio, 680. Humming-bird moths caught by the tongue, 1388. HOP-VINE. INSECTS AFFECTING. Grapta interrogationis, 420, 444. Hundred and fifty million dollars. That, 1647. Hypena humuli = H. scabra. legged worms, 219. scabra, 979, 1001. Hybrid between a grape-vine and a hickory, 1284. Hyperchiria io, 420. Hydrachna belostomæ n. sp., 1632. Phorodon humuli, 235, 279, 1001, 2393, 2394, Hydrophilus piceus, 2016. 2396, 2400. triangularis, 2016. Saturnia io = Hyperchiria io. Notes on, 2016. Vanessa interrogationia — Grapta interroga-Hylecœtus lugubris, 575. tionis. Hylesinus opaculus, 1656, 1721. trifolii, 1690, 1721, 1777, 1846, 2512. Hop-vine. Insect foes of the, 420. Hoplophora arctata n. sp., 1363, 1370. Hylobius confusus, 700, 1168. Hopper in Iowa, 1585. stupidus, 845. Hylurgope, Packard, A.S. Development of, 2267. Horinus lævis = Merinus lævis. Hylurgus pinifex, 2267. Horizontal insect boxes, 1963. Hymenoptera, 543, 1329, 1736, 2267. Hormaphis spinosus, 1678. Description of North American, 385. Horn-bug, 1636. Parasitio, 310. Cocoon of, 784. Horned Passalus, 1311. Willow galls made by, 46. Workers among, 311. Egg of, 1329. Hymenorus obscurus, 2105. Hornia n. g., 1601, 1643. rufipes, 2105, 2226. minutipennis n. sp., 1601, 1651. as a myrmecophilous species, Structure and development of, 1651. 2226. Horn's classification of the Carabide, 2051. Hypena humuli, 9 = H. scabra. Horse bot-fly, 2238. scabra, 979, 1001, 1976, 2343. chestnut. Flat-headed apple-tree borer in, scabralis, 2343. Hyperchiria io, 420, 809, 1264, 1329, 1352, 1389, 2343. Gregarious worms on, 1192. varia = H. io. HORSE-CHESTNUT. INSECTS AFFECTING.

Cacœcia rileyana, 1192.

Chrysobothris femorata, 1316.

Tortrix rileyana = Caccecia rileyana.

Hypermetamorphoses of Meloidæ, 2191.

textor = H. cunea.

Hyphantria cunea, 112, 242, 454, 456, 460, 619, 1301,

1733, 1849, 1995, 2238, 2378, 2379, 2394.

Hypoderma bovis, 898, 1563, 2238 Imported carpet-heetle, 1690. Hyponomeuta, 2000. currant-worm, 1127, 1570, 2238. malinella, 2000. fly and its parasite, 389 multipunctella, 2000. 381. elm-leaf beetle, 2232, 2304, 2325, 2378, 5-punctella, 1608, 1804. Hypopus, 1708. 2394. Icerya. Bull. No. 15 on, 2389. . gooseberry saw-fly, 228, 332. Life-history of, 2886. werms, 156. in New Zealand. Enemies of, 2479. insects, 140, 225. purchasi, 2386, 2394, 2400, 2401, 2415. and native American insects. an insect injurious to fruit-1115, 1127, orchard Scolytus, 2238. trees, 2401. Original habitat of, 2415. plants and insects, 1339. sacchari, 2415. Importing European parasites, 208. Improved method of spraying trees for protection Ichneumon, 1541. brevipennis, 1570. against insects, 2211. In memoriam B. D. Walsh, 1098. cæruleus, 1802. Cocoons of, 68, 183. Index to reports State entomologist of Missouri, flies, 662, 679, 1648, 1625. 2026. Cocoons of, 851. Indian-corn insects, 23, 188. fly. Army-worm, 1127. Indiana, Butler, A. W. The periodical Cicada in fly mistaken for a wasp, 477. southeastern, 2364. Webster, F. M. The season's observaleucaniæ, 1670. obsoletus n. sp., 1570 = var. of brevitions in, 2418. Indigo. Walshia amorphella on false, 1127. pennis. signatipes, 1802. Industry in the United States. Silk, 2208. from stomach of bluebird, 1878. Inexpert defense, 2476. wing, 50, 197, 385. Inflating Chalcis, 1059. Ichneumonidæ, 2516. Information wanted, 1111. Descriptions of new, 385. Injurious caterpillars, 151. Habits, 158. insects, 126. Ichthyura inclusa, 856. in California, 2074. Inquilines in galleries of common white ant, 1728. Icy lady-bird, 1311. Identity. Mistaken, 1593. and Psenides. Relations of, 41. Ignorance in the North. Entomological, 421. Inquiries answered, 180. South. Entomological, 390. Inquiring friends, 1642, 1656. Illinois. Entomological tour in Southern, 372. Insect boxes. Vertical vs. horizontal, 1963, First report noxious insects of, 378. catching habits of Sarracenia variolaria, Flying locusts in, 1437, 1443. 1385, 1390. Natural History Society. Address bechanges, 388. fore, 5. collection for sale, 2046. New Cynipids of, 41. collections. Naphthaline conce for, 2073. Protection of, 2180. Pseudoneuroptera, 24. A rare capture in, 1211. damage to the corks of wine-bottles, 2477. Six worst insect enemies of the fruitdefoliators. Shade-trees and their, 2378, growers in northern, 377. State entomologist of, 327. destroyer. Paris green as an, 1447. Horticultural Society. Report of oggs, 243, 286, 291. Growth of, 2241. committee on entomology of. 1056. enemies, 1583. Imbricated anout-beetle, 1801, 2291. of the Colorado potato-beetle, 411. Imitative butterflies, 613. growing rice, 1911. Impatiens fulva. Galls on, 852. Imperial moth. Larva of, 1087, 1268. to the rice-plant, 1949. extinguisher. Treat's (review), 581. Importation of insect parasites, 2461. foe of the apple-tree borer, 421. Lestophonus. Further on the, to green corn. A new, 1655. 2541. foes of the apple-tree, 467. Imported cabbage-butterfly, 1127, 2291. army-worm, 12. worm, 2232, 2238, 2291. bark-louse, 417. Bacterial disease of the, hop-vine, 420. 2251. pes, 14. potate, 1558. New remedy for, 2298. in the South, 1714. found about orange-trees, 1798. friend. Another, 879. Successful introduction of a parasite of the, friends and insect foes, 38, 2291, on grape, 1569,

Insect injurious to junipers, 1713. wheat. A new, 2288. killer. Worthlessness of the sparrow as an, 2413. life. 5. and sun spots, 2094. locomotion, 1955. named, 2297. pest. A new, 1686. plagues, 2198. A plant growing out of an, 332, 478. powder. Directions for cultivating pyre thrum for, 1996. Persian, 1485. powders and their use, 1692. ravages, 1464. Trade in, 1986. world, 1466, 1467. Figuier's (review), 408. Remarkable peculiarity in the, 622. Insecticide. London purple as an, 1725. Oxeye daisy as an, 1861. Pyrethrum an important, 2131. ite use as an. 2119. Use of naphthaline as an, 2274. Paris green as an, 1497. Insecticides. Emulsions of petroleum as, 2134. and their value as. 2126, 2200, on garden insects , Alwood, W. B. Tests with, 2382. upon insects affecting garden crops. Experiments with, 2344. Quelques mots sur les, 2283. Two valuable, 1743. Insectivorous plants. Food of, 1499. Insects attracted to light, 178, 2352. Bill providing for the extermination of, 1842. Birds vs., 199, 369. Bluebirds feeding on parasitic and predaceous, 1885. in California. Injurious, 2074. Californian orange, 2373. Catalogues and monographs of, 1765. Centennial, 1511, 1611. clustered on apple-trees, 1263. collecting and preserving, 881, 1057. Comstock's classification of, 2890. Damage to silver plate by, 2154. Death of mules caused by, 1811. is directed. How flight in, 1891. Directions for collecting and preserving. rearing, 14.

and drought, 2100. effects of severe cold on, 1818. enemies of fruit and fruit-trees. Review of Trimble's, 187. in northern Illinois. Six worst, 377. the honey-bee, 1059.

rice-plant, 1949. feeding on sap of black-walnut, 1195. in the flower garden, 936.

27 ENT

Insects as food for man, 2166. found on apple-trees, 731. Fungus diseases of beneficial, 1813. Harmless, 1526. Imported, 140. plants and, 1339. Improved method of spraying trees for protection against, 2211. Injurious, 136. injurious to agriculture. Legislation in regard to, 1468. New, 2055. cereals and forage crops. 2238. cotton in Brazil, 2277. field crops, 2238. fruit and fruit trees, 2238. trees, 16, 31. garden vegetables, 2238. grape-vine, 267, 1081, 1086, 1091, 1102, 1118, 1127, 1130, 1158, 1175, 1202, 1221, 1255, 1281. 1301, 1311, 1329, 1363. live-stock, 2238. orange in Brazil, 2277. sugar-cane in Brazil, 2277. vegetation. Review of Harris's, 568. in Illinois, 6, 52. Legislation to control, 1946. the vine, 2238. of interest to fruit-growers.

Introduction and spread of scale, 2282. by malodorants. Repelling, 2001. Maple-tree, 2279. Mind how you pack your, 435. most destructive to the orange. Successful management of the, 2088. named, 163, 200, 316, 858, 359, 413, 447, 463, 490, 540, 560, 562, 582, 631, 673, 714, 721, 736, 748, 746, 785, 789, 805, 832, 847, 849, 858, 1140, 1147, 1183, 1226, 1273, 1287. to be named, 488, 514, 598.

in the National Museum. The collection of. 2299. Report on the collection of, 2282, 2830.

2368. and native American insects. Imported, 1115, 1127.

Nervous system of, 2225. Notes on our commoner, 1802. on the cleander, 780. in the orchard. Review of Houghton's,

Packard, A.S. Larve of injurious forest, 2267.

Notes on forest, 2253. Number of segments in the head of winged, 2267.

around peach-trees, 660. Pitcher-plant, 1385.

Poisoning noxious, 205. Insects. Poisonous, 2399. Popular remedies for noxious, 73. Preserving, 881, 1057. Regulation of sex in, 1415. in relation to agriculture, 2238. Remedies for various, 2284. Retarded development of, 2040. Salt and vinegar for, 937. Simulium feeding on other, 2177. Some interesting, 1232. Stings of, 116. from stomach of lark, robin, and sunfish, 1926. rock-bass, 1792. stripping the burr-oak, 966. in timber, 918. Trade in, 1986. Transformation of, 528. Unity in coloration of, 50. Use of buckwheat to destroy, 1744. fungus growths to destroy, 1918. poisons to destroy, 1887. used as food. Salt-water, 2203. White willow, 907. of the year, 2289, 2331. Destructive, 2322. Insekten. Einige unserer schädlicherer, 1325. Insidious flower-bug, 1127, 1423, 2418. Instinct of Cicada septendecim, 2144. Instructions to agents of the U.S. Entomological Commission. Supplementary, 1888. Interaction of organisms, 139, 1954. Interest felt in economic entomology in California, Interim committees. Ad, 394. Intermittance of phosphorescence in fire-flies, 1805. Internal mite in fowls, 2157. Introduction of Phylloxera. Laws to prevent the, Invigorator again. Best's fruit-tree, 530. once more. Best's, 545. Io moth, 1329, 1389. caterpillar, 809. Iowa. Economic entomology in, 2197. The hopper in, 1585. Osborn, H. Report upon the insects of the season in, 2418. The seventeen-year Cicada in, 1737. Ipomæa commutata. Aletia feeding, 2343. Ips in calyx of pear. Banded, 1239. fasciatus, 4, 214, 1239. 4-signatus = L fasciatus. Isabella tiger-moth, 1311. Isosoma, 2394. allynii, 2060, 2063. grande n. sp., 2288, 2291, 2316, 2348, 2363, Habits of 2348. hordei, 563, 923, 2060, 2238, 2316, 2394. Larger wheat-straw, 2291. lineare, 2060, 2119.

nigrum == I. bordei.

2363, 2394.

tritici n. sp., 2060, 2063, 2119, 2123, 2316,

Isosoma vitis, 482, 1059, 1127. Wheat, 2119. straw, 2291, 2394. Ithycerus noveboracensis, 16, 306, 652, 659, 869, 1633, 1085, 1188, 1301. Iulus, 430. cœruleo-cinctus, 236. marginatus = Spirobolus marginatus. infested with Gamasus juloides, multistriatus n. sp., 193 = Cambala annulata. virgatus, 261. Ixodes bovis, 404. sp., 834. unipunctata, 1133. Jaeger's North American insects, 84. Japanese mode of packing silk-worm eggs, 1616. Jarring down infested fruit. Codling-moth, 1318. Jassidæ, 737. Jassus, 1766.. sexnotatus, 1766. Jiggers, 412. Johnson, L. Report on cotton-worm, boll-worm, and other insects, 2164. Joint-worm, 49, 62, 563, 923, 2238. Appendix to article on, 1223. fly, 1127. Joint-worms, 2119, 2394. Notes on, 2316. Joints of wheat. Worm in, 1848. Jones, R. W. Observations and experiments on cotton-worm, 2164. Report on cotton insects, 2343. W. J. Report on cotton insects, 2343. Joppidium n. g., 385. ruficeps n. sp., 385. Journal of a State entomologist. One day's, 383. Jumping to conclusions, 253. seeds and galls, 1496, 2163, 2173. spiders, 2302. sumach-beetle, 1363. tree-cricket, 1329. Juniper. Dapsilia rutilans on, 1713, 1721. web-worm, 1721. Junipers. Insects injurious to, 1713. Juniperus sabina. Insects injurious to, 698. Junonia lavinia, 753. Kansas bombardier-beetle, 1311. The locust in, 1591. and Missouri this fall. No locust injury in, 1433. Report on grasshopper question to the governor of, 1573. Scorpion in, 1119. Silk culture in, 1542. Kartoffel-Käfer. Neue, 919. Katydid. Angular-winged, 1363. Broad-winged, 1329, 1363. Domesticated, 1536. Eggs of the, 516, 1005. angular, 1518. oblong, 569. Narrow-winged, 1363. Oblong-winged, 1829, 1863. See Catydid. Katydids, 1363.

Kentucky. Coleopterous cave fauna of, 2033. Kermes galliformis n. sp., 1972. Kerosene emulsion, 2291. as a means against cotton insects, 2164. orange insects, 2164. Kiesenwetter. Obituary, 1819. Killer. Cottonwood, 1654. Killers. Bee, 1543. Killing apple-worms by machinery, 769. Kingdom. Animal, 393. Klippart's wheat plant (review), 186. Knot once more. Black, 980. Knots on apple-tree roots caused by root-lice, 1187. Knotweed geometer, 1059. Hæmatopis grataria on, 1059. Knowledge useless ! Is any, 1135. Koebele, A. Experiments on cottony cushionscale, 2394. the red scale, 2394. Notes on locusts about Folsom, Cal., Report of experiments against scale insects, 2418. Koeppen, F. T. Account of the Hessian-fly, 2267. Labena. Useful, 1423. Labia, 2267. Lac insects, 2119 Lace-wing fly, 533, 1428. Lace-wing larva, 1059. Weeping, 1127. Lachnosterna fusca, 3, 54, 68, 300, 332, 410, 478, 500, 594, 640, 865, 1020, 1059, 1064, 1072, 1307, 1313, 1329, 1436, 1440, 1522, 1803, 1812, 2238, 2394. pilosicollis = L. tristis. quercina = L. fusca. quercus, 372. tristis, 966. Lachnus caryse, 27. plantanicola n. sp., 2138. strobi, 265, 320, 1039. Lackey moth. American, 1301. Lacordaire, J. T. Death of, 1285. Ladder spider, 1299. Lady-bird. Ashy-gray, 2119. Blood-red, 2119. Cactus, 2119. Fifteen-spotted, 959, 1811. Icy, 1311. and its larva. Northern, 1289, Spotted, 599. Twice-stabbed, 38, 1883. Lady-birds, 1059, 1127, 1423, 1558, 2119. Swarms of, 824. Lafayette, Ind., Webster, F. M. Experiments at, Lagoa opercularis, 145, 796, 1748. Lake Superior. Simulium from, 2032. Lamellicornia, 1440. Lampronota, 1878. amphimilæna n. sp., 385. breviventris n. sp., 385.

imitatrix n. sp., 385.

interpellata n. sp., 385.

pictiventris n. sp., 385.

Lampyridæ, 1705, 1805, 2036.. Revision of the, 1819. Lampyris noctiluca, 1097, 1917. Laphria thoracica = Dasyllis thoracica. Laphygma autumnalis = L. frugiperda. frugiperda, 1127, 1256, 1267, 1282, 1301, 1400, 2011, 2343. fulvosa n. v., 1301 = var. of L. frugiobscura, n. v., 1301 = var. of L. frugiperda. Laporte. Obituary, 1817. Lappet caterpillars on the apple, 972. apple-trees, 812. Larch. Nematus erichsoni on, 2232. saw-flv, 2232. Large Asilus fly, 1269. black potato-beetles, 1206. compound gall on grape-vine, 720. dragon-fly, 759. fish-fly, 712, 903. gray straight-horned snout-beetle, 1033. green caterpillars on the apple, 1076. worms in a peach, 661. moth on apple-trees, 1028. phosphorescent larva, 1874. saw-fly, 1514. silken cocoon, 604. water-beetle, 750, 816. white-scale on Acacias, 1730. willow-worm, 1380. worm on apple-trees, 1048. Larger cabbage butterfly, 2232. wheat straw Isosoma, 2291. Lark. Insect from stomach of a, 1926. Larva boring along the axis of apple twig, 1850. injurious to cotton squares. Butterfly, 1872. Large phosphorescent, 1874. Moth issuing from a, 1779. Larve. Aquatic, 1851. Habits of, 346. in the human bowels, 382. of injurious forest insects, 2267. named, 1264. Preserving, 1300. in stomach of black-bass, 1792. blue-bird, 1871. Larval characteristics of Corydalus and Chauliodes. 1652. characters and habits of blister-beetles, 1600. Epicauta, 1600. Macrobasis, 1600. habits of bee-flies. Bombylidæ, 1947, 1970. Dexidse, 2260. Epicauta and Henous, 1387. Sphenophori that attack corn, 2030. life as influenced by food. Number of molts and length of, 2167. stages and habits of bee-fly. Hirmoneura, Larviform females in the Phengodini. Luminous, 2397, 2402,

Lasius latipes, 27.

Lasioderma serricorne, 551.

Lasioptera vitis, 720, 1329.

Law. Carrying out the, 1993.

Lawn. Beetles swarming about, 718.

Lepidoptera of the Outer Hebridee, 2058.

Lepidopterological notes, 1999, 2613, 2166.

Laws to prevent the introduction of Phylloxera, Lepidopteron. A myrmecophilous, 2214. Lepidopterous case-bearer, 269. 2019. Leaf-beetle. Streaked cottonwood, 2201. larvæ. Dried leaves as feed for, bug. Ash gray, 1127, 1423. 2150 bugs, 400. Fleas feeding on, 2116. crumpler. Apple, 574, 1580. Remarkable, 44. in Georgia. Rascal, 1962. Lepidosaphida, 1201. Rascal, 341, 373, 1311. Lepiopomus pallidus. Insects from the stemach folder. Cranberry, 2291. of, 1792. Grape, 1301, 1579. Leptobatus illinoiensis n. sp., 385 = Excesses galls and caterpillars on sugar berry, 762. illinoiensis. Leptostylus aculiferus, 673. on the grape-vine, 1716. hopper, 36. Leptura capitata, 200. injurious to small grains. A new, Leptus americanus n. sp., 1226 = Tetranychus americanus. hoppers, 2362. irritans n. sp., 1326 = Tetranychus irrion celery, 452. tans. Lesser apple-leaf folder, 1311. of the grape, 399. Grape-vine, 484. locust, 2363. migratory locust, 2232. injuring wheat, 1766. miner on white-oak, 1879. pine-burer, 2267. miners of the locust, 451. Prionus, 2267. roller. Strawberry, 1574. Lestes insequalis n. sp., 24. Leucania albilinea, 1507, 1570, 1610. rollers. Descriptions of new, 1969. tyer. The green, 1311. unipuncta, 6, 7, 8, 9, 10, 11, 12, 17, 52, 326, Leafy oak-gall, 774. 647, 670, 879, 906, 1127, 1267, Leather-beetle, 2863. 1282, 1400, 1442, 1482, 1481, Lebia grandis, 365, 1218. 1551, 1570, 1679, 1809, 1835, 1856, 1877, 1885, 1886, 1953, Great, 1301. Lecanium, 1. 2001, 2009, 2086, 2087, 2020, 2119, 2122, 2235, 2239, 2253, acericola n. sp., 389 = Pulvinaria innumerabilia. 2262, 2367, 2843. macluræ n. sp., 389 = Pulvinaria innu-Additional notes on, 2001. merabilia. Complete life-history of, 1678. Hibernation, 7, 8, 2006. oleæ, 1303. rosse == L. olese. Parasites of, 6, 10, 17, 1679. Leucopis, 154, 160, 174, 1539. sp. on blackberry, 117. Phylloxera, 2150. china-tree, 1964. magnolia, 1377. Libellulidæ, 1707, 1709. plum, 107. Library pest. Croton bug as a, 1717. Lice on calves, 889. sugar-maple, 1004. vitis = Pulvinaria vitis. horses, 258. pigs, 900. Leconte, J. L. Death of, 2246. snow-balls, 1184. Tribute to the memory of, 2264. Leconte's pine-worm, 1570. Lichtenstein, J. Obituary, 2370. Lederer, J. Death of, 1285. Lichtenstein's theory as to dimorphic asexual females, 2072. Le Duc, W. G. Letter to, 1684. Life. Tenacity of, 1755. Legged maple-borer, 1363. Light. Insects attracted to, 178, 2352. Legislation to control insects injurious to vege-Lightning-hopper. Frosted, 1329. tation, 1946. in regard to insects injurious to agrihoppers, 737. Lignified anake of Brazil, 2136. culture, 1468. Lignivorous. Rhyssa not, 2286. Lema trilineata, 119, 126, 135, 136, 138, 185, 401, 565, Ligyrus relictus, 3, 54, 68, 1440. 925, 1059, 1328, 1558, 1593. rugiceps, 1794, 2080. Length of larval life as influenced by food. Number of molts and, 2167. Lilac-borer, 844. the thread of the silk-worm, 1359. LILAC. INSECTS APPECTING. Leopard moth. Great, 1311. Egeria syrings = Podosesia syrings. Podosesia syringæ, 844. Lepidium vs. bed-bugs, 1741. Limacodes, 847, 1150. Lepidoptera, 13.9, 2267, 2340. by alinus n. sp., 40 = Phobetron byali-Hudson Bay, 1985. Migrations of, 1770. New lists of North American, 2132. pithecium = Phobetron pithecium. Notes on South American, 1784. scapha, 40.

Locust, important observations on the Rocky Limacodes tetradactvlus n. sp., 40. Limenitis disippus, 613, 1193, 1217, 1301, 1305, 1306, Mountain, 1571. injuries. New method of counteracting, 1340. ursula, 145, 1217, 1305, 1806, 1840. Limneria fugitiva, 418, 1311. injury in Kansas and Missouri this fall, lophyri n. sp., 1570. pallipes, 2378. the near future. Probabilities Limpophilus, 2267. of, 2355. Lina lapponica, 1884. injury next spring, 1555. populi, 1834. summer. Anticipated, 1615. scripta, 1664, 1884, 2291. LOCUST. INSECTS AFFECTING. Arhopalus robiniæ = Cyllene robiniæ. tremulæ, 1834. Linden and ash destroyers, 1849. Cossus robiniza, 37, 1121. LINDEN. INSECTS APPECTING. Clytus robinise = Cyllene robinise. Hyphantria cunea, 1849. Cyllene robinise, 87, 472, 827, 921, 1196, 1250, textor = H. cunes. 1288, 1516. Odontota rubra, 1849. Hispa scutellaris = Odontota dorsalis. Linoceras junceus, 543, 1827. Nitidulidæ, 87. L'insectologie agricole, (review), 506, 568. Odontota dorsalis, 451. Lintner's first report, 2244, 2254. Spermophagus robiniæ, 1026, 1474. Linum. Acridide that eat, 1645. Xyleutes robinize = Cossus mobinize. Lissorhoptrus simplex, 273, 1911, 1973, 2119. Logust invasion with the occurrence of drought. Listotrophus cingulatus, 805. Connection of, 1422, Lithocolletis cincinnatiella, 1879. Leaf miners of the, 45L guttifinitella. Mandible of, 1984. Lecture on the Rocky Mountain, 1493. Lithophane antennata, 1182. Lesser migratory, 2232, 2863. cinerea, 1301. mite, 1423, 1625, 1643, 1959. Little Cicada, 1242. in Montana in 1880, Bruner, L. Rocky known facts about well-known animals, 2071. Mountain, 2267. multiplication and migration, Swinton, Turk and its crescent, 829. Live-stock. Insects injurious to, 2238. A. H. Solar physics and earthquake Lives. Two useful, 2370. commotion applied to, 2267. Lixus. Larval habits of, 2404. Natural history of the Rocky Mountain. macer, 2404. notes. Miscellaneous, 2267. parcus on Amelanchier, 2404. Lobesia botrana = Eudemis botrana. in 1876, 1521. Loco weed. Walshia amorphelia bred from, 1127, 1880, Martin, J. Rocky Mountain, 2267. 2356. 1885, Bruner, L. Abundance of the Locomotion. Insect, 1955. Rocky Mountain, 2363. Locust, 2366. and other insects in the Northwest during Ash-colored, 2363. the summer of 1883, Bruner, L. Obser-Atlantic migratory, 1423, 1625. vations on the Rocky Mountain, 2277. borer, 921, 1121, 1196, 1288. Packard, A. S. Development of the, 2267. Hickory w., 1516. The periodical Cicada alias the seventeenborers, 37, 472. year and the thirteen-year, 1159. Bruner, L. Observations in the Northpest, 1501. west on the Rocky Mount-Philosophy of the movements of the ain, 2165. Rocky Mountain, 1009. California migratory, 2368. plague, 1473. Californian, 1959. in the United States, 1625. Devastating, 2863. probabilities for 1882, 2057. Differential, 1423, 2363. prospects, 1527, 1562, 1582. eggs, 1548. in southwest Missouri this fall, in Asia Minor. Bombylid larvæ 1596. destroying, 2118. ravages in California, 1959. Condition of, 1567. 1880 and 1881. Chronology of, 2267. Experiments with, 1572. Red-legged, 2363, 1423, 1625. Expected advent of the, 2307. experience, 1617. report to governor of Kansas, 1573. flights, 1590. Rocky Mountain, 1423, 1451, 1452, 1482, 1588, in Dakota, 2007. 1557, 1570, 1625, 1643, 1959, 2267, 2291.

east of the Mississippi, 1549, 1618.

Mountain, 2267.

seed-weevil. Honey, 1026. Geographical distribution of the Rocky Seventeen-year, 370, 884, 1489. or grasshopper. Governors of Western swarms that devastate the trans-Missis-States on the Rocky sippi country, 1674. Mountain, 1557. theory wanted. New, 1532.

scourge. Rocky Mountain, 1629.

Locust. Two-striped, 1423, 2363.

weevil. Honey, 1474. in Wyoming, Montana, etc., in 1881, Bruner, L. The Rocky Mountain, 2267. Yellow, 2368. Locustidæ, 929, 1453. Locusts, 926, 1349, 1863, 2418. again, 1550. Bruner, L. List of North American, 2267. Changes in vegetation caused by, 1495. in Dardanelles. Sarcophaga lineata destructive to, 2075. Destruction of young or unfledged, 1577. Destructive, 2363. Dimorphism of, 1889. Ditching for young, 1488. eat the castor-bean, 1645. about Folsom, Cal., Koebele, A. Notes on, 2363. as food for man, 1481. Grasshoppers and, 26, 147, 929. Habits of young or unfledged, 1578. hatching? Are the, 1566. Mistaken identity. Are the, 1598. How to destroy, 1446. in Illinois. Flying, 1437, 1443. in Kansas, 1591. Literature of destructive, 1959. and locusts, 1994. in Nevada, 2024. New remedy against destructive, 2328. next spring. Injury by, 1555. Notes on, 1456. 1880. Chipman, A. J. Notes on, 2287. Prairie fires and hateful, 1453. Professor Riley and the, 1434. of San Joaquin Valley, Cal. Coquillett, D. W. Report on the, 2363. sting ! Do, 371. in Texas in spring of 1886, Bruner, L. Report on, 2382. in the West, 2044. and the Western cricket, Bruner, L. Notes on other, 2267. in western Missouri. Ravages of young, 1492. Loew, H. Description of the rye gall-gnat, 2267. Lombardy poplars. Caterpillars on, 571. London purple as an insecticide, 1725. and Paris green, 2021. Long-horned Diabrotica, 1905. legs. Grand-daddy, 838. sting. Delicate, 1329. tailed Ophion, 1311. Longicorn beetles. Food of, 1902. borers, 95. Longicorns in pine and cedar, 319. Longitarsus, 636. Look out for the eggs of the apple-tree plantlouse, 507. Lopha 4-maculata = Bembidium 4-maculatum. Lophyrus abbotii, 465, 927, 956, 1057, 1570. abietis, 115. lecontei, 985, 1011, 1057, 1570.

Lozotænia rosaceana — Cacœcia rosacean Lubber grasshopper, 2119. Lucanus dama, 784. elaphus, 305, 755, 957, 1517. Lucidota atra, 358. Lucilia macellaria, 209, 1890, 1921, 2158, 2199, 2255. Ludius attenuatus, 224. Lumbricus, 1304. Luminosity of fire-flies, 1840. Luminous larviform females of Phengodini, 2307, 2402. Luna moth, 776. silk-worm, 1311. Lungs. Larva of Scenopinus from human, 1348. Luperus brunneus, 1799. morulus, 1799. noxius = L. brunneus. Lure for moths, 1695. Lyda sp., 656. Lydella doryphore n.sp., 1059 = Exorista doryphoræ. Lygus lineolaris = L. pratensis. pratensis, 2, 81, 76, 682, 1127, 1213, 1219, 2235, 2291, 2363. Lymexylidæ, 575. Lymexylon navale, 575, 1135. Lytta ænea = Pomphopæa ænea. atrate = Epicauta pennsylvanica. cinerea = Macrobasis unicolor. fabricii == Macrobasis unicolor. marginata = Epicauta cinerea. murina = Macrobasis unicolor. sayi = Pomphopœa sayi. tarsalis = Pomphopœa tarsalis. vittata = Epicauta vittata. Machine. A satisfactory grasshopper, 1592. Machinery for destroying the cotton-worm, Barnard, W. S. Tests of, 2253. Killing apple-worms by, 769. McLain, N. W. Apicultural experiments, 2362. Experiments in apiculture, 2394. Report on experiments in apiculture, 2363. Maclura aurantiaca as food for Sericaria, 2234. Silk-worms fed on, 1341, 1542, 1609. McMurtrie, W. Tests of silk-fiber from coccoms raised at the Department, 2253. Macrobasis, 1600, 1648, 1651, 2238. albida, 1796. Macrobasis, Larval characters and habits, 1602. murina = M. unicolor. unicolor, 38, 134, 185, 347, 362, 401, 470, 736, 912, 1044, 1209, 1558, 2248. Macrocentrus delicatus, 1334. Macrodactylus subspinosus, 249, 361, 373, 565, 748, 1075, 1278, 1329, 1357, 1375, 1478, 1583, 2248. Macrogomphus! spiniceps n. sp., 24. Macromia flavipenuis n. sp., 24. illinoiensis n. sp., 24. Macronema zebratum, 372. Macrosila carolina = Protoparce carolina. cluentius, 1784. 5-maculatus = Protoparce celeus. rustica = Protoparce rustica.

100

Louisiana. Destructive cricket im, 2384.

Mad. Entomology indeed run, 227, 1224.

Madarus ampelopsidos = Ampeloglypter ater.

vitis n. sp., 1050 = Ampeloglypter sessetris.

Madras. Coffee-borer in, 498.

Macklin, F. W. Obituary, 2161.

Maggots in sauce, 1607.

Magnolia. Lecanium on, 1377.

Scale-insect on, 1377.

Maia moth, 1329.

Mails. Queen bees in the, 1762.

Maine. New potato-bug in, 119.

MAIZE. INSECTS AFFECTING, 2119.

Achatodes zeze, 1927.

Agrotidæ, 1522.

Elateridæ, 1522.

Lachnosterna fusca, 1522.

quercina = L. fusca.

Malaria. Mosquitoes vs., 2162.

Mallodon melanopus, 2291.

Mallophaga, 258, 497.

Mallophora orcina, 1959.

Malodorants. Repelling insects by, 2091.

Malva sylvestris. Eryunis alcese boring in stems

Mameetra, 2238.

Cabbage, 2232.

chenopodi = M. trifolii.

picta, 179, 1056, 1127, 2232.

subjuncta, 2291.

trifolii, 229, 281, 2232.

Man. Effect of Paris green on, 1427.

Insects as food for, 2166.

Locusts as food for, 1481.

Mantidæ, 2267.

Mantis, 2267.

carolina = Phasmomantis carolina.

Preying, 457.

or rear-horse. Eggs of, 1060.

Supposed eggs of preying, 1002.

Mantispa, 2267.

brunnea, 1243.

Mantispian. Brown, 1243.

Many-banded robber, 1059, 1423.

Maple-bark lice, 1004.

lonse, 344. Beetle on sugar, 1014.

borer. Legged, 1363.

A new (f) segerian, 1360.

Eggs on sugar, 350.

Maple Galls on leaves of soft, 445.

MAPLE. INSECTS AFFECTING.

Acarus aceris-crumena, 1265.

Acronycta americana, 841.

Actony cua americana, est.

Ægeria acerni, 743, 1063, 1360, 1363.

Anisota rubicunda = Dryocampa rubicunda.

Arachnid, 445.

Attacus cecropia, 841.

Catydid, 665.

Ceratocampa imperialis = Eacles imperialis.

Chrysobothris femorata, 1250.

Clisiocampa americana, 350.

Clytus speciosus == Plagionotus speciosus.

Coccid, 344.

Dryocampa rubicunda, 841, 915, 1329.

Escles imperialis, 1268.

MAPLE. INSECTS AFFECTING-Continued.

Eburia 4-geminata, 1014.

Lecanium acericola — Pulvinaria innumera-

sp., 1004.

Orgyia antiqua, 1797.

Plagionotus speciosus, 915.

Pseudococus aceris, 1890.

Pulvinaria innumerabilis, 1515, 1816, 2279.

Sciara ocellaris, 2119.

Teles polyphemus, 841.

Trochilium acericolum = Ægeria acerui.

acerni = Ægeria acerni.

Maple. Mite-gall on sugar, 1265.

Ocellate leaf-gall of the red, 2119.

scale. Cottony, 1816, 2291.

Scale insect on, 1890.

tree insects, 2279.

twigs. Rows of eggs on, 665.

worm. Green striped, 1329.

worms, 841.

Maples. Cottony scale on, 1515.

Flat-headed borer in soft, 1250.

Mare's nest. Finding a, 86.

Margined blister-beetle, 1059, 1558.

Martin, J. Report on the Rocky Mountain locust, 1959.

> Report on the Pocky Mountain locust in 1880, 2267.

Mary Chalcis-fly, 1311.

Masicera archippivora n. sp., 1301.

Mason-bee cells. Remarkable new genus of Meloids infesting, 1601.

Mass of eggs, 240.

mistakes, 234.

Massachusetts. Periodical Cicada in, 2321.

ern, 2216.

Massospora cicadina infesting Cicada, 1809, 2112.

May-beetle, 300, 865, 1020.

Egg of the common, 1329.

beetles swarming in Alabama, 1812.

fly, 283.

Meadow enemy, 1368.

lark. Beetles in stomach of, 1013.

worms, 947.

Meal sack. Worm infesting, 1896.

worms, 191.

Mealy bugs. Structure of, 2119.

Measuring-worms, 75.
Megastizus brevipennis, 375.

Megathymus coloradensis n. var., 1602.

yuccæ, 1420, 1465, 1482, 1570, 1602.

Additional notes, 1602.

Note on, 1465.

Megatoma serra, 1352.

Megilla maculata, 6, 94, 188, 589.

Food-habits of, 2145.

Melampsalta parvula, 1242. Melancholy chafer, 522.

in apples, 842.

Melanactes, 1874.

Melanippe montanata, 2224.

Melanolestes picipes, 314, 1253. Melanophila sp., 2267.

Melanophora? diabrotica, 2260.

Microgaster, Military, 1127. Melanoplus atlanis = Caloptenus atlanis. destructor == Caloptenus destructor. devastator = Caloptenus devastator. Melanotus communis, 351, 358. incertus, 40, 846. Melasoma lapponicum = Lina lapponica. populi = Lina populi. Microlepidoptera. Works on North American, scriptum = Lina scripta. tremulæ = Lina tremulæ. Melissopus n. g., 1969. aurichalceana n. sp., 1969 latiferreans, 1969. Melittia ceto, 125, 126, 248, 377, 1083. gloriosa 2410. Meloe, 1500, 1600, 2267. angusticollis, 1387. barbarus, 1600. proscarabæus, 2083. Meloidæ, 912, 1230, 1390, 1600, 1601, 1651, 1858, 2002, 2072, 2118, 2267. Fire cure for, 121. Hypermetamorphoses of, 2191. infesting potato, 48. Means against, 794. Remarkable genus of, 1601. Remarks on, 1600. Triungulin of, 2082. Meloini, 1601. Melolontha philophaga = Lachnosterna fusca. Melon. Bug on, 897. bugs. Satisfactory remedy for, 2236. Diabrotica vittata on, 897, 2236, 2238, Membracidæ, 737. Membracis ampelopsidis, 1183. Memoriam B. D. Walsh. In, 1098. Meracantha contracta, 1871. Merinus lævis, 1153. Meriaus destructor, 1581, 2382. subapterus n. sp., 2832. Mermis, 2363. Meromyza, 1461. americana, 727, 1058, 1059, 1506, 1589, 1848, 1875, 2291, 2894. Mesochorus. Glassy, 1127. vitreus n. sp., 6, 1670. Mesoleius, 385. Metamymar n. g., 2343. aleurodis n. sp., 2343. Metapodius femoratus, 373, 775. pasulus = M. femoratus. Meteorus hyphantrize n. ep., 2378, 2394. Methods of destroying scale-insects, 2119. Mexican honey-ants. Peculiarities of, 1417. Mexico. Report on cotton crop and its enemies in. 2343. Mezium americanum, 2346. Michigan apples and codling-moth, 1677. Microcentrum retinervis, 241, 247, 1363, 1518, 1536. Microdon globosus, 1789. Microgaster, 77, 158, 183, 490, 662, 679, 717, 861, 1217, 1290. Disippus, 1301. gelechise n. sp., 1959. Life-habits of, 64. limenitidis n. sp., 1301 = Apanteles

limenitidia.

ceratomise n. sp., 1125, 1134, 1969. gortynæ n. sp., 1900. Micropterus salmoides. Larvæ in stomach of Micropus leucopterus = Blissus leucopterus. Midae fulvipes n. sp., 40. Midge, 167. Clover-seed, 2291. Pear, 2363. Wheat, 109, 110, 142, 216, 280, 292, 428, 711, 1512, 2238, 2291, 2363. Migration of butterflies, 1770, 1991. and hibernation of Aletia argillacea. 1689. of plant-lice from one plant to another, 2017. Migratory butterflies, 1622, 1635. locust. Atlantic, 1423. Lesser, 2232. Military Microgaster, 1127. Milkweed. Danais archippus on, 1535. weevil, 711. Million dollars. That hundred and fifty, 1617. Plums for the, 501. Mills. The same in flouring 1261. Milyas cinctus, 777, 1294, 1431, 1450, 1672, 2048. Mimicry and protective resemblances, 1991, 1340. Mind how you pack your insects, 435. Minings on apple-twigs. Galls and, 552. Minneapolis. Entomolo y at, 2212. Minnesota. A State entomologist for, 1198. Minot, C. S. Anatomy of Aletia, 2243. Minute borers in cherry, peach, and plum tr.es, 1940. Miscellaneous, 735. Miscellaneous notes on orange insecta, 2164. Misnamed gall-moth, 1127. Mississippi country. The locust swarms that devastate the trans, 1674 Description of Aphididze from west of the, 1678. floods. One effect of the, 2080. Locust flights east of the, 1549, 1613. valley. The Buffalo-gnat problem in the lower, 2416. Ailanthus silk worm in, 1460. Missouri. bee-killer, 1127. entomological reports, 1680. Noctuide in the, 2069. Entomology in, 1361. this fall. Locust prospects is southwest, 1596. No locust injury in Kansas and, 1433. Murtfeldt, M. E. Notes from, 2291.

Notes on North American, 1960.

1264

Microgasters. Notes on, 2097.

Microplitis actuosus, 1125, 1134.

parasitic on Protoparce celeus, 155.

sp. parasitic on Thecla (poca?), 1872.

Missouri. Murtfeldt, M. E. Notes for 1886 from, Mosey rose-gall, 570, 1166. Oyster-shell bark-louse in, 1169. Ravages of young locusts in western, 1492. Report State entomologist of, 1, 1059. 2. 1127. 3, 1301. 4, 1811. 5, 1329. 6, 1363. 7, 1428. 8. 1482. 9. 1570. Index, 2026. State Horticultural Society. Report committee on entomology, 1113, 1127. Supposed bark-lice eggs in, 1984. Mistaken identity, 1598. Mistakes. A mass of, 284. Mite. On an extensile penetrating organ in a gamasid, 1626. in fowls. Internal, 2157. gall on sugar-maple, 1265. Locust, 1423, 1625, 1648, 1959. parasites of the Colorado potato-beetle, 1505. Phylloxers, 1363. Red, 1959. Scarlet, 1470. Silky, 1423, 1625. transformations, 1618. Mites, 356. Cannibal, 624. Classification of, 1992. Descriptions of new subterranean, 1870. Egg-infesting, 1637. on grasshoppers, 146. mistaken for plant-lice, 59. parasitic on beetles, 315. Transformations of the red, 1632. Mniszech, G. V. Obituary, 2054. Mold and Phylloxera, 1807. Mole crickets, 1270. Molts and length of larval life as influenced by food. Number of, 2167. Monell, J. Notes on Aphidida, 1678. Monocesta coryli, 1721. Monocrepidius vespertinus, 351. Monographs again, 1765, 1828. Monohammus titillator, 1324. Monophadnus rubi, 663, 761, 1212, 1641. Monostegia rosse, 672, 984, 1780. Monstrosities. Rare, 2224. Montana in 1880, Bruner, L. The Rocky Mountain locust in, 2267. 1881, Bruner, L. The Rocky Mountain locust in Wyoming and, 2267. Monterey again. Butterfly-tree of, 2052.

More good words, 532.

Mosquitoss, 608.

universal remodies, 278.

Morus as food for Sericaria, 2234.

w. malaria, 2162.

Morning-glory. Coptocycla aurichalces on, 1510.

Sphinx. White-lined, 630, 1198, 1301, 2291.

Most precious bug, 575. Moth on apple-tree. Large, 1028. Bee, 716, 904, 1059, 1508, 1728. breeding. Pedigree, 2381. eggs, 611. Handmaid, 2222. issuing from a larva, 1779. named, 807, 1241. Viviparity of a, 2153. Motherless race. That fatherless and, 1650. Moths attracted by falling water, 2108. and butterflies caught by the tongue, 1761. caught in Alabama, 1774. in cushions. Ravages of, 1731. Descriptions of two new, 1411. Galls made by, 1127. Lure for, 1695. mistaken for Aletia, 1976. by Physianthus albens. Capture of, 1888, 1419, 1761. Sugaring for, 1300. Mottled tortoise-beetle, 1127. Moulting in Orgyia. Variable, 2379. Mounding peach-trees, 587. again, 617, 653. Mountain ash. Ash and, 705. MOUNTAIN ASH. INSECTS APPECTING. f Aphie mali, 579. Aspidiotus harrisii = Chionaspis furfurus. Chionaspis furfurus, 579. Mountain ash. Plant-louse eggs on apple and, 579. Mud-wasp and parasites, 1827. spider egg-nest, 1847. Mulberry. Sericaria mori on, 1311, 1341, 2234. silk-worm, 1311. Osage orange for the, 1220, 1286. for the silk-worm. Osage orange vs., 9224 Mulch hay. Worms under, 1161. Mules caused by insects. Death of, 1811. Murgantia histrionica, 849, 1009, 1207, 1311, 1534, 2238, 2263, 2291. Murky ground-beetle, 486. Murmid.us. Habits of, 2217. Murmidius ovalis, 2217. Murtfeldt, M. E. Notes from Missouri, 2291. for the season of 1886,2382. Musca, 1390. domestica, 815, 864, 2071, 2078. harpyia = M. domestica. Muscid, 459. from stomach of Ohio shad, 1853. Muscidse, 153, 1880, 2256. Muscle shaped bark-louse on apple-trees South, Museum pest. Dinoderus pusillus as a, 2104. A new, 2109. once more, 492. Trogoderma tarsale as a, 2139. pests, 246, 462, 572, 726. again, 483. Mutilla coccinea = Sphærophthalma occidentalis. Mycetophila peraics n. sp., 653, 660, 963, 1056.

Necrological, 1817.

Mycetophila sp., 419. Mychocerus, 2217. Mygale hentzii, 466, 493, 521, 823, 1178. and Pepsis formosa, 1619. Myochrous denticollis, 2418. Myriapoda, 219, 224. Poisonous, 2399. Myriapoda, 1625. Myrmecocystus hortusdeorum, 1942. mexicanus, 1417. Myrmecophila, 2202. Myrmecophilous coleopters, 2105. lepidoptera, 2214. species. Hymenorus rufipes as Myrmica minuta, 1413. Myrmicocela ochracella, 2261. Mysia 15-punctata = Anatis 15-punctata. Mystery. Dr. Hagens', 1943. in reference to Pronuba yuccasella, Mytilaspis, 1. conchiformis = M. pomorum. pomicorticis n. sp., 1329 = M. pomopomorum, 188, 201, 345, 372, 878, 377, 423, 565, 896, 944, 951, 973, 982, 989, 994, 1007, 1008, 1012, 1040, 1056, 1059, 1084, 1169, 1246, 1329, 1362, 1397, 1552, 1583, 1774, 2088, 2238. M v zus cerasi, 678, 960, 1067, 1207, 1251. ribis, 304, 322, 735, 970. Nails into fruit-trees. Driving, 87. Nanodes tamarisci, 1496. Nantucket. Pine-moth of, 2183. Naphthaline cones, 2133. for insect collections, 2073. as an insecticide. Use of, 2274. Napping. An entomologist caught, 1100. Narrow-winged katydid, 1363. National Museum. The collection of insects in, 2299, 2390. Reports on the department of insects in the, 2282, 2330, 2368. Native American insects. Imported and, 1115, apple-tree bark-lice, 1152. bark-lice on apple-trees, 1061. currant-worm, 1570, 2238. grape-vines. Cause of deterioration in some of our, 1342. home. Colorado potato-beetles, 1462. plums. Wier, D. B., 2388. Naturaliste Canadien. Le (review), 546. Naturalists. Entomological speculations of the New England school of, 44. Neal, J. C. Observations and experiments, 2164. Nebraska bee-killer, 1127, 1311. Bruner, L. Notes from, 2291. Report of the season's observations in, 2418.

insects, Bruner, L. Report on, 2382.

rufipes, 1363, 1367, 1703.

Necrobia ruficollis, 1363.

Necrophorus marginatus, 315, 845. Nectarophora granaria, 126, 1806, 2394. rudbeckiæ, 27, 2205. Needle. Devil's darning, 1709. Negro bug. Flea-like, 1127, 1423, 2291, 2418. Nematus erichsoni, 2198, 2232, 2355. fur n. sp., 197. hospes n. sp., 197. inquilinus n. sp., 197. integer, 2232. mendicus n. sp., 197, 821. ribesii, 78, 140, 228, 333, 340, 364, 380, 381, 756, 772, 1031, 1224, 1570, 1696, 2238. ribis = N. ribesii. s.-desmodioides n. sp., 197. s.-pisum n. sp., 197. s.-pomum n. sp., 197. Spru 3e, 2232. ventralis, 907, 917, 924, 1211. ventricosus = N. ribesii. Nemestrinidæ, 2186. Nemobius vittatus, 1287. Nemorsea leucanise, 6, 12, 647, 1112, 1127, 1670. Neoclytus capræa, 560,582, 1849. erythrocephalus, 1798. Nephelodes violans, 1885, 1990. Nephila plumipes, 830. Peculiarities of, 1416. Nepidæ, 797. Nerium oleander. Moths caught by, 1419, 1761. Nervous system of articulates. Yersin. Function of the, 1959. insecta, 2225. Phylloxera, 1687. Neuroptera, 1329, 1736, 2267. North American, 24, 39. Neuroterus q.-saltatorius, 1496, 2163, 2173. Nevada. Locust in, 2024. New Curculio humbug, 680. New England and New York. Packard, A. S. Causes of destruction of evergreen and forest trees in, 2232, 2291, 2363, 2364, 2382. school of naturalists. Entomological speculations of the, 44. entomological journal, 2056. Hampshire. Colorado potato-beetle in, 1859. humbug, 122. Mexico. Blister beetles from, 1796. philosophy. Old and, 392. York. Chinch-bug in, 2237, 2252, 2271. Colorado potato-beetle in, 1379. and Eastern States. Supposed armyworm in, 1990. Entomology in, 2244, 2254. Packard, A.S. Causes of destruction of evergreen and forest trees in New England and, 2232, 2291, 2363, 2364, 2382. State entomologist for, 1863. weevil, 306, 652, 659, 1301. in apple trees, 1085. without a State entomologist, 1330. News. Recent cotton-worm articles in the, 1719. Nicotiana tabacum. Flea-beetle eating, 1782.

Night. How the Curculio flies by, 706. Nursery. Holes around the roots of young ash Ninth report State entomologist of Missouri, trees in the, 471. 1570. Nuts. Gall, 1560. Niptus hololeucus, 2154. Nymphalidæ, 1301. Nitidula bipustulata, 214. Pupation of, 1704. Nitidulidæ, 37, 214. Nysius angustatus, 1329, 1357, 1402, 1453, 2048, 2291. Noctuid. A unique and beautiful, 2189. destructor n. sp., 1329 = N. angustatus. Noctuidæ, 455, 787, 1419, 1761. n. sp., 1317. of the Missouri reports, 2069. Oak apple, 1448. North America. Notice of Grote's bark louse, 159. borer, 2267. iliustrated essay on the, 2148. Parasites of, 849. Clytus, 2267. taken at Orono, Me., 2160. coccid mistaken for a gall, 1972. Nola sorghiella n. sp., 2119. fig gall, 745. Nomenclature. Scientific, 303, 768. gall, 212, 1925. None so blind as those who shut their eyes, on acorn cups. A new, 1606. Honey-producing, 1942. Nonsense about the Phylloxera, 1457. Leafy, 774. North America. Dilar in, 2023. OAK. INSECTS AFFECTING. Early references to the Hes-Amphibolips q.-inanis, 1448. sian-fly in, 2267. q.-prunus, 1606. Notes on Psyllids of, 2272. q.-spongifica, 1448. Packard, A. S. Zoo-geograph-Andricus q.-californicus, 1967. ical map of, 2267. q.-punctatus, 1822. American Anthomyidæ, 1966. seminator, 739, 1037. Coleoptera. Classification of, Balaninus rectus, 1311. 2085. Biorhiza forticornis, 745. Hymenoptera. Descriptions of, Cecidomyia q.-pillulæ, 799. q. symmetrica, 799. 385. Lepidoptera. New lists of, 2132. Cecidomyidæ, 799. Microgasters. Notes on, 1960. Cerambycid, 2267. microlepidoptera. Works on, Ceroptres ficus, 745. Coccus, 159. Entomological ignorance in the, 431. Cossus robiniæ, 1003. Northern army-worm, 1835. Cynipid, 212. Cynipidæ, 41, 799. brenthian, 1363. Illinois. Six worst enemies of fruit-Cynips fecundatrix, 1606. growers in, 377. forticornis = Biorhiza forticornis. lady-bird and its larvæ, 1289. gallæ-tinctoriæ, 1560. Northwest in 1883. Bruner, L. Rocky Mountain q.-californicus = Andricus q.-califorlocust in the, 2277. nicus. on the Rocky Mountain locust. Bruq.-decidua, 1925. ner, L. Report of observations in q.-frondosa, 774, 1606. the, 2165. q. glandulus, 1606. Note, 184, 954. q.·inanis = Amphibolips q.·inanis. Notes. Entomological, 2, 1396, 1554, 1609, 1610, q.-mellaria, 1942. q. prunus = Amphibolips q. prunus. 1681, 1958, 2015, 2058, 2066, 2135, 2147, 2168, 2184, 2192, 2219, 2228, 2247, 2261. q.-punctata = Andricus q.-punctatus. by B. D. Walsh, 174. q.-spongifica = Amphibolips q.-spongi-Nothrus ovivorus, 1357. fice Notices, 196. seminator = Andricus seminator. Notodonta concinna = Œdemasia concinna. вр., 799. unicornis = Cœlodasys unicornis. terminalis, 1448. Nottiglossa undata = Oncometopia undata. Edema albifrons, 422. Noxious insects increase upon us. Why, 766. Elaphidion parallelum = E. villosum. named, 454. putator = E. villosum. Poisoning, 205. villosum, 288, 793, 2267. Popular remedies for, 73. Euclemsis bassetella, 1972. Nozzle. Cyclone, 2327. Galla, 1269. Nuisance. Caterpillar, 1995. Holocera glanduella, 1310, 1311. made useful, 983. Ithycerus noveboracensis, 1033.

Kermes galliformis, 1972.

Lachnosterna fusca, 1812.

pilosicollis = L. tristis.

quercina = L. fusca.

tristis, 966.

Number of entomologists in Europe, 1865.

fluenced by food, 2167.

sects. Packard, A. S., 2267.

molts and length of larval life as in-

segments in the head of winged in-

OAK. INSECTS AFFECTING-Continued. Œdipodini, 1959. Lithocolletis cincinnatiella, 1879. Enothera grandiflora. Moths caught by, 1419, Mallodon melanopus, 2291. 1761. Myrmecocystus hortus-deorum, 1942. Œstridæ, 114, 153, 290, 2256, Mytilaspis pomicorticis = M. pomorum. Æstrus hominis = Dermatobia hominis. ovis, 456, 887, 914, 1057, 1059, 1367, 1568, pomorum, 973. Ozognathus cornutus, 1967. 2238. Phylloxera lichtensteini, 1421. Œta compta = Œ. punctella. Tenthredinid, 108. punctella, 1059, 1343. Xylotrechus colonus, 2267. Ohio insects. Alwood, W. B. Report on, 2382. Oil-beetles, 1053, 1643. Oak. Insects stripping the bur, 966. leaf gall, 790. Old-fashioned potato-bugs, 186. Leaf-miners on white, 1879. and new philosophy, 392. Phylloxera, 1363. question of species, 2201. Oleander. Chilocorus bivulnerus on, 730. pruner, 793. Coccid on, 218. and rose galls, 1987. tree borer, 1003. Insects on, 730. caterpillars, 422. Olfersia americana, 2307. twigs. Cynipid galls on, 1822. Omaha. Conference of western governors at, Woolly galls on white, 739. 1557. OATS. INSECTS AFFECTING. Oncideres amputator, 1856. cingulata, 442, 443, 476, 489, 746, 848, 1566, Aphia avenæ = Nectarophora granaria. Nectarophora granaria, 1806. 1938. Silvanus surinamensis, 12:9. putator, 1556. Oats and rye. Beetles working in wheat, 1259. Oncometopia undata, 36, 79, 164, 399, 1013. One day's journal of a state entomologist, 383. Oberea bimaculata, 302, 783, 1363. perspicillata = 0. bimaculata. effect of the Mississippi floods, 2088. half the vine area of France affected by Phylвр., 1059. loxera, 2020. f boring in apple twigs, 1860. tripunctata, 1503. Onion-fly, 225. Obituary. Bazille, L., 2370. ONION. INSECTS AFFECTING. Belfrage, G. W., 2161. Anthomyia ceparum, 225. Chambers, V. T., 2218. Cambala annulata, 236. Clemers, B., 223. Iulus multistriatus - Cambala annulata. Glover, T., 2218, 2258. Ortalis flexa = Tritoxa flexa. Tritoxa flexa, 225. Lichtenstein, J., 2370. Maeklin, F. W., 2161. Onion-maggot. Means against the, 12. pest. Chester, 2319. Mniszech, G. V., 2054. Putnam, J. D., 2054. Ontario. Index to the entomological reports of Oblong winged katydid, 1329, 1363. the province of, 2228. Eggs of the, 569, 1157. Onward march of the Colorade potato-beetle, Ocellate leaf gall of the red-maple, 2119. Odonata. Venation of, 39. Ophiogomphus mainensis n. sp., 39, Odontota, 2267. Ophion. Long-tailed, 1311. dorsalis, 451. macrurum, 1112, 1802. rubra, 1849, 1876. pargatum, 1442, 1670. Supposed eggs of, 1876. Purged, 1127. Odor in butterflies, 2025. Opsiccetus personatus, 497. Odynerus, 770, 1827. Orange. Bark-louse on osage, 289. birefimaculatus, 1827. in Brazil. Branner, J. C., insects injuriflavipes, 770, 1827. ous to the, 2277. Œcanthus, 2267. California. Scale-insects of the, 2389. latipennis n. sp., 2026, 2276. Experiments upon scale-insects affectniveus, 143, 251, 286, 414, 554, 728, 958, ing, 2164. 961, 999, 1006, 1059, 1122, 1323, in Florida. Voyle, J., the effects of cold 1329, 1333, 1503, 1691, 2195, on the scale-insects of the, 2277, 2238, 2276, Hubbard, H. G. Rust of the, 2291. Habits of, 207. ORANGE. INSECTS AFFECTING. Œcodoma fervens = Atta fervens. Coccid, 218. Œdemasia concinna 61, 413, 454, 779, 788, 922, Mite, 2291. Phytopus oleivorus, 2308. Œdipoda atrox, 1959. Orange insects. Californian, 2373. carolina = Dissosteira carolina. and cotton-worm, 2067. cruciata, 2075. Kerosene as a means against, obliterata, 1969. 2164. pellucida, 1959. Miscellaneous notes on, 2164.

Orange insects. Successful management of the most destructive, 2088.

> for the mulberry silk-worm. Osage, 1220, 1286.

Rust of the, 2308.

Scale-insects of the, 2119.

seed. Worms in osage, 507.

Silk-worms fed with osage, 1341, 1542.

Successful management of insects most destructive to the, 2088.

trees. Insects found about, 1798.

Orchard giving out. An, 578.

Injured, 1594.

Scolytus. Imported, 2233.

Heughton's insects in the, 188.

Orchelimum, 118, 861.

glaberrimum, 1329.

gracile, 447.

Oregon. Apple-plant lice in, 1899. Organisms. Interaction of, 139, 1954.

Orgy1a, 872.

antiqua, 1797.

Glands in, 2412.

leucostigma, 65, 480, 535, 600, 762, 1000, 1059, 1151, 1227, 1352, 2198, 2238, 2378, 2379, 2380. moth issuing from a larva, 1779.

Variable moulting in, 2380.

Ori atidæ, 1363.

Original habitat of Icerya purchasi, 2415. Orocharis saltator, 513, 1287, 1323, 1329, 2276.

Ortalis flexa = Tritoxa flexa.

Orthocentrus pusillus, 385.

stigmaticus n. sp., 385.

trifasciatus n. sp., 385.

Orthoptera, 1329, 2267, 2276.

Orthoptera, Packard, A. S. systematic position of the, 2267.

Orthopterous insects. Packard, A. S., embryological development of, 2267.

Orthosoma brunneum, 95, 397, 2267.

Cylindrical, 1059.

cylindricum = 0. brunneum.

unicolor = 0. brunneum.

Oeage orange. Bark-louse in. 389.

OBAGE ORANGE. INSECTS AFFECTING.

Dipteron, 597.

Lecanium acericola = Pulvinaria innumerabilia.

> maclure = Pulvinaria innumerabilia.

Pulvinaria innumerabilis, 389.

Sericaria mori, 1341, 1542, 1609, 2234.

Sphinx hageni, 2119.

Osage orange for the mulberry silk-worm, 1220, 1286.

seed. Worms in, 597.

Silk-worms fed with, 1341, 1542.

Sphinx, 2119.

vs. mulberry for the silk-worm, 2234.

Osborn, H., report of experiments at Ames, Iowa, 2344.

> upon the insects of the season in Iowa, 2418.

Oscinia, 2394.

brassics: n. sp., 2291.

Oscinis. Cabbage, 2291.

Other insects affecting cheese, 1703.

Otiorhynchidæ injurious to cultivated plants. 2117.

Otiorhynchus picipes, 1788.

Our bugs, 2374.

table. On, 395, 408, 506, 546, 568, 623, 827, 1110.

Out of evil cometh good, 618.

Outbreaks of army-worms. Recent, 2262.

Overcrowded, 654.

Overflow bugs in California, 2099.

Oviposition in Carabidæ. Mode of, 2341.

Prodoxus decipions, 2049.

round-headed apple-tree borer, 2266.

Saperda bivittata, 1621. Tortricids, 1922.

the Yucca moth, 1354.

Owlet moth. Spider-wort, 1301.

Owls' pellets, 601.

Ox insects, 290.

Oxalis. Aleurodes on, 1791.

Ox-eye daisy as an insecticide, 1861.

Oxyopes viridans, 2343.

Oxyptilus periscelidactylus, 1059, 1175, 1301.

Oyster shell bark-louse, 373, 1059, 1127.

of the apple, 1329, 2238.

in Missouri, 1169.

Ozognathus cornutus, 1967.

Pachylobius picivorus, 845.

Pachypsylia n. g., 2208, 2272.

c.·gemma n. sp., 2272.

c.-mamma n. sp., 2208, 2272. venusta, 2208, 2272.

Pachytylus migratorius, 1959, 2044. Pacific coast. Entomologist for the, 2022.

Wheat-stalk worm on the, 2123.

Pack your insects. Mind how you, 435.

Packard, A.S. Additions to the third report on

the causes of the destruction of the evergreen and other forest trees in northern New England, 2364.

> Causes of destruction of evergreen forests in New England and New York, 2232.

> Descriptions of larvæ of injurious forest insects, 2267.

> Development of the bark-boring beetles Hylurgops and Xyleborus, 2267.

Embryological development of Caloptenus atlanis, 2267.

Embryological development of Caloptenus spretus, 2267.

Embryological development of the locust, 2267.

Embryological development of orthopterous insects, 3267.

Fourth report on insects injuring forest and shade trees, 2382.

The Hessian fly, 2267.

Narrative of the first journey in

1877, 1643.

a second journey in

1877, 1643.

Notes on forest-tree insects, 2253.

Sheep, 2043.

Packard, A.S. Notes of a journey to Utah and | Parasites, 66. Idaho in 1878, 1959. on bees, 1500. Number of segments in the head bred from cotton-worm, 1722. of winged insects, 2267. of Colorado potato-beetle. Mite, 1505. Second report on the causes of cotton-worm, 1712. the destruction of the evercut-worms, 349. Efficacy of Chalcid egg, 2115. green and other forest trees in northern New England and on eggs of Caloptenus spretus, 161 New York, 2291. fowls, 1308. Systematic position of the Orthe hateful grasshopper, 728. thoptera in relation to the other Hessian-fly, 2332. orders of insects, 2267. .Hog-caterpillars of the vine infested Third report on the causes of the with, 1247. destruction of the evergreen of the human animal, 728. and other forest trees in north-Importing European, 208. ern New England, 2363. on its larva. The Abbot Sphinx, 1277. Packard's Guide to the study of insects (Review), of the larva of Lachnosterna fusca, 2391. 395, 479, 827. Leucania unipuncta, 6, 10, 17, 1670. Pædisca celtisana n. sp., 1968. of the plum Curculio, 1751, 1795. giganteana n. sp., 1968. Synonyms of, 1932. obfuscata. Girdling habits of, 2405. Parasitic coccons, 158, 1125, 1134. scudderiana, 2356. Not eggs but, 1290. Notes on, 2213. coleoptera, 2353. Painted borer, 89. diptera, 2076. lady-bird, 1329. flies, 310. fungus. Remarkable, 1313. Pale thighed tortoise beetle, 1127. Paleacrita n. g., 1438. Parasitized tomato-worm, 155. vernata, 75, 86, 172, 173, 377, 1021, 1066, Paria aterrima, 1887, 1904, 2229. Paris green absorbed † Is, 1487. 1127, 1357, 1363, 1423, 1438, 1482, 1539, 1604, 1847, 2012, 2149, 2238, 2267. for cotton-worms. How to use, 1544. Palingenia bilineata = Hexagenia bilineata. the curculio, 1258. its effects on plants, soil, and man, flavescens n. sp., 24. puichella n. sp., 24. 1427. terminata n. sp., 24. as an insect destroyer, 1447. vittigera n. sp., 24. insecticide. Use of, 1497. Paris green, London purple and, 2021. Pamphila ethlius, 1897. Paniscus geminatus, 1059. purple, 2021. Pangus caliginosus = Harpalus caliginosus. Parorgyia. Glands in, 2412. Panorpa, 2267. Synonyms and food-plants, 2412. Panorpidæ, 2267. Parsnip caterpillars, 453, 804. Paper makers, 505. PARSNIP. INSECT AFFECTING. Paphia glycerium, 1117, 1127, 1329, 1704, 1711. Papilio asterias, 453, 804, 863, 1273. Papilio, 1352. Parenip worm, 863. asterias, 453, 804, 863, 1272, 1276, 1512. Parthenogenesis of the Hessian-fly. Probable, cresphontes, 1237, 1300, 1914. 1787 glaucus, 28, 1048. in insects Occurrence of, 1029. marcellus, 666. of Mytilaspis pomicorticis, 1956. philenor, 1127, 1321, 1774, 1957. Passalus cornutus, 1311, 1329, 1636. thoas = P. cresphontes. Horned, 1311. troilus, 469, 857. Egg of, 1329. turnus, 28, 361, 1048. Passer domesticus as an insect destroyer, 1667. Parandra brunnea, 1733, 1876. PRA. INSECTS AFFECTING. Paraphia. Fir, 2363. Bruchus pisi, 434, 1120, 1301, 1916, 2238. Pararhyssa n.g., 385 = Rhyssa. Cerotoma caminea, 2418. Parasa chloris, 1233. Gryllus sp., 2383. Parasite of cabbage-worm, 2221. Pea and its insect foes, 14. canker-worm, 717. weevil, 1301, 2238. the imported cabbage-worm. Peach borer, 1017, 1059. Introduction of a, 2291. Flat-headed, 2267. Imported current-worm fly and its, 380, borers, 871, 1366. 381. Apple and, 1475, 1513. Mud-wasp and, 1827. Means against, 121. PEACH. INSECTS AFFECTING, 514. Phora not a, 1923. Ægeria exitiosa = Sannina exitiosa. on Prodoxus decipiens, 1831. or a scavenger! Is Cyrtoneura a, 2102. Asilus sp., 660.

Callimorpha lecontei, 1301-

PEACH. INSECTS AFFECTING-Continued. PEAR. INSECTS AFFECTING-Continued. Carpocapsa pomonella, 1334. Eriocampa cerasi, 1031, 1382. Cetonia inda == Euphoria inda. Euphoria melancholica, 372. Dicerca divaricata, 871, 2267. Euryomia melancholica = Euphoria melan-Ephestia interpunctella, 325. cholics. see = E. interpunctella. Ips fasciatus, 1239. Eriocampa cerasi, 1258. Lygus pratensis, 2, 31. Euphoria inda, 447. Mytilaspis pomicorticis = M. pomorum. Gortyna nitela, 632, 1210. pomorum, 973, 982. Helops æreus, 751. Notodonta concinna = Œdemasia concinna. , pullus = H. æreus. Œdemasia concinna, 779. Larva, 661. Oncideres cingulata, 848. Lepidopteron, 697, 1210. Platycerus querous, 1162. Lithophane antennata, 1182. Polycaon confertus, 2103. Mycetophila persicæ, 653, 660. Prionus laticollis, 561. sp., 419. Rhopalus sp., 372. Œcanthus niveus, 554. Scolytus rugulosus, 2233. Phlœotribus liminaris, 2047. Selandria cerasi = Eriocampa cerasi. Sannina exitiosa, 587, 617, 871, 1017, 1059, 1070, Tremex columba, 928. 1475, 1513, 2238. Typhlodromus pyri, 1739, 1759. Scolytus rugulosus, 1940. Pear leaf blister, 1739. Selandria cerasi = Eriocampa cerasi. midge, 2363. Thyridopteryx ephemeræformis, 1189. root-borer, 561. Xylina cinerea = Lithophane antennata. shoots. Bug gathering on, 741. Peach. Large green worm in a, 661. alug. Blood-sucker and, 1253. tree bark borer, 2047. Current worm and, 1031. borer, 1070, 2238. tree borer. Elm and, 928. borers, 871. insect. A new, 1162. trees again. , Mounding, 617, 653. msects, 277. Beetle, around, 751. Rows of eggs in, 577. Insects around, 660. alug, 1382. Minute borers in, 1940. worms, 779. Mounding, 587. twigs. Girdled, 848. Sulphur cure for, 176. Two new foes of the apple and, 22. Supposed cause of yellows in, 515, Pearl wood nymph, 1127, 1301, 1363. twig borer, 632, 697. Pears. Curculios on, 1358. twigs. Eggs in, 554. Peas. Buggy, 1916. worm, 825. from bugs. To keep seed, 434. Pecan trees girdled by Oncideres cingulata. Blue spangled, 1301. boring into, 1182. Young, 489. Pear. Apple-bark lice on, 973. Pediculina, 258, 497. Banded Ips in calyx of, 1289. Pediculus capitis, 497. Bark-lice on the, 962. cervicalis = P. capitis. and cherry trees. Slug on, 1222. humanus = P. vestimenti. Diplosis, 2363. pubis = Pthirius pubis. vestimenti, 497. PEAR. INSECTS AFFECTING. Pedigree moth breeding, 2381. Anametis grisea, 2117. Pelargonium. Gall on, 1764. Anthonomus quadrigibbus, 1358. Pelidnota punctata, 40, 113, 129, 354, 358, 725, 1221, Aragnomus griseus, 2428. 130L Aspidiotus harrisli = Chionaspis furfurus. Spotted, 1301. Capsus oblineatus — Lygus pratensis. Carabid, 692. Pelopæus cementarius, 543, 1371. Habits of Polistes and, 1371. Carpocapsa pomonella, 1334. lunatus = P. cementarius. Cerambycid, 288. Pempelia hammondi n. sp., 810, 1311, 1322. Chionaspis furfurus, 372. lignosella, 2119. Chloroneura malefica = Emposeca viridescens. Pemphigins. Biological notes on gall-making, maligna = Emposeca obtusa. Clisiocampa americana, 863 1653, 1678. Conotrachelus cratægi, 1358. Pemphigini, 2017. nenuphar, 1358. Pemphigus acerifolii n. sp., 1678. Corimelæna pulicaria, 741. formicarius n. sp, 27. Diplosis nigra = D. pyrivora. formicetorum n. sp., 27. pyrivora, 2363, 2392. fraxinifolii n. sp., 1678, 2361. Dolerus unicolor, 1989. imbricator, 449, 1396, 2361. Empoasca obtusa, 22. populicaulis, 446, 713.

populi-monilis n.sp., 1678.

viridescens, 22.

Pemphigus populi-ramulorum n. sp., 1678. Pesomachus, 158. populi-transversus n. sp., 1678. Diminished, 1127. pyri = Schizoneura lanigera. minimus n. sp., 12, 1442, 1670. rhois, 518. Pezotettix, some species dimorphic forms of Cs tessellata, 1948. optenus, 1889. ulmifusus n. sp., 518. pacifica n. sp., 1959. vagabundus, 446, 518. Phakellura nitidalis — Eudioptis nitidalis. Phalæna zea, 2343. vitifolim = Phylloxers vastatrix. Penetrating organ in a gamasid mite. On an ex-Phalenide. Description of a new genus of HR tensile, 1626. Phalangidæ, 838. Pennsylvania ground beetle, 1059. Phaneroptera curvicauda - Scudderia curvicauda Phasmida, 2267. soldier beetle, 1059. Phasmomantis, 457, 590, 859, 971, 979, 1659, 166. Pentagenia n. g., 39. quadripunctata n. sp., 39. Pentarthron n.g. (See p. 379) = Trichogramma. Phongodes, 1874, 3397. Penthina fullerea n.sp. (See p. 378) = P. hebelaticollis, 2412. and Zarhipis. Further notes on, 2411. vitivorana == Eudemis botrana. 2412 Pepsis formosa, 466, 521, 543, 823. Notes on, 2408. Mygale hentzii and, 1619. Phengodina. On the luminous larviform females Periclistus sylvestris, 1131. in the, 2397, 2402. Perilitus indagator, 1311. Phenomena. Vegetable, 436. Pheocyma. Pine, 2363. Perimegatoma variegatum, 2109. Periodical Cicada, 474, 527, 619, 1059, 1811, 1489, 1624, Phemonone 5-caudata, 2410. 1836, 2014, 2312, 2314, 2315, 2363. Philampelus achemon, 1791, 1127. pandorus, 1102, 1127. alias the seventeen-year locust, satellitia == P. pandorus. 1971, 1979. alias the seventeen-year and Philenor swallow-tail, 1127. thirteen-year locust, 1159. Philonthus, 1311. Belated individuals of the 648. Philosophy of the movement of the Bocky Mountain locust, 1669. in Massachusetts, 2321. Old and new, 392. Notes on the, 2318. of the pupation of butterflies, 1711. Our first brood established, 707. Phlœotribus liminaris, 2047. Premature appearance of the, Phlox. Heliothis phloxiphaga injuring, 936, 945. 2320, 2326, worm, 945. in savin to igs. Eggs of the, 698. Phoberia atomaria, 1976, 2343. Song notes of the, 2334. Phobetron hyalinum, 40. Periodical Cicada in southeastern Indiana. Butler A. W., 2364. pithecium, 777, 1272. Massachusetts, tetradactylus, 40. Phora aletim, 1923, 1932, 2102, 2343. a scavenger, not a parasite, 1923. Twigs punctured by, 1055. Phorodon humuli, 285, 279, 1001, 2396, 2394, 2396. or seventeen-year Cicada, 2312, 2314. 2400. Perla decipiens n. sp., 24. dorsata, 903. Phorostoma latum, 2260. Phosphorescence of fire-flies. Intermittence of elongata n. sp., 24. the, 1905. flavescens n. sp., 24. fumipends n. sp., 24 = Chloroperla fumithe glow-worm. Nature of the, 1810. producta n. sp., 24. Phosphorescent larvse. Large, 1874. varians n. sp., 24. Photinus. Food of Calopteron and, 2358. Perlidse, 2267. pyralis, 396, 1097, 1706, 1917, 2358. Perny silk-worm, 1311. Photuris pennsylvanica, 1097, 1705, 1917. Persian insect powder, 1485. Phoxopteris cornifoliana n. sp., 1968. PERSIMMON. INSECTS AFFECTING. fragarize, 499, 984, 1959, 1574, 2324. Ceratocampa regalis = Citheronia regalis. murtfeldtians n. sp., 1968. Citheronia regalis, 1275. nubeculana, 1721. Pests. Carpet, 1663. Phryganeidse, 1188. Drug-store, 510. Phyciodes nycteis, 1410, Museum, 246, 462, 572. Phycita nebulella n.v., 1811 = var. of Acrobasis Potato, 1554. indiginella. of the strawberry, 2324. Phycita nebulo n. sp , 2 = Acrobasis indiginella. West. Winged, 2313. Phyllobrotica longicornis = Diabrotica longi-Petroleum as insecticides. Emulsions of, 2134. and their value as insecticides. Emul-Phylloptera oblongifolia = Amblycorypha oblon-

gifolia.

sions of, 2126, 2200.

Phyllotreta albionica, 2291. Phylloxera vastatrix, 127, 373, 565, 724, 840, 916, striolata = P. vittata. 1281, 1295, 1301, 1311, 1325, vittata, 348, 564, 2291. 1829, 1842, 1353, 1363, 1373, zimmermanni, 2291. 1376, 1394, 1463, 1421, 1423, Phylloxera, 1355, 1396, 1404, 1406, 1428, 1449, 1644. 1435, 1439, 1449, 1454, 1457, 1671. 1469, 1482, 1586, 1531, 1564, in California, 2140. 1623, 1671, 1716, 1721, 1727, Grape, 1727. 1792, 1808, 1841, 1900, 1906, Winged, 1895. 1907, 1952, 2015, 2019, 2020, caryse-avellana n. sp., 1901. 2140, 2150, 2208, 2250, 2283, -caulis, 360, 685, 1439. 2855. -fallax n. sp., 1423. Dipterous enemies of, 2150. -folise, 1311, 1863. Impregnated egg of, 1974. vitifoliæ = P. vastatrix. -globuli n. sp., 27, 300. ·gummosa, n. sp., 1423. work, 1900. ren n. sp., 1423. Phyloptera, 2267. -scissa n. sp., 1901. Phyloscia, 1900. -semen, 373. Phymata erosa, 778, 1660. on chestnut, 353. Physianthus albens, 1418, 1419. congress in Spain, 1907. Capture of meths by, 1388, convention at Berne, 2147. 1419, 1761. correction, 1355. Peculiarities of, 1418. enemy. Grape-leaf, 1530. Physianthus. Insects affecting, 1388. in France. Grape, 2193. Deilephila lineata, 1419. with fungus disease. Infesting, 1808. Noctuidæ, 1419, 1761. in Geelong, Victoria. Grape, 1732. Sphingidæ, 1418, 1419, 1761. Grape, 1329, 1863, 1373, 1423, 1482, Physicians. Quacks or, 282. 1721. Physocnemum brevilineum, 696. and grape-rot, 1623, 1628. Physonota quinquepunctata n. sp. 767 = P. uni-Half the vine area of France affected punctata. by, 2020. unipunctata, 767. laws. 2019. Phytocoridæ, 1301. in Europe, 2248. Phytolacca not fed on by Aletia, 2943. German, 2432. Phytonomus nigrirostris, 2855. lichtensteni, 1421. opimus = P. punctatus. mite, 1863. punctatus, 2010, 2028, 2061, 2119, 2355. Further notes on, 2061. Mold and, 1807. Nervous system and salivary glands rumicia, 2119. of. 1687. Phytophagic species, 45, 50. New biological facts regarding the Phytophagic varieties, 45, 50. grape, 1421. Phytopus, 1739. hickory galls made by, 1901. oleivorus, 2306. Newest facts of grape, 1404. Pickle-worm, 1127. Nonsense about the, 1457. Pieridæ, 613, 1801. not at the Cape. Grape, 1841. Dimorphic, 28. permanently destructive. The Pieris menapia, 2121. grape, 1907. monuste, 1991, 2232. notes, 1406. oleracea, 835, 1127, 2282, 2236. Notes on the grape, 1952, 2291. protodice, 836, 1127, 2282, 2288. in the natural history of, 1439. rapse, 1127, 1257, 1279, 1204, 1469, 1694, 1714, Nouveau remède contre, 2283. 1720, 1721, 1867, 1948, 2221, 2282, 2288, Oak, 1363. 2251, 229<u>8</u>. One-half the vine area of France afinto Alabama. Spread of, 1720. vernalis, 1684. fected by, 2020. Piesma cinerea, 398, 2048. querous, 1363. Remedies for, 1408. Pigeon Tremex, 458. riloyi, 1363, 1428, 1530. in apple, 1126. in sandy soil, 2250. Pigs. Lice on, 900. Pimpla annulicornia, 2343. Some recent discoveries in reference to, 2209. annulipes, 1884. atrata = Thaloesa atrata. epinosa, 2209. in Switzerland. Extermination and cœlebs n. sp., 385. restriction of, 2207. conquisitor, 2343. Use of guano for the grape, 1887. inquisitor, 1059.

Pimpla investigatrix n. sp., 385., Plant-feeding habit of Feniseca tarquinius, 200. pictipes n. sp., 385. habits of predaceous beetles, 1988. pleurivinctus = P. conquisitor. growing out of an insect, 332, 478. Red-legged, 1329. lice, 23, 80, 265. sp. on Parandra brunnea, 1876. again, 1008. Apple tree, 270, 335, 591, 996. vidua n. sp., 385. Pine Amorbia, 2363. on berberry, 644. borer. Common longicorn, 2291. Cherry-tree, 1067. Lesser, 2267. Current, 304. Caterpillars on the, 927. To destroy, 1148. and cedar. Longicorns in, 319. Destroying cherry, 1207. False caterpillars on the, 965. enemies, 151. PINE. INSECTS AFFECTING. Evergreen, 1039. Amorbia, 2363. Mites mistaken for, 59. Asemum mæstum, 2267. from one plant to another. Migratisa Aspidiotus pinifolii — Chionaspis pinifolii. of. 2017. Borers, 1049. in Oregon. Apple-tree, 1899. Callidium antennatum, 319. Plum-tree, 678. janthinum, 319. on potatoes, 1679. Scab in apple us. apple-tree, 1139. Chionaspis pinifolii, 97, 1329, 2079. Cybocephalus nigritulus, 2079. and their foes. Cherry, 1251. Hylobius confusus, 1168. their friends and foes, 198. Lachnus strobi, 320. louse again. Eggs of the apple-tree, 550. Lophyrus abbotii, 465, 927, 956, 1570. Apple, 883. root, 373, 495, 1859. lecontei, 985, 1011, 1570. Monohammus titillator, 1824. tree, 1127. Pheocyma, 2363. Cabbage, 2291. Pieris menapia, 2121. Corn-root, 23. Pissodes strobi, 781, 2363. Current, 322. Podapion gallicola, 2231. eggs on apple and mountain ask Ptinus brunneus, 191. 579. Therina, 2363. Eggs of the apple-tree, 507, 539. Pine-leaf scale insect, 1829. enemy, 301. moth of Nantucket, 2183. Grain, 99, 126, 1127. Pheocyma, 2363. Look out for the eggs of the apple-Therina, 2368. tree, 507. White-pine, 320. trees. Butterfly larvæ injurious to, 2121. killed by borers. White, 1949. Plants. Fertilizing, 324. worm. Leconte's, 1570. Food of insectivorous, 1499. White, 465, 956. and insects. Imported, 1329. Pinion. Ash-gray, 1301, Otiorhynchids injurious to cultivated. Pinus elliotii. Chionaspis pinifolii on, 2079. inops. Gall on, 2231. soil and man. Effects of Paris green on. 1427. strobus. Insects affecting, 97, 320. Piones, Cabbage, 2232. Platamodes unicolor, 573. rimosalis, 1867, 1915, 2232. Platycerus quercus, 1162. Piophila casei, 1141, 1266, 1700, 1734. Platygaster herricki, 2332. radicum, 1357. Platyhypena scabra = Kypena scabra. Pipisa femoralis, 495. Platynus æruginosus, 1013. radicum n. sp., 495 = P. femoralis. excavatus, 1013. Pirates biguttatus = Rasahus biguttatus. maculicollis, 2099. picipes = Melanolestes picipes. punctiformis, 1798. Pissodes strobi, 781. Platyphyllum concavum = Cyrtophyllus con-PITCHER-PLANT. INSECTS AFFECTING, 1885. Sarcophaga sarracenise, 1385, 1390. Platypsyllus castoris, 2417. Xanthoptera semicrocea, 1885, 1390. as determined by the larva. 874 Pithy blackberry gall, 1131. tematic relations of, 2417. galls on blackberry-twigs, 610. Platyptera, 2267. Pityophthorus micrographia, 295. Platyptilus carduidactylus, 1059. minutissimus, 295. Plectrodes scalator, 1330, 1525. Plagionotus speciosus, 547, 915. Plenty of specimens. Send, 593. Plagues. Insect, 2198. Pleurotropis phyllotretæ n. sp., 2291. Planipennia, 2267. Plug-ugly theory, 233, 1383. Plant-bug. Dotted legged, 1311. Plum bark-louse, 107. Curculio, 373, 1059, 1129, 1301, 2296. Tarnished, 1127, 1219, 2291.

breeds in the apple, 1214.

feeders. Carnivorous propensity of, 1867.

INDEX.

Plum Curculio, natural history and how to catch Podosesia syringæ, 844. him, 1372. Poduræ, 1764. Parasite for the, 1751, 1795. Podura nivicola, 608. gouger, 33, 34, 373, 1301. Poduridæ, 317. gum. Larvæ in, 419. Pœcilloptera prumosa, 737, 1329. PLUM. INSECTS AFFECTING. Pœcilocapsus lineatus, 715. Acarid, 1265. Poison. Potato pest, 1524. Ægeria exitiosa - Sannina exitiosa. Poisoning noxious insects, 205. Anthonomus prunicida = Coccotorus scutel-Poisonous Arachnida, 2399. Is the Colorado potato-beetle, 1483. flour, 586. Aphie cerasi = Mysus cerasi. prunifoliæ, 678. Hexapoda, 2399. Attacus cecropia, 1249. insects, 2399. Bibio albipennia, 683. Myriapoda, 2399. Carpocapsa pomonella, 1834. qualities of the Colorado potato-beetle, Cerambycid, 559. Coccotorus scutellaris, 33, 373, 1301, 1378. qualities of the Colorado potato-bug, Colaspis sp., 372. Conotrachelus geminatus, 34. spider, 2306. nenupher, 83, 372, 373, 501, 566, Tomato-worm not, 771. 1059, 1071, 1301, 1372, 1477, 1685, worm, 710. Poisons to destroy insects. Use of, 1887. puncticollis n. sp., 34 = C. Polistas fuscatus = P. metricus. geminatus. metricus, 1371. Dipteron, 419. and Pelopseus. Habits of, 1371, Elaphidion parallelum = E. villosum. rubiginosus, 543. villosum, 606. Pollenia rudia, 2128, 2174. Elaterid, 81. Pollination of Yucca, 1603, 2000. Grapholitha prunivora, 373. Polycaon confertus. Habits of, 2103. Hylobius stupidus = Pachylobius picivorus. Polydesmus, 261. Lecanium sp., 107. complanatus, 215. Lyda sp., 656. serratus, 842. Mytilaspis pomicorticis = M. pomorum. вр., 834. pomorum, 973. virginiensis, 219. Myzus cerasi, 678. Polymitarcys alba, 1926, 1961. Pachylobius picivorus, 845. Polyphemus moth, 542. Phorodon humuli, 2393, 2394. Caterpillar of, 765. Sannina exitiosa, 81. Cocoon of, 1164. Scolytus rugulosus, 1940. silk-worm, 1311. Semasia prunivora = Grapholitha prunivora. Polyrhabdus n. g., 385 = Choringeus. Teles polyphemus, 765. cariniger n. sp., 385 = ChorinsonsPlum leaf worms, 656. cariniger. moth, 373. Polysphincta nigriceps n. sp., 385. saw-fly, 2291. nigrita n. ep., 385. pimploides n. sp., 885. tree insects, 683. Minute borers in, 1940. Pomace-fly. Pretty, 2119. plant lice, 678. Vine-loving, 2119. twigs. Borer in, 559, 606. Pomphopœa senea, 289, 301, 1056. Plume. Grape vine, 1059, 1301. sayi, 188, 301. moth. Spruce, 2363. Synoptic table of, 289. Plums and cotton, 1477. tarsalis, 289. from Curculio. To protect, 1071. Pomphilus formosus = Pepsis formosa. for the million, 501. Poplar borer, 71. Preventing rot in, 1685. POPLAR. INSECTS AFFECTING. Acronycta acericola = A. americana. Wier, D. B. Native, 2388. Plusia brassicæ n. sp., 1127, 1867, 2232, 2238. americana, 575. Cabbage, 1127, 2232, 2238. Saperda calcarata, 1849. dyaus, 2348. Poplars. Caterpillars on Lombardy, 571. Plutella. Cabbage, 1291, 2232. Popular delusion, 525. cruciferarum, 614, 1291, 1297, 2232. names, 100, 430, Pod-like willow gall, 1170. remedies for noxious insects, 73. Podabrus modestus, 358, 673. Populus monilifera. Insects on, 1459. rugulosus, 353. Porcelio, 1900. Podapion n. g., 2231. Porizon conotracheli n. sp., 1751 =Thersilochus gallicola n. sp., 2231. conotracheli.

Curculio parasite, 1301.

Podisus spinosus, 372, 456, 1378, 1431, 1450, 1672.

Potamanthus ! odonatus ». sp., 24. Potato beetle, 946, 968, 995, 1019, 1027. abroad. The Colorado, 1364, 1407. again. Colorado, 1301. Colorado, 21, 334, 991, 1059, 1311, 1329, 1363, 1423, 1431, 1558, 1570, 2291, 2294 2418. enemies, 313, 366. in Europe. Colorado, 1507. New Hampshire. Colorado, 1859. Experience with, 1519. Inquiries concerning, 1491 Insect enemies of the Colorado, 411. Mite parasites of the Colorado, 1505. in New York. Colorado, 1379. The onward march of the, 1218. poisonous? Is the Colorado, 1483. Poisonous qualities of the Colorado, 1425. progress, 1480. Specific for the Colorado, 1490. Sweet, 1697. Ten-lined, 925. Three-lined, 1059, 1558. beetles, 247, 470. Driving, 1044. Fire cure for, 121. Large black, 1206. native home. Colorado, 1462. Safe remedies for, 1948. Some notes on, 1528. Sweet, 681, 1510. Potato-bug, 742, 905, 1450, 1672. Bogue Colorado, 519. Colorado, 293, 834, 366. in Maine, 119. New, 48, 88, 171, 293. Poisonous qualities of the Colorado, 1101. bugs, 401, 794, 1096. To destroy Colorado, 650. Old-fashioned, 185. and corn insects, 138. Insect foes of the, 1558. POTATO. INSECTS AFFECTING, 767, 1050. Acarina, 1444. Aphia solani, 1679. Baridius trinotata = Trichobaris trinotata. Cassida clavata = Coptonycla clavata. Ceresa bubalus, 2107, 2285. Coptocycla aurichalcea, 1510. clavata, 693, 742, 1558. Crepidodera brevis, 1782. cucumeris, 401, 1066, 1059, 1558. parvula, 1782. Doryphora 10-lineata, 21, 48, 121, 138, 171, 293, 834, 347, 365, 366, 401, 565, 760, 905, 925, 946, 968, 995, 1019, 1027, 1044, 1059, 1218, 1301, 1311, 1329, 1363, 1364, 1379, 1405, 1407, 1423, 1425, 1431, 1450, 1462, 1480, 1482, 1483, 1490, 1491, 1505, 1519, 1524, 1528, 1558, 1570, 1597, 1672, 1859, 2081, 2294. Epicauta cinerea, 401, 1056, 1558. corvina, 1206. pennsylvanica, 38, 401, 1056, 1558. Preying Mantis, 457.

puncticollis, 1381.

POTATO. INSECTS AFFECTING-Continued. Epicauta vittata, 401, 470, 794, 1056, 1230, 1558. Epitrix brevis = Crepidodera brevis. hirtipennis = Crepidodera parvula. Gortyna nitela, 401, 1059, 1558, 2229. Gryllus sp., 2383. Haltica cucumeria = Crepidodera cucumeria. Lema trilineata, 138, 401, 565, 925, 1059, 1558. Lepidopteron, 794. Lytta atrata == Epicanta pennsylvanica. cinerea = Macrobasis unicolor. marginata = Epicauta cinerca. murina = Macrobasis unicolor. vittata = Epicauta vittata. Macrobasis unicolor, 38, 347, 401, 470, 1044, 1558. Melanotus incertus i, 846. Meloidæ, 48, 121, 794, 1059, 1230. Nysius sp., 1317. Protoparce celeus, 1059, 1558. 5-maculata, 401. Sciara sp., 239. Sphinx 5-maculata = Protoparce 5-maculata. Trichobaris trinotata, 401, 1059, 1558, 2172. Meloidse injurious to, 48, 121, 794, 1859. 1230. New insect foe of the, 693. pest poison, 1524. pests, 1558. Scab on, 239. stalk-borer, 1059, 2172, 2229. weevil, 1059, 1558, 2291. worm, 1059, 1329, 1558, Potatoes. Black blister beetles on, 1381. Buffalo tree-hopper injurious to, 2107. Plant-lice ou, 1679. Scabby. 1444. Wire-worms in, 846. Pot-herb butterfly, 1127, 2232, 2238. Potter wasp. 770. Fraternal, 1127. Powder. Manufacture of pyrethrum, 1984. Persian insect, 1485. Powders and their use. Insect, 1692. Practical entomology in reality, 914. Prairie belt. First appearance of the cotton-worm in, 1852. fires and hateful locust, 1453. Prasocuris. Table of species, 156. varipes, 156. Predaceous beetles. Plant-feeding habits of, 1956. Predictions verified, 1494. Premature appearance of the periodical Cicada 2320, 2326, Preparation of Diptera, 2041. Preserving insects, 881, 1057. Directions for collecting and, larvæ, 1300. President. Washington Entomological Society. Address as, 2355. Presidential address to St. Louis Academy of Sciences, 1564, 1629. Pretty pomace-fly. 2119. and unique gall-making tortricid, 2176.

Supposed eggs of, 1002.

Prickly ash larva, 1914.

Papilio cresphontes on, 1914. rose gall, 1194, 1245.

PRIMROSE. INSECTS AFFECTING.

Alaria florida = Rhodophore florida. Rhodophora florida, 1273.

Prionide. Food habita, 1902.

Prionidus cristatus, 262, 491.

Prionotus novenarius = Prionidus cristatus.

Prionus. Eggs of broad-necked, 1329.

imbricornis, 95, 1127,

laticollis, 95, 561, 1059, 1081, 1127, 1329,

2287.

Lesser, 2267.

sp. 736.

Tile horned, 1127.

Prisopus sp., 2267.

Pristiphora grossulariæ n. sp., 140, 176, 189, 364, 772,

sycophanta n. sp., 197.

Probabilities of locust or grasshopper injury in the near future, 2335.

Problem of the hop-plant louse in Europe and America, 2400.

fully solved, 2396.

Proboscis of the common house-fiv. 1783.

Proconia undata = Oncometopia undata.

from stomach of Sturnella magna, 1013.

Procris. American, 1127.

americana = Harrisiana americana. falsarius = Acoloithus falsarius.

Grape vine, 213.

Protacanthus milberti, 1127, 1269, 1543.

missouriensis = P. milberti.

Proctrotrupidæ, 2343.

a new genus of, 1749.

Prodenia autumnalis n. sp., 1301 = Laphygma frugiperds.

commelinæ, 850, 1282.

fulvosa = var. of Laphygma frugiperda. lineatella, 1775.

obscura = var. of Laphygma frugiperda. sp., 2238.

Prodoxidse, 2000.

Prodoxus n. g., 1830, 2000.

ænescens n. sp , 2000.

cinereus n. sp., 2000.

decipiens n. sp., 1804, 1830, 1854, 1983, 1943, 2000, 2049.

> mistaken for Pronuba yuccasella, 1804.

Oviposition of, 2049.

A parasite on, 1831.

Evolution of, 1804.

Further notes on Propuba and, 2000. remarks on the difference be-

tween Pronuba and, 1854.

intermedius n. sp., 2000.

marginatus n. sp., 2000.

Pronuba vs., 1844.

Structural and anatomical peculiarities of. 2171.

Profits of silk-culture, 2338.

Promachus, 1127.

apivorus, 668, 1059, 1543. bastardii, 1269.

Promachus vertebratus, 1269.

Promethea silk-worm, 1311.

Prominent. The red-humped, 2291.

Pronuba n. g., 1829, 1336, 2000.

and its connection with the pollination of Yucca, 2391.

unicorn, 1280.

Evolution of, 1804.

maculata n. sp., 2000.

A new tineid genus allied to, 1830. and Prodoxus. Further notes on, 2000.

> remarks on the differences be

tween, 1854.

Structural and anatomical peculiarities of, 2171.

vs. Prodoxus, 1844.

yuccasella n. sp., 1329, 1336, 1337, 1363, 1603, 1854, 1983, 1943, 2000, 2171.

Chrysalis of, 1414.

Further remarks on, 1603.

Mr. Hulst's observations on, 2871.

mistaken for Prodoxus decipiens, 1804.

Mystery in reference to, 1983. Supplementary notes on, 1337.

Prophecy fulfilled, 709.

Proscopia, 2267.

Prosopis affinis, 1171.

Prospects. Locust, 1527, 1562, 1582. Protection of insect collections, 2180.

Protective device employed by a glaucopid cater-

pillar, 2242. resemblances. Mimicry or, 1301, 1340.

Proteoteras n. g., 1968.

zeculana n. sp., 1968, 2114, 2206.

Protoparce carolina, 131, 933, 1089, 1262, 2238.

celeus, 131, 152, 401, 490, 710, 815, 851, 885, 993, 1059, 1089, 1264, 1329, 1558, 1661.

Prunus insects, 1503.

Psenides, 821.

and inquilines. Relations of, 41.

Psenocerus supernotatus, 481, 1458.

Pseudococcus aceris, 1890.

Pseudoneuroptera, 2267.

of Illinois, 24.

New, 24, 39.

Pseudoscorpion, 1884.

Psinidia wallula n. sp., 1959.

Paoci, 2133.

Psocidæ, 1147, 2267.

Psocina. Structure of the claw in, 2045.

Paocus, 1078.

amabilie n. sp., 24.

bifasciatus n. sp., 39. confluens n. sp., 39.

conterminus n. sp., 39.

geologus n. sp., 24.

lichenatus n. sp., 39.

madescens n. sp., 39.

permadidus n. sp., 39.

perplexus n. sp., 24.

pollutt s n. sp., 24.

Psocus purus n. sp., 24.

Pyrethrum willemoti, 2343.

rufus n. sp., 39. semistriatus n. ep., 24. venosus, 176, 1078, 1147, 1195, 1263, 1526, 1798. Psyche confederata, 1441. Psychidse, 1650. Psychomorpha epimenis, 1301, 1363. Psylla. Box, 2291. rabi, 669. Quedius, 1311. Psyllid galls. Hackberry, 2208. Psyllids of North America. Notes on, 2272. the United States, 2210. Pteromalus puparum, 2221. Pteronarcys, 2267. Pterophorus carduidactylus n. sp., 1059. periscelidactylus = Oxyptilus periscelidactylus. Pthirius pubis, 497. Ptinus brunneus, 191, 516, 1141. Publication. Discontinuance of, 2106. Publilia concava, 163. Puccinia graminia, 1605. Pulex irritans, 497. penetrans - Sarcopsylla penetrans. sp. feeding on lepidopterous larvæ, 2110. Pulvinaria acericola = P. innumerabilis. / innumerabilis, 389, 1515, 1816, 2279, 2291. macluræ = P. innumerabilis. vitis, 106, 1212. Punctured grape-caues, 513, 1333. Punctures of Hemipters upon shrubs, fruits, and grains. Webster, F. M. Experiments on the effects of, 2382. Punctures on rose twig, 1167. Pups of the disippus butterfly, 1193. Pupæ. Habits of, 346. How to hatch, 509. Pupation of butterflies. Philosophy of the, 1711. Nymphalidæ, 1704. Ranatra, 1709. Purged Ophion, 1127. Purple willow Gracilaria, 2363. PURSLANE. INSECT AFFECTING. Deilephila lineata, 1402. Putnam, J. D. Obituary, 2054. Pyralid infesting seed-pods of trumpet-vine, 1929. Pyralis farinalis = Asopia farinalis. olinalis = Asopia olinalis. Pyrameis atalanta, 540. cardui, 1635, 1770. Pyramidal grape-vine worm, 1801. Pyrethrum. Additional experiments with, 1903. cinerariæfolium, 1996, 2235. Cultivation of, 1984. Directions for cultivating, 1996. raising, 1862. Experiments with, 1948, 2353. an important insecticide, 2131. powder. Manufacture of, 1984. roseum, 1996, 2119, 2343. for the screw-worm, 1921. in the United States. Additional notes on the cultivation of, 2377. The use of, 1858.

ite use as an insecticide, 2119.

Pyrophila conspersa, 1301. pyramidea, 785, 1301. pyramidoides, 671, 785, 1301, 1471. Hibernation of, 1471. Pyrrharctia isabella, 244, 558, 909, £153, 1311, 1842. Pyrrhia exprimens, 677. Quackery. Entomological, 439. Quacks and physicians, 282. Queen bees in the mails, 1762. Quelques mots sur le insecticides, 2268. Quercus acuminata. Cynips on, 1822. coccifers. Phylloxers on, 1421. douglassi. Gall on, 1967. Galls on the prinus group of, 1606. infectoria. Gall on, 1560. muhlenbergi. Cynips on, 1925. palustris. Coccid on, 1972. undulata. Galls on, 1942. Queries answered; 217. Bundle of entomological, 1463. Questions answered, 1246. Quick traveler, 752. Quince Curculio, 1301. Quince. Insects affecting. Capsus oblineatus = Lygus pratensis. Conotrachelus cratægi, 1301. Eriocampa cerasi, 2243. Lygus pratensis, 2. Selandria cerasi = Eriocampa cerasi. Quince. Saw-tly on the, 2243. Rabbit insects, 290. Race. Some further facts regarding that fatherless, 1658. That fatherless and motherless, 1650. RAGWEED. INSECTS AFFECTING. Epicauta pennsylvanica, 732. Lytta atrata = Epicauta pennsylvanica. Railroad worm, 2238. fusca, 797, 1191. Range of the rear-horse, 859. Ransom chip-trap for Conotrachelus, 1177. Ranunculus acris. Insecte affecting, 156. Curculio remedy, 1201. Rapacious soldier-bug, 1050. Rape butterfly, 836, 1294. our new cabbage pest, 1257. Rare capture in Illinois, 1271. monstrosities, 2224. Rasahus biguttatus, 497. Rascal leaf-crumpler, 341, 373, 1311, 1580. in Georgia, 1962. Raspberries. Do bees injure, 529. destroyed by weevils, 1788. Raspberry borer, 783. brand, 676. canes dying, 1085. Eggs of snowy tree-cricket in, 1122. tree-cricket in, 1006. Gouty gall on blackberry and, punctured by Orchelimum glaberrimum, 2249.

gouty gall, 1124. and grape-vines. Egg punctures in, RASPBERRY. INSECTS AFFECTING. Ægeria rubi = Bembecia marginata. Agrilus ruficollis, 1124, 177L Bembecia marginata, 1863, 1453. Chionaspie furfurus, 1968. Diaspis harrisii = Chionaspis furfurus. Dipteron, 676. Monophadnus rubi, 663, 761, 1212, 1641. Oberea bimaculata, 783, 1363. perspicillata = 0. bimaculata.tripunctata, 1503. (Ecanthus niveus, 961, 1006, 1122, 1503, 1691, Orchelimum glaberrimum, 2249. Otiorhynchus picipes, 1788. Rhodites radicum, 1149. Selandria rubi = Monophadnus rubi. Synchlora rubiyoraria, 1059. Raspberry-root borer, 1963. gall, 1149. saw-fly, 1212, 1641. Scale insect on, 1968. worms, 663, 761. Rat-tailed larva, 168. Ratseburg. Forest trees and weeds of Germany (revicw), 1110. Rear-horse, 1059. Eggs of the Mantis or, 1060. Range of the, 859. Rear-horses vs. grasshoppers, 590. Record of American entomology (review), 827. Red ants, 2309. cedar caterpillars, 427. clover. Bumble bees and, 2376. humped caterpillar, 788. prominent, 61, 2291. leg. The cussed, 1598. legged ham-beetle, 1363. locust, 1423, 1625, 2368. maple. Occilate leaf-gall of the, 2119. mites, 1959. Transformation of the, 1632. scale. California, 2394. Koebele, A., experiments on the, 2894. shouldered Sinoxylon, 1311. spider, 1146, 1225, 1710. on roses, 1710. tailed Tachina fly, 1127. weevil, 711. Reddish amout beetle on apple. Small, 1244. yellow spruce bud-worm, 2291. Reduvius personatus = Opsicœtus personatus. raptatorum = Sines diadems. Regal walnut caterpillar, 425. Regulating sex in insects, 1415. Remarks, 206, 255. Remedies and appliances, 2375. More universal, 278. Universal, 204, 503, 818. Remedy. Another universal, 232. Report of committee of entomology, Ill. State Hortic. Soc., 1056.

Raspberry Geometer, 1059.

Report of committee of entomology, Mo. State Hortic. Soc., 1113, 1127. experiments, chiefly with kerosene, upon insects injuriously affecting the orange-tree and cotton-plant, 2164. to governor of Kansas on the grasshopper question, 1573. on Lucilia macellaria, 2199. of State entomologist of Illinois, 1, 373, Missouri, 1, 1059. 2, 1127. 3, 1301. 4. 1311. 5, 1329. 6, 1863, 7, 1423. 8, 1482. 9, 1579. Index, 2026. U. S. Ent. Commission, 1, 1643. 2, 1969, 8, 2267. 4, 2343. U. S. Entomologist for 1878, 1721. 1881-'82, 2119. 1883, 2232. 1884, 2291. 1885, 2363. .1886, 2394. 1887, 2418, Reports. Missouri entomological, 1680. U. S. Entomological Commission, 1757. Resemblances. Mimicry or protective, 1301, 1840. Resistance of grape-vines to Phylloxers in sandy Retarded development in a blister-beetle, 1860. of insects, 2040. Remarkable case of, 2003, 2005. Retirement of Mr. Fuller, 1910. Revision of the Lampyridæ, 1819. Rhagium lineatum, 2267. Rhenm rhaponticum. Asilus sericeus feeding on, Rhinoceros beetle. Gigantic, 580, 1216, 1292. Rhinopsylla n. g., 2272. schwarzii n. sp., 2272. Rhipiphorus sp., 2301. Tiphia and, 2851. Rhizotrogus sp., 2260. Rhodites, 1235. bicolor, 1037, 1194, 1245. radicum, 137, 1149. rosse, 570, 1166, 1989. Rhodobsenus 13-punctatus, 1301, 2119. Rhodophora florida, 807, 1278. Rhopalocera, 1635, 1784. Rhopalosiphum, 1678. Rhopalus, 372. Rhus glabra. Insects on, 320. Rhynchophora, 254, 1301. Rhynchophorus simmermanni, 2119. Rhyssa, 385. atrata = Thalessa atrata. not ligalvorous, 2286. lunator = Thalessa lunator.

Rocky Mountain locust. Philosophy of the move Ribes insects, 772. Rice crop. Enemies of the, 278. ments of the, 1669. grub, **2**119. scourge, 1620. RICE. INSECTS AFFECTING. in Wyoming, Montana, etc., in 1881. Bruner, Acridium obscurum, 2119. L., the, 2267. Cecidomyia oryzæ, 1949. Rocky Mountains. Excursion to, 988. Chalepus trachypygus, 273, 1911, 1949, 1973, 2119. Chilo orysmellus, 2119, 2120. Rogas n. sp., 879. Laphygma frugiperda, 2011. sp., 1002. Lissorhoptrus aimplex, 273, 1911, 1973, 2119. Roller. Strawberry-leaf, 984, 1050, 1574. Murmidus ovalis, 2217. Roman-nosed pupa, 1217. Rice. Insects affecting stored, 2190. Rooms. Swarms of minute flies in, 506. Root Aphis, 990, 1038. enemies of growing, 1911. plant. Another enemy of the, 2011. Apple-boter and, 1638, blight. Apple, 997. Insect enemies of the, 1949. Insects affecting, 2119. borer, 981. Water weevil of the, 1973. Raspberry, 1363. stalk borer, 2119. New, 2120. 1187. weevil, 1663. louse. Barley, 2394. Ricinis communis. Acridide feeding on, 1645. Syrphus-fly, 1059. Ridding the ground of cut-worms, 2810. 1.iley to Dr. Schaffer. Entomology. Professor, 2360. in favor of the birds. Professor, 1675. insects, 1149. Letter from Prof. C. V., 1676. Rose. Bedeguar of the, 1939. and the locusts. Professor, 1434. beetle, 361. Yucca moth. Professor, 2068. Riley's report to the governor of Kansas. Professor, 1573. researches, 2007. Ring-banded soldier-bug, 1311. bug, 373, 1075. legged Pimpla, 1329. on apples, 748. Robber. Many-banded, 1059, 1423. Rose bug remedy, 1478. Robin. Cut-worms from the stomach of, 1873. bugs, 1278. Insects from the stomach of, 1926. Rock-bass. Insects from stomach of, 1793. chafer, 1329, 1583. Rocky Mountain grasshopper, 1557. gall, 1235. locust, 1423, 1451, 1452, 1482. 1588, 1557, 1570, 1625, 1648, Mossy, 570, 1166. 1959, 1998, 2267, 2291. Prickly, 1194, 1245. Bruner, L., observations galls. Oak and, 1037. in the Northwest on ROSE, INSECTS AFFECTING. the, 2165. in 1880. Martin, J., on the, 2267. Diaspis ross, 1303. 1885. Bruner, L., report on the abundance Lecanium olese, 1303. of the, 2363. rosse, 1308. Geographical distribution of the, 2267. 1875, 1478, 1583. or grasshopper. Governors of Western States Pyrrhia exprimena, 677. on the, 1557. Rhodites bicolor, 1037. Important observations on, 1575. 1989. Lecture on the, 1493. sp., 1235. in Montana in 1880. Bruner, L., 2267. Natural history of the, Rose-slug, 672, 1780. 1578. and other insects in the worms, 677. Northwest during the Roses. Failure of tea, 1768. summer of 1883. Bruner, L., observations on

the, 2277.

lice. Knots on apple-tree roots canced by, Roots of Ampelopsis. Swellings on, 1428. Insects affecting apple-tree, 963. Rosacese. Diastrophus confined to, 1121. in California. Fuller's, 1749. Fuller's, 1708, 1721, 2291. Habits of Fuller's, 1708. Not Fuller's, 1825. bushes. Bark-lice on, 1303. chafers on grape-vines, 1375. Aramigus fulleri, 1708, 1721, 1749. Cynips bicolor = Rhodites bicolor. Heliothis marginidens = Pyrrhia exprimens Macrodactylus aubspinosus, 361, 373, 1675, 1329, Monostegia rosse, 672, 1780. rose, 570, 1687, 1166, 1194, 1245, Selandria rosæ == Monostegia rosæ. Tetranychus telarius, 1710. twig. Puncture on, 1167. The red spider on, 1710. Rot in plums. Preventing, 1685. Rotten root, 758.

Round-headed apple-tree borer, 1059, 1608, 2238. New facts about the, 1630. Oviposition of the, 2266. Rove beetle. Spotted, 1123. beetles, 1558. Royal horned caterpiller, 702, 803, 800, 1275. Rubus insects. 1503, Rudbeckia. Aphid on, 2205. RUDBECKIA. INSECTS AFFECTING. Nectarophora rudbeckiæ, 2205. Siphonophohra rudbeckiæ = Nectarophora rudbeckiæ. Rue. Papilio asterias on, 1512. worms, 1512. Rumex insects, 1165. Russia. Excessive injury done by a beetle in, 1985. Rust. Grain Aphis vs., 1806. and Hessian fly. Wheat, 1605. of orange, 2308. Hubbard, H. G. Report on, red social wasp, 1329, 1558. Rustic. Corn, 1059. Ruta graveolens. Insects on, 1512. Rye. Beetle working in wheat, oats, and, 1259. gall-gnat, Loew, H. Description of, 2267. RYE. INSECTS AFFECTING. Cecidomyia secalina, 2267. Silvanus surinamensis, 1259. Tenebrio molitor, 72. Rve. Worms in flour and, 72. Sack bearers, 1052. Saddle-back caterpillars, 60, 161, 424, 829, 1092. Sage. Galls growing on wild, 1347. St. Louis Academy of Science. Presidential address, 1564, 1629. Sale of silk-worm eggs, 1908. Salivary glands of Phylloxera, 1687. Salix. Galls of Cecidomyidæ on, 46, 197. nigra. Vanessa antiopa on, 1609. Salt and vinegar for insects, 937. water insects used as food, 2203, Salutatory, 386. Samia cecropia - Attacus cecropia. columbia = Attacus columbia. cynthia = Attacus cynthia. The ailanthus silk-worm, 1718. Food-plants of, 2204. ricini = Attacus ricini. San Joaquin Valley, California. Coquillett, D. W. Report on locusts of, 2363. Sandy soil. The Phylloxera in, 2250. Sannina exitiosa, 55, 587, 617, 871, 1017, 1059, 1070.

1475, 1513, 2238. Saperda, 379, 934, 2267. bivittata = 8. candida. Oviposition of, 1621. calcarata, 71, 963, 1849. candida, 2, 6, 55, 57, 277, 377, 421, 578, 714, 870, 939, 962, 1023, 1038, 1056, 1059, 1213, 1227, 1332, 1367, 1441, 1513, 1608, 1621, 1630, 2230, 2238, 2266. lateralis, 583.

Sapenda. Variation in the two-striped, 1213. on willow, 2267. Sarcophaga, 1390. carnaria, 1390, 1452, 2843. lineata destructive to locusts in Dardanelles, 2075. sarraceniæ n. sp., 1885, 1890, 2843. Sarcophagidæ, 2256. Sarcoptes scabiei, 497. Sarcopsylla penetrans, 412, 497. Sarracenia variolaris. Description and natural history of two insects which brave the dangers of, 1390. Insect-catching habits of, 1385, 1390. Sarracenias, 1499. Satellite Sphinx, 1127. Saturnia io - Hyperchiria io. maia = Hemileuca maia. Sauce. Maggots in, 1607. Savin twigs. Eggs of periodical Cicada in, 698. Saw-flies, 294. Spruce-tree, 115. Saw-fly. Ash, 2291. egga, 108. Elm-tree, 699. Gooseberry, 140. Imported, 228, 333, Grape, 2291. Larch, 2232. Large, 1514. larvæ on the quince, 2243. Plum, 2291. Raspberry, 1212, 1641. Strawberry, 2418. Scab in apple vs. apple-tree plant-lice, 1139. potato, 239. Scabby potatoes, 1444. Scale on acacias. Large white, 1730. Cottony maple, 1816, 2291. on Euonymus latifolia, 2403. insect on maple, 1890. Cottony, 1515. Pine-leaf, 1329. on raspberry, 1968. Scale insects affecting the orange. Experiments on, 2164. Coquillett, D. W., gas treatment for, 2418. Experiments on, 2232. Introduction and spread of, 2232. Koebele, A., experiments against, 2418. on magnolia, 1377. Methods of destroying, 2119. New species of, 1919. of the orange, 2119, 2869. in Florida. Voyle, J., report on the effects of cold, 2277. The use of gases against, 2389.

White-pine, 97.

Apple-tree, 69.

Scarites subterraneus, 805.

Scales, 218.

S. Number of, 2267.

Scarlet mite, 1470. Selandria cerasi = Eriocampa cerasi. Scarred apple trees, 951. ross = Monostegia ross. Scavenger! Is Cyrtoneura a parasite or a, 2102. rubi = Monophadnus rubi. vitis = Blennocampa pygmæa. mistaken for a foe, 416. not a parasite. Phora a, 1923. Self-taught entomologists, 276. Scelio ovivora, 1643. Semasia helianthana n. sp., 1968. Scenopinus from human lungs. Larva of, 1348. prunivora n. sp., 373 = Grapholitha pruвр., 2354. nivora. Schadlicherer Inseckten. Einige unserer, 1325. Semiotellus chalcidiphagus n. sp., 384, 563. Schizoneura americana n. sp., 1678. destructor, 1581. cornicola, 27. Send plenty of specimens, 592. fungicols, 27. Senometopia atropivora, 2343. lanigera, 58, 867, 872, 873, 467, 495, 963, militaris n. sp., 6. 990, 997, 1038, 1059, 1187. Series iricolor, 74. vespertina, 1140. rilevi, 1059. tessellata, 2361. Sericaria mori, 45, 542, 1311, 1346, 1609, 1616, 1648, 1718, 1721, 1724, 1753, 1945, 2062, ulmi, 1959. Sciapteron polistiformis, 373, 1301, 1509. 2234, 2381. robiniæ, 2410. Length and weight of thread of, Sciara, 239, 596, 1547, 1662, 1950. 1250 mali, 211. Sericornia claypoleana n. sp., 2114 = Steganoocellaris, 2119. ptycha claypo'eana. Scientific names, 101, 430, 826. instrutana = Steganoptycha clayponomenciature, 303, 768. aymbols, 405. Serrell automatic silk-real. Walker, P., 2388. Seeia pelasgus = Hemaris thysbe. Scolopendra castanipes, 834. Scolytus carys n. sp., 220, 938 = S. 4-spinosus. Seventeen-year Cicada, 1034, 1093, 1698, 2312, 2314. destructor, 938, 1329. in Iowa, 1787. fagi n. sp., 220. Periodical or, 2312, 2314. Food-habits of, 1940. Sting of the, 407. Hickory, 1754. locust, 370, 884, 1489, 1971, 1979. Imported orchard, 2233. or thirteen-year locust. Periodical Cicada, alias the, 1150. pyri = Xyleborus pyri. Seventh report State entomologist of Missouri, quadrispinosus, 220, 938, 1329, 1401, 1754. rugulosus, 1940, 2233. Synoptic table of, 220. Severe cold on insects. Effects of, 2037. Sex in butterflies. Controlling, 1352. Scorpion in Kansas, 1119. insects. Regulating, 1415. Scorpions, 453. Shad. Terrestrial insects in stomach of, 1863. Scotch and Austrian pines. False caterpillars on Shade trees and their insect defoliators, 2378, Scrape our trees! Shall we, 1399. Screw-worm, 209. Shaffer. Entomology. Prof. Riley to Dr., 2000. in Central America. Prevalence of Shagreened cut-worm, 2291. Shall we scrape our trees, 1399. Shedding of traches and double coc vir. 1215. its parentage in doubt, 1880. Pyrethrum for the, 1921. Sheep bot, 450. Scudderia curvicauda, 1363, 2241. fly, 2238. Scurfy apple-bark louse, 2305. gad-fly, 887. Scymnus cervicalis, 1059. Sheldon's borer remedy, 193. l.æmorrhous, 151. Shell-bark hickory. Citheronia regalis on, 775. Shimer, H., criticised by B. D. Walsh, 174. sp., 151. Scyphophorus yucce, 1602. Shrube, fruit, and grains. Webster, F. M. Experiments on the effect of punture of Hemipters Second report State entomologist of Missouri, 1127. on, 2386. U. S. Entomological Commission, Sialidæ, 2267. 1957. Sialis infumata, 39. Secretion on stems of bitter-sweet. White, waxy, Sigalphus Curculio parasite, 1301. curculionis, 1751, 1796. Seed corn maggot, 657, 1059, 1065. rufus n. sp., 1301 = var. of curculionis.grain. • Chinch-bug not in, 888. Silesia in 1869. Cohn, F. Hessian-fly in, 2267. Silk-culture, 1668, 1753, 2119, 2291, 2363, 2394. peas from bugs. To keep, 434. in California. Promotion of, 2000. ticks under bark of apple-trees, 1138. weevil. Honey locust, 1026. the colonies, 2278. Seeds and galls. Jumping, 1496, 2163, 2173. how to dispose of cocoons, 1881. Segments in head of winged insects. Packard, A. in Kansas, 1542.

Profits of, 2338.

Silk-culture in the United States, 1668, 1945, 2070, 2196, 2836. Walker, P., 2418. fiber from cocoons raised at the Department. McMurtrie, W. Test of, 2253. industry in the United States (Review), 2268. producer. Antherma yama maia as a, 1346. reel. Walker P. Serrell's automatic, 2386. spiders, 830. Silk-worm, 2095. Ailanthus, 899, 1811. breeding, 1648. Cecropia, 1311. Circular, 2303. cocoons, 1486. ogga, 1114, 1758. Japanese mode of packing, 1616. prices and where obtained, 2062. Sale of, 1908, 2303. Length and weight of the thread of the, 1359. Luna, 1311. Manual of the, 1721, 1724. in Missouri. Ailanthus, 1460. Mulberry, 1311. naturalized. Ailanthus, 1179. notes, 2232. Osage orange for the mulberry, 1220, vs. mulberry for the, Perny, 1311. Silk-worm. Polyphemus, 1311. Promethes, 1311. Tusseh, 1311. Yamamai, 1311. Silk-worms fed with osage orange, 1341. Food for, 1318. Nomenclature of American, 303. Silky mite, 1423, 1625. Silpha americana, 817, 1226. peltata = S. americana. surinamensis, 1259, 1260, 1261. Silphurus femoratus, 39. Silvanus. Grain, 1259. in dried English currants, 1260. in flouring-mills, 1261. Silver plate by insects. Damage to, 2154.

Simulium, 1160, 2291, 2416. attacking mules, 1811. feeding on other insects, 2177. from Lake Superior, 2032. meridionale n. sp., 2394. molestum, 1174. pecuarum n. sp., 2394. pictipes, 1958, 2032.

piscidium n. sp., 1174, 1283, 1345. Sines diadema, 34, 638, 731.

Singular caterpillar, 890. Sinoxylon basilare, 1311, 1747.

Red shouldered, 1311.

Siphonophora, 1678.

avenæ = Nectarophora granaria. rudbeckiæ = Nectarophora rudbeckise.

Sitaris, 1600, 1643.

443 Sitones flavescens, 2394. Sitophilus granarius = Calandra granaria. oryzæ = Calandra oryzæ. remotepunctata = Calandra remotepunctata. Six worst insect enemies of fruit-growers in northern Illinois, 377. Sixth report State entomologist of Missouri, 1363. Skeletonizer. Apple-leaf, 1311, 1322. Skipper. Cheese, 1700. Skippers injuring smoked ham, 1734. Slug on pear and cherry-trees, 1222. Pear-tree, 1382. Rose, 672, 1780. worm. Spined, 1150. Small apple-leaf worm, 747. borer in apple-twig, 1458. galls and minings in apple-twigs, 552. A new leaf-hopper injurious to, grains. 1767. and grasses. Webster, F. M. Insects affecting, 2394. reddish snout beetle in apple, 1244. white bristly cut-worm, 1059. Smaller cornstalk borer, 2119. Smart bugs. Concerning certain, 565. Smeared dagger, 1301. Smerinthus. Blind-eyed, 1912. excæcatus, 1912.

Smicra albifrons, 6. Smilax. Agrotis saucia injuring, 1941. injured by cut-worms, 1941.

Smilia auriculata, 1183. Smith, E. A. The cotton belt, 2343.

Smith, J. B. Report upon cranberry and hop in sects, 2277.

Report upon insects affecting the hop and the cranberry, 2291. Smith's patent Curculio-trap, 969.

Smoked hams. Skippers injuring, 1734. Smut in wheat, 1479.

Cause of, 1461. Snake of Brazil. Lignified, 2136. worms, 1547.

Snakes. Hair, 612, 861, 1143. Horse hair, 612.

Snellen von Vollenhoven. Obituary, 1817.

Snout beetle, 700, 1168, 1302. on apple. Small reddish, 1244.

> Imbricated, 1301, 2291. Large gray straight horned, 1033.

beetles injurious to fruits, 1302. Snow-balls. Aphidida on, 1184.

Lice on, 1184.

flean, 608.

Snowy tree-cricket, 1059, 1329, 2238, 2291.

in raspberry canes. Eggs of, 1122

So-called army-worms. Three, 328. web-worm of young trout, 1283.

Soap. Apply, 1367. against borers, 47.

Social wasp. Rust-red, 1329, 1559.

Social wasps, 616. Sod-worm, 2418.

Soft maples. Flat-headed borer in, 1250.

Soil. The Phylloxera in sandy, 2250.

Soils. Effects of Paris green on, 1427

Solanum carolinense. Cassida texana on, 2096.

Solanaceze insects, 1238.

Species. Theory of, 384.

Specific for the Colorado potato beetle, 1490.

names. Capitalizing, 2170, 2257.

value of Apatura aliciæ, 1977.

Specimens lost, 1210. Doryphora juncta on, 2096. elæagnifolium. Cassida texana on, 2096. Send plenty of, 593. Flea-heetle eating, 1782. Speckled cut-worm, 1059, 2291. insects, 1238. Specters. Walking-sticks or, 1395. melongena. Cassida texana on, 2215, 2235. Spectrum bivittatum, 1298. Doryphora juncta on, 2096, femoratum = Diapheromera femorata. 2215, 2235. Speculations of the New England school of nat-Solar physics and locust multiplication and miuralists. Entomological, 44. gration. Swinton, A. H., 2267. Professor Dana and his entomolog-Soldier-beetle larvæ, 1643. ical, 82. Pennsylvania, 1059. Spermophagus robiniæ, 1026, 1474. bng, 1558, 2291. Sphæria morbosa, 930. Bordered, 1059. Sphærophthalma occidentalis, 814, 832, 858. Glassy-winged, 1301. Sphecidæ. Table of genera of, 375. Rapacious, 1059. Sphecius speciosus, 371, 372, 407, 543, 858, 2014. Ring-banded, 1311. Sphenophori that attack corn. Larval habits of, Spined, 456, 1059, 1311, 1329, 1523. 2030. Sphenophorus. Cocklebur, 1301. Solenobia, 38, 943. Solenopsis geminata, 2105. Corn. 1301. xylonii = S. geminata. Grain, 2363, 2394. parvulus, 2394. Solidago. Aphid on, 2205. robustus, 2030, 2119. gall moth, 1059. sculptilis, 337, 1042, 1301. Grapholitha olivaceana bred from, 2285. zero n. sp., 337 = S. sculptilis. leaves. Galls on, 1924. Sphex ichneumonea, 543. nemoralis. Cecidomyia carbonifera on, 1924. Sphida obliquata, 2357. Sphingicampa n. g., 40. Song notes of the periodical Cicada, 2334. Sorghum. Nola sorghiella on, 2119. bicolor, 40, 45. Sorghum web-worm, 2119. Sphingicampa distigma n. sp., 40 = S. bicolor.Sphingid pupa. Probable sound organ in, 2101. Sound organs in sphingid pupse. Probable, 2101. South America. Report on cotton crop and its Sphingidæ, 771, 1089, 1277, 1418, 1419, 1761, 1784, 2101. enemies in, 2343. Sphinx. Abbot, 1127, 1248, 1277. American lepidoptera. Notes on, 1784. Achemon, 1127. Entomological ignorance in the, 390. atropos, 2101. Field for the entomologist in the, 1109. carolina = Protoparce carolina. Fruit-culture in the, 2265. catalpæ = Ceratomia catalpæ. The imported cabbage-worm in the, 1714. hageni = Ceratomia hageni. Muscle-shape bark-louse on apple trees in Larva of Abbot, 1248, 1277. the, 1774. Aquatic, 1951. Notes from the, 1657. Osage orange, 2119. 5-maculata = Protoparce celeus. Pass, Ill. The bag-worm at, 1156. side of trees. Apple-tree borers on, 634. Satellite, 1127. west. Economic investigations in the, 1864. White-lined morning, 630, 1198, 1301, 1643, Southern buffalo-gnat, 2291, 2394, 2418. 2201 Sphyracephala brevicornis, 299. cabbage-butterfly, 1127, 2232, 2238. grass-worm, 1127. Spider egg-nest. Mud-wasp and, 1847. Illinois. Entomological tour in, 372, Egg sacks of some unknown, 1144. Sowing cotton seeds in hot-beds, 1772. Ladder, 1299. Spain. Phylloxera congress in, 1906. and nest, 1869. Span-worms. Gooseberry, 1068, 1570. Poisonous, 2306. Sparrow. Anent the English, 1667. Red, 1146, 1225, 1710. on roses. Red, 1710. as an insect killer. Worthlessness of the, 2413. wort owlet-moth, 1301. Spathius trifasciatus n. sp., 1329. Spiders, 217. Spattered-copper underwing, 1301. Grape leaf-folders eaten by, 468. Spearman. Larva of the ten-striped, 866. Jumping, 2902. Ten-striped, 32. Silk, 830. Species. Geographical range of, 1614. Trees injured by, 2409. Spilochalcis maries, 1112. Old question of, 2201. of Otiorhynchidæ injurious to cultiva-Spilonota oculana = Tmetocera ocellana ted plants, 2117. Spilosoma acræa, 2343. Phytophagic, 45, 50. virginica, 296, 454, 1202, 1307.

State entomologist of Illinois, 327. Spindle-worms, 331. Spined slug-worm, 1150. for Minnesota, 1108. soldier-bug, 456, 1059, 1311, 1329, 1523. New York, 1863. spider, 813. One day's journal of a, 383. Spirobolus marginatus, 166, 963, 1025. Status and future prospects of silk-culture in the Spittle insect, 93. United States, 2336. Spotted lady-bird, 599. Steele, J. P., report on cotton insects, 2343. Pelidnota, 1301. Steganoptycha claypoleana n. sp., 2206. rove-beetle, 1123. Stick-bug, 448. Cecidonyia impatientis Stictonotus isosomatis n. sp., 2119. touch-me-not. on. 852. Still they come, 908. Galls on, 852. Sting? Do locusts, 371. rumpet-leaf. Insects associated with, of the seventeen-year, 407 1385, 1390. Stinging bug, 778. Spraying trees for protection against insects. Im caterpillars, 1748. larvæ, 760, 811, 1329. proved method of, 2211. Spread of Pieris rapæ into Alabama, 1720. Stings of bees, 116. scale insects. Introduction and, 2232. insects, 116, 217. Spring canker-worm, 1423, 1482, 2238. Stiretrus anchorago, 372. Howell, M. A. Experience fimbriatus = S. anchorago. with, 2267. Stizus brevipennis n. sp., 375 = Megastizus brevi-Locust injury next, 1555. pennis. tails, 317. grandis = Sphecius specionus. Sprinklers and atomizers, 1857. speciosus = Sphecius speciosus. Spruce-borer. Flat-headed, 2267. Stock. Insect injurious to live, 2238. bud. Tortrix, 2232. Stomach of black-bass. Larvæ in, 1792. worm. Reddish yellow, 2291. blue-bird. Ichneumon in, 1878. Epizeuxia, 2363. Larvæ from, 1871. cat-bird. Tipula eggs in, 1735, 1745. SPRUCE. INSECTS AFFECTING. Epizeuxis, 2363. lark, robin, and sun-fish. Insect from, Lophyrus abietis, 115. 1926 Melanophila, 2267. meadow-lark. Beetles in, 1013. Nematus integer, 2232. robin. Cut-worms from, 1873. rock-bass. Insects from, 1793. Plume moth, 2363. shad. Terrestrial insects in, 1853. Therina, 2363. Tortrix fumiferana, 2332. Stomoxys calcitrans, 864. Spruce. Nematus, 2232. Stored corn. Serious injury to, 1683. plume-moth, 2363. rice. Insects affecting, 2190. Therina, 2363. Strachia histrionica = Murgantia histrionica. tree saw-flies, 115. Straight-horned snout-beetle. Large gray, 1033 Squares. Butterfly larvæ injurious to cotton, 1872. Strange bug, 1078. Squash-borer, 378, 1127. Strawberry beds. White-grub in, 1236. bug, 409, 867, 1059. borers, 2385. does not touch the white bush scolbugs, 637. lop, 825. crown-borer, 1301. Glorified, 262. Notes on, 1393. SQUASII. INSECTS AFFECTING. destrover, 689. Anasa tristis, 409, 825. Eggs of bugs on, 690. Coreus tristis = Anasa tristis. enemy, 342. Diabrotica vittata, 355. and grape-vines. Injured, 682. STRAWBERRY. INSECTS AFFECTING. Epilachna borealis, 125. Melittia ceto, 125, 248, 378. Ægeria impropria, 2385. Trochilium cucurbita = Melittia ceto. Agrotis tricosa, 2324. Squash-vine borer, 248. Anarsia lineatella, 2235. insects, 125, 355. Anchylopera fragariæ = Phoxopteria fragaгiæ. Squirrel-bot, 526. Anthonomus musculus, 2363 Stag-beetle, 957, 1517. Horns of the, 755. Capsus oblineatus = Lygus pratensis. Stalk-borer, 1558, 1595, 1646, 2291. Chrysomelidæ, 2235. Dahlia, 862, 1009. Colaspis flavida, 1904. and aster, 940. Corimelæna pulicaria, 637. Rice, 2119. Cyclocephala immaculata, 1236. Eccopsis permundana, 2324. Tomato, 694, 976. weevil. Potato, 1558. Emphytus maculatus = Harpiphorus macula-Staphylinide wanted. American, 1786. tus. Staphylinus maculosus, 1123, 1186. Graphops, 2229.

Sunfish. Insects from stomach of, 1926. STRAWBERRY. INSECTS AFFECTING-Continued. Sunspots and insect life, 2004. Harpiphorus maculatus, 499, 955, 964, 1570, 1586, 2324. Supposed army-worm in New York and the cast-Julus sp., 834. ern States, 1990. Lygus pratensis, 682, 2235. bark-lice oggs in Missouri, 1984. cause of yellows in peach trees, 515. Monostogia rosm, 984. Myriapod, 2285. dock. Galls on, 1165. Paria aterrima, 1904, 2229. eggs of the preying Mantis, 1902. Phoxopteris fragarise, 499, 984, 1059, 1574, 2324. hibernating Aletia chrysalids, 1927. Polydesmus serratus, 342. trout enemy, 1141. sp., 834. Swallow-tail. Larva of those, 1237. Saw-fly, 2418. Philenor, 1127. Selandria rosse = Monostegia rosse. Thoas, 1268. Tyloderma fragarize, 1301, 1393. Swallows, 878, 1502. Strawberry leaf-roller, 984, 1059, 1574. Dipterous larva on, 153. A new enemy to, 1904. Swarms of butterflies, 406, 1050. Pests of the, 2324. lady-birds, 824. saw-fly, 2418. minute flies in rooms, 596. weevil, 2363. that devastate the trans-Mississippi worm, 955, 965, 1570. country. Locust, 1674. and remedy, 1586. Sweet-potato beetles, 681, 1510, 1697. worm, 499. SWEET-POTATO. INSECTS AFFECTING. Streaked cottonwood leaf-beetle, 2291. Cassida bivittata 681, 1127, 1510. Striped-beetle, 2291. nigripes, 1127, 1510, 1697. blister-beetle, 1059, 1230, 1558. sp., 94. Cassididre, 993, 1056, 1082, 1127, 1510. bug, 128, 175. cucumber-beetle, 773, 1127, 2238. Coptocycla aurichalces, 681, 1127, 1510. bivittata = Cassida bivittata. bug, 148. flea-beetle, 2291. guttata, 1127, 1510. Sturnella magna. Contents of the stomach of 1013. nigripes = Cassida nigripes. Subangular ground beetle, 1959. Gryllus sp., 2384. Subscribers. To our, 1106. Swellings in apple scions, 548. Subterranean mites. Description of new, 1370. roots of Ampelopsis, 1428. Successful management of the most destructive Swinton, A. H., Solar physics and locust multipliorange insects, 2088. cation and migration, 2267. Successor. Mr. Walsh's, 1105. Switzerland. Extermination and restriction of Sucking organs of bees, wasps, and flies, 2182. Phylloxers in, 2207. Sugar-berry. Leaf-galls and caterpillars, 762. SYCAMORE. INSECTS AFFECTING. Orgyia leucostigma on, 762. Cerambycid, 2267. Lachnus platanicola, 2138. cane in Brazil; Branner, J. C. Insects injurious to, 2277. Symbols. Scientific, 405. SUGAR CANE. INSECT AFFECTING. Synchlora rubivoraria, 721, 1059. Ligyrus rugiceps, 1794. Synergus albipes = S. lana. Sugar-maple. Beetle in, 1014. lana, 41. borer, 2291. lignicola, 41. Eggs on, 350. mendax n. sp., 41. Mite gall on, 1265. rhoditiformis n. sp. 41 = S. lignicole. tree-borer, 547. Synonyms of parasites, 1932. SUGAR-TREE. INSECTS AFFECTING. Synophrus albipes n. sp. = Synergus lana. Arhopalus speciosus = Plagionotus speciosus. Synopsis of North American Heliothinæ (Review), Plagionotus speciosus, 547. Sugaring for moths, 1300. Syntomeida sp., 2412. Natural, 2138. Syringa. Citheronia regalis feeding on, 863. Sulphur cure on peach trees, 176. Syrphids. Larval habits of, 168. in trees, 98, 228, 931, 1154, 1383. Syrphus-fly. Root-louse, 1059. Sumach-beetle. Jumping, 1363. mellinus, 1813. Eggs in, 118. вр., 1251, 1798. SUMACH. INSECTS AFFECTING. caught by flower of Bidens chrysan-Blepharida rhois, 1363. themoides, 1761. Coccid. 320. Systematic position of the Orthoptera. Packard. Orchelimum sp., 118. A. S., 2267. Xiphidium sp., 118. relations of Platypsyllus as deter-Summer. Anticipated locust injury next, 1615. mined by the larva, 2417. dormancy of butterfly larvæ, 1410. Systechus leucophæus, 2355. larva of Phyciodes nycoreas, 1541, 1947, 1959, 2002.

Tabanus atratus, 1127.

teis, 1410.

Tabanus, sp., 40. Test of machinery for destroying the cotton-Table. On our, 395, 408, 506, 546, 568, 623, 827, worm. Barnard, W. S., 2253. 1110. Tests of silk-fiber from cocoons raised at the Department. McMurtrie, W., 2253. Tachina aletise n. sp., 1712, 2843. anonyma n. ep., 1311. Tetranychus americanus, 1326. archippivora n. sp., 1301 = Masicora arirritans, 1326. telarius, 1146, 1225, 1710. chippivora. concinnata, 2343. Means against, 508. flice, 1643, 1736. Tetraopes femoratus, 323. fly, 1558. 5-maculatus, 1140. tetraophthalmus, 323. Anonymous, 1423, 1625. tornator = T. tetraophthalmus. Cecropia, 1311. Red-tailed, 1127. Tetrastichus esurus, 1712, 2843. Yellow-tailed, 1127, 1625. productus n. sp., 2332. Tettigonia coagulata, 1024. fraterna, 2343. phycitæ, 1311. sp., 951. sp., 2378. vitis = Typhlocyba vitis. Tettigonidæ, 452. villica, 2343. Tachinid eggs, 1914. Tettix granulata, 1566. Tachinidæ, 66, 1801. Texas again. Tarantula of, 493. fever. Ticks and, 404. Teniopteryx fasciate, 249. Tamarack. Pieris menapia feeding on, 2121. in 1883. Anderson, E. H. Cotton-worm in south, 2253. Tamariscus. Nanodes tamarisci on, 1496. spring of 1886. Bruner, L. Locuste in, Tanay for borers, 1016. Tarantula killer. Notes on the, 823. 2282 of Texas, 466, 521, 1178. Tarantula of, 466, 521, 1178. again, 493. Thalessa, 2286. Tarnished plant-bug, 1127, 1219, 2291, 2363. atrata, 338, 477. Tarred paper for fruit trees, 1568. lunator, 1111, 1126, 2350. Tat. Tit for, 432. Thecla posas, 1872. Tawney emperor, 1363. Thelaxes ulmicola = Colopha ulmicola. Tea roses. Failure of, 1708. Thelia bimaculata, 787. Teles polyphemus, 244, 542, 629, 635, 841, 1164, 1311. Theory. Plug-ugly, 238. Caterpillar of, 765. wanted. New locust, 1532. Theridula sphærula, 2343. Cocoons of, 2181. Telenomus, 2115. Therina, Pine, 2363. bifidus, 2578, 2394. Spruce, 2363. Telephoridae, 374. Thersilochus conotracheli n. sp., 1301, 1751. Telephorus bilineatus, 1311, 1825. Thick-thighed walking-stick. 1701, 1721. Temnochila virescens = Trogosi ta virescens. Third report State entomologist of Missouri, 1301. Ten-lined potato beetle, 925. U. S. Entomological Commission, striped spearman, 32. Larves of, 866. Thirteen-year locust. The periodical Cicada alias Tenacity of life, 1755. the seventeen-year and the, 1159. Tenebrio molitor, 72, 191, 980, 2167. Thistle. Beetles on, 169. obscurus, 191, 2167. THISTLE. INSECTS AFFECTING. Tenebrionidæ, 2105. Diabrotica longicornis, 169. Tent caterpillar, 642. Phyllobrotica longicornis = Diabrotica longiof the apple tree, 336, 1301, 2238, 2291. Platyptilus carduidactylus, 1050. Eggs of the American, 1329. Thistle-plume, 1059. of the forest, 645, 688, 1181, 1200 Those swallow-tail, 1268. 1301, 1331. Larva of the, 1237. caterpillars, 363. THORN. INSECTS AFFECTING. and fall web-worms, 819. Anthonomus crategi, 376. Tenthredinidæ, 197, 294. Cecidomyid, 376. Descriptions of new, 385. Thorn-leaf gall. Mr. Couper's, 376. Larvæ of, 108, Those centennial insects, 1511. Tenthredo. Venation of, 197. Thousand-legged worm, 193, 236, 261, 834, 1025. Three-banded Spathius, 1329. Teras cinderella, 1311. malivorana, 1311. lined leaf-beetle, 1059, 1558. Willow, 2363. so-called army-worms, 328. Terias, 1711. worms and their work, 1504. Termes, 2267. Thripe, 1127, 2362. flavipes, 1620, 1729. Food-habits of, 53, 280, 685. Terrestrial larvæ in stomach of shad, 1853.

sp., 203.

Thrips. True and bogus, 203. TOBACCO. INSECTS AFFECTING. Crepidodera cucumeria, 1782. of the vine-grower, 53. Thrushes. Food-habits of, 1726. parvula, 1782. Thyreus abbotii, 763, 1018, 1118, 1127, 1248, 1277. Epitrix cucumeris = Crepidodera cucumeria. Thyridopteryx ephemerseformis, 182, 206, 266, 271, hirtipennis = Crepidodera parvula 427, 538, 629, 641, 738, 815, 1036, 1059, 1090, 1156, 1189, Gryllus sp., 2384. 1852, 1424, 1472, 1650, 1658, 2272, 2378, 2379. Macrosila carolina = Protoparce carolina. Thyridopteryx ephemeræformis. Remarks on, 5-maculata = Protoparce celeus. Protoparce carolina, 131, 933, 1262, 2238. celeus, 131, 771, 885, 933, 1329. Thysania zenobia, 1291. Thysanopters, 174. Sphinx carolina = Protoparce carolina. Thysanura, 2267. 5-maculata = Protoparce celeus Tibicen cassinii, 1057, 2367. Tobacco moth, 131. septendecim, 51, 189, 213, 870, 407, 474, 527, plants. Flea beetles on young, 1782. 618, 619, 648, 698, 707, 746, 884, 920, 1034, worm, 885, 2238. 1037, 1057, 1059, 1093, 1159, 1311, 1489, 1609, moth, 1059. 1624, 1698, 1737, 1809, 1836, 1971, 1979, 1994, Counterworking the, 1329. 2014, 2112, 2144, 2216, 2305, 2307, 2312, 2314, Tilden tomato and the, 933. Tolype velleda, 972, 1773. 2315, 2318, 2320, 2321, 2829, 2334, 2363, 2367. tredecim, 474, 746, 1034, 1057,1159,1609,1624, Tomato feeding worm, 850. 1809, 1836, 1979, 1994, 1997, 2014, 2112, 2307, fruit worm, 1136. 2312, 2329, 2367. gall. Grape-vine, 1329. Ticks under bark of apple-trees. Seed, 1133. TOMATO. INSECTS AFFECTING. and Texas fever, 404. Doryphora 10-lineata, 21. Gortyna nitela, 694, 734, 976. Tiger beetle. Common, 1763. larva, 719. Heliothis armigera, 1136. Prodenia commelinæ, 850. Virginia, 1059. beetles, 628, 1643. Protoparce carolina, 933, 1089. moth. Chrysalis of the virgin, 674. celeus, 152, 155, 933, 1089, 1661. Isabella, 1311. Splring 5-maculata = Proteparce celeus. Tomato stalk borer, 694, 976. Tilden tomato and tobacco-worm, 933. Tile-horned Prionus, 1127. and the tobacco worm. Tilden, 933. worm, 152, 1059, 1558, 1661. Tilia gall, 1116. Odontota rubra on, 1849. again, 1104. Timber borers, 191. Worm eating into green, 734. worm. Parasitized, 155. encourages apple-tree borers, 1441. That venomous, 1089. Insect in, 918. worms not poisonous, 771. TIMOTHY. INSECTS AFFECTING. Tomicus pusillus = Pityophthorus minutissimus. Hadena devastatrix, 2394. Sphenophorus parvulus, 2394. ramulorum = Pityophthorus microgra-Timothy: Webster, F. M. Insects affecting, 2394. phus. Tinea, 1633. Tongue. Butterfly's, 1815. Tineid. Apple-tree, 77, 92. Humming-bird moths caught by the, genus allied to Pronuba. A new, 1830. Tineidæ, 70. Moths and butterflies caught by the, New genus of, 1329, 1336. 1761. Tineids in bee-hives, 252. Too fond of honey, 764. Fossil, 2084. Toothed Dermestes, 2363. Torrubia, 1064. Tineina, 1246. Tingis amorphæ n. sp., 45 = Gargaphia amorphæ. elongata, 1430, 1486. ravenelii, 1803, 1823. ciliata = Corythuca ciliata. tiliæ n. sp., 45 = Gargaphia tiliæ. Tortoise beetle. Black-legged, 1127. Tiphia femorata, 1363. Clubbed, 1227, 1558. inornata, 1363, 2301. Golden, 1127. and Rhipiphorus, 2351. Larva of clubbed, 1238. Unadorned, 1363. Mottled, 1127. Tipula, 947, 1161, 1278, 1368. Pale thighed, 1127. eggs in stomach of cat-bird, 1735, 1745. beetles, 105, 767, 1082, 1127. sp., 512. Tortricid. A pretty new gall-making, 2176. trivittata, 512, 1735. Tortricidse, 918, 1975. Tit for tat, 432. Description of new, 1969. Tmetocera ocellana, 747. Oviposition of, 1922, Toads, 217, 262. Tortrix cinderella n. sp., 1311 = Teras cinderella.eat worker bees! Do, 544. fractivittana = Cacocia fractivittana in gardens, 822. fumiferana, 2232. vs. bugs, 1103. rileyana = Cacœcia rileyana.

Tortrix. Spruce bud, 2232. Walnut, 1050. Touch-me-not. Cecidomyia impatientis on spotted. 852. Galls on spotted, 852. Trachese and double cocoons. Shedding of, 1715. Trade in insects, 1986. Tragidion fulvipenne, 490. Tragocephala viridifasciata = Chortophaga viridifasciata. Transformations of insects, 528. Mite, 1618. of red mites, 1632. the tumble bug, 1386. Trap. Thomas Wiers's apple-worm, 1312. Trapping the carpet beetle, 1752. Treat's insect extinguisher (Review), 531. Tree borers of the family Cossidse. Bailey, J. S., 2253. cricket, 251, 953, 961, 999. on grape vines. Eggs of, 723. Grapes cut off by, 414. Habits of, 207. Jumping, 1329. in raspberry canes. Eggs of, 1006. Snowy, 2238. Snowy, 1329, 2291. cut-worms, 229, 281. hopper. Buffalo, 415, 1829. injurious to potatoes. Buffalo, 2107. Trees. Apple-tree borers of the south side of, 634. injured by spiders, 2409. for protection against insects. Improved method of spraying, 2211. Shall we scrape our, 1399. Tremex columba, 458, 928, 1126, 2286. Pigeon, 458, 1126. Trenton, N. J. Bennett, T. Report of experiments at, 2344. Trichius delta, 372. Trichobaris trinotata, 401, 1059, 1558, 2172. Trichodectes ovis, 2048. Trichogramma minutum n. sp., 1301. pretices n. sp., 1712, 2115, 2343. Trichopsenius depressus, 1729. Trichopters, 2267. Trifolium insects, 1459. Trimble's insect enemies of fruit, etc. (Review). Trimerotropis cœruleipes n. sp., 1959. latifasciata n. sp., 1969. similia n. sp., 1950. Triodites, 1947. mus, 1959, 2002. Trioza tripunctata, 669. Triphleps insidiosus, 1853, 2048. Triplax thoracica = Tritoma thoracica. Tritoma thoracica, 114. Tritoxa flexa, 225. Triungulin of Meloidse, 2083. Trochilium acericolum = Egeria acerni. acerni = Ægeria acerni. caudatum = Alcathe caudatum. cucurbitm = Melittia ceto.

denudatum = Fatua denudata.

29 ENT

Trochilium hospes, 197. sp., 1063. tipuliformis = Ægeria tipuliformis. Trogoderma tarsale, 2167. as a museum pest, 2139. Trogosita mauritanica = Tenebrioides mauritanica. virescens, 721. Trogus obsidianator, 1802. Troilus butterfly caterpillar, 469. Trombidium, 624. developed from Astoma, 2071. giganteum n. sp., 1632. holosericeum, 1470. locustarum n. sp., 146, 728, 1287, 1451, 1521, 1598, 1618, 1632. muscarum n. sp., 1632. sericeum, 1451, 1470, 1598, 1618. telarium = Tetranychus telarius. tinctorium, 1470. Trout. Death-web of young, 1138, 1160, 1174. enemy. Supposed, 1141. Food for, 1142. So-called web-worm of young, 1283. True army-worm, 647, 1127. and bogus Yucea moth, 1804. Thrips and bogus Thrips, 203. Trumpet grape-gall, 791, 1116, 1329. leaf. Insects associated with the spotted, 1385, 1390, vine. Clydonopteron tecomæ in seedpods of, 1929. Pyralid infesting seed pods of. 1929. Trupanea apivora = Promachus apivorus. Truths in applied entomology. General, 2291. Trypeta pomonella n. sp., 177, 367, 373, 454, 1320, 1553, 1654, 2119, 2238. solidaginis, 137, 180, 798, 1059. Tryphon atricoxus n. sp., 385. Trypoxylon albitarse, 543. Tulip-tree bark-louse, 271. Tumble-dung. Transformation of the common, 1396. Turf web-worm, 2418. Turk and its crescent. The little, 329. Turkey-gnat, 2894. Turnip enemy, 215. Polydesmus complanatus an enemy to. Tussch silk-worm, 1311. Tussock-moth. Caterpillars of the white-marked, 1227. Eggs of the white-marked, 480. White-marked, 1059, 1363, 2238, 2378. again, 535. Twelve-spotted Diabrotics, 687, 1127, 2418. Twice-stabbed lady-bird, 38, 1329, 1883. Twig-borers, 1052, 1185. girdler, 476, 1938. Twigs amputated by some unknown animal, 442. Berry and cherry, 1503.

Eggs in or on oanes and, 1329.

Twigs girdled by some animal, 443. United States. Status and prospects of silkpruners, 288. culture in the, 2336. Unity of coloration in insects, 50. punctured by periodical Cicada, 1055. Two-striped locust, 1423, 2363. Universal remedies, 204, 503, 818. Saperda. Variation in, 1213. More, 278. sweet-potato beetle, 1127. remedy. Another, 232. walking-stick, 1298. Unjust accusation, 1537. Tyloderma fragariæ, 1301, 1393. Unknown corn pest, 1073. Typhlocyba aurea n. sp., 22. larvæ, 853. binotata n. sp., 22. moth, 753. pallidula n. sp., 22. worm, 986. tricincta, 25. Unmasked. A friend, 374. vitis, 203, 686, 1892, 1853. Unnatural secretion of wax, 782. Typhlocybini. New species of, 22. Unsightly galls on cotton-wood, 446. Untrue. Attractive but, 1644. Typhlodromus pyri, 1739, 1759: Tyroglyphus phylloxeræ n. sp., 1363, 1370. Urena Anomia, 2119. siro, 1703. lobata. Anomis erosa feeding on, 2343. Uji parasite, 1311. Uroceridæ injurious to apple-twigs, 893. Uloma-impressa, 805. Urocerus flavicornis, 2347. Ulmus. Gall-making Pemphiginæ on, 1653. Uropoda americana n. sp., 1505, 1626. vegetans, 1626. Unadorned Tiphia, 1363. Unfledged locusts. Destruction of young or, Useful Labena, 1423. lives. Two, 2370. Habits of young or, 1578. A nuisance made, 983. Unicorn apple-tree caterpillars, 749, Useless! Is any knowledge, 1135. prominent, 1280. Ustilago segetum, 1461, 1479. Unimpregnated eggs hatch! Will, 1029. Utah in 1878. Packard, A.S. Notes of a journey Unique and beautiful noctuid, 2189. to, 1959. United States. Agricultural advancement in the. Utilization of ants in horticulture, 2089, 2137. 1750. Vagabond Crambus, 2119. Clothes moths observed in the. Valedictory, 326. Valery Mayet on Phylloxera, 2207. 2146. Valuable insecticides. Two. 1742. The cotton-worm in the, 1769. Vanessa antiopa, 540, 907, 1234, 1327, 1609, 1704, 171L Department of Agriculture. Di-. comma == Grapta comma. vision of Entomology-Bulletin 1, 2164; 2, 2165; interrogationis - Grapta interrogationis 3, 2253; 4, 2277; 6, 2304; Variable molting in Orgyia, 2379. 8, 2315; 10, 2378; 11, 2344; Variation in Anisopteryx, 1540. 12, 2364; 13, 2382; 14, 2388; the two-striped Saperda, 1213. Variegated cut-worm, 1059, 2291. 15, 2389. Entomological Eccopsis, 2363. Commission. Appropriation for, 1843. Varieties. Phytophagic, 45, 50. Bulletin 1, 1577; 2, 1578; Varying Anomala, 2291. 8, 1736; 6, 2026. Vegetable phenomena, 436. Circular 1, 1575; 2, 1576. Vegetal-feeding ground beetles, 1738. Report 1, 1643; 2, 1959; Vegetation caused by locusts. Changes in, 1495. Harris. Insects injurious to (Re-3, 2267 : 4, 2343, Reports, 1757. view), 568. in Illinois. Insects injurious to, 6, 52. Supplementary instructions Legislation to control insects injurito agents of the, 1888. W.S. Entomologist. Report for 1878, 1721. ous to, 1946. 1881 -'82, 2119. Venation of wings of Anisopteryx, 1540. for 1883, 2282, Venomous tomato-worm. That, 1089. 1884, 2291. Vera Cruz, Mexico. Interesting cotton-worm notes from, 1845. 1885, 2363... Verified. Predictions, 1494. 1886, 2894. 1887, 2418. Vertical insect-boxes, 1963. Vesicants. Meloids as, 912. United States. New source of wealth to, 1668. Vespa crabro, 616. Notes on Aphidina of the, 27, 1678. Psyllids of the, 2210. maculata, 543. settled fact. Hibernation Vespidæ, 190, 505, 770. Victoria. Grape Phylloxera in Geelong, 1732. Aletia xylina in the, 2141. Vilfa vaginæfiora, 402, 1495, 1588. Silk-culture in the, 1668, 1945, 2070, 2196. Vine area of France affected by Phylloxera. Half the, 2020. industry in the (Review), 2266. Blue caterpillars of the, 1127, 1363.

Vine-growers. Thrips of, 53. Water. Moths attracted by falling, 2108. infested with parasites. Hog caterpillar of weevil, 2119. the, 1247. of rice-plant, 1973. Insects injurious to the, 2238. Wavy-striped flea-beetle, 348, 564, 2291. Wax-beans. Epilachna corrupta eating, 2135. loving pomace fly, 2119. insect. New, 2119. Vinegar for insects. Salt and, 937. Virgin tizer-moth. Chrysalis of the, 674. Unnatural secretic n of, 782. Virginia tiger beetle, 1059. worm, 1059. Waxy secretion on stem of bitter-sweet. White, Vitis lituus, 1116. vinifera. Phylloxera destructive to, 1727. Ways of bag-worms, 1424. viticola, 1116. Wealth to the United States. A new source of V-marked Cacocia, 2363. Viviparity of a moth, 2153. Web-worm. Fall, 1301, 1733, 2238, 2378, 2394. Volume. Close of the first, 701. Voyle, J. Experiments on orange scale insects, Garden, 2363. 2164. on hickory. Fall, 460. Report on the effects of cold on the Juniper, 1721. scale insects of the orange in Florida, Sorghum, 2119. Turf, 2418, Wagner, B. Observations on the new crop gallof young trout. So-called, 1283. gnat, 2267. of young trout. Death, 1:38, 1160, 1174. Walker, P. Serrell's automatic silk-reel, 2388. Webster, F. M. Insects affecting barley, 2394. Silk-culture, 2418. buckwheat, 2394. Walking-stick, 144. fall-wheat, 2291, Thick-thighed, 1701, 1721. 2363, 2394, Two-striped, 1298. small grains and sticks. Cicadas and, 920. grasses, 2394. or specters, 1395. timothy, 2394. Walnut case-bearer, 1311. white clover, caterpillars, 1229. 2394 Gregarious, 1045. Record of experiments on the Insect feeding on the sap of black, 1195. effect of punctures of hemip-WALNUT. INSECTS AFFECTING. tera upon shrubs, fruits, and Caccecia rileyana, 1059. grains, 2382. Clisiocampa diastria, 363. Report on Buffalo gnats, 2388. sylvatica = C. disstria. of experiments at La-Datana ministra, 1045, 1229, 2222, 2333. fayette, Ind., 2344. Notodontid, 757. upon the season's obser-Phycis juglandis, 1311. vations in Indiana, Paocus venosus, 1195. especially upon corn Walnut Tortrix, 1059. insects, 2418. tree. Worm on bark of, 757. Weeping lace-wing, 1127. and willow. Enemies of the black, 2333. Weevil, 1673. Walsh, B. D. Entomological collection of, 1107, on apple-trees, New York, 1085. 1203 Clover, 1777. In memoriam, 1098. Honey-locust seed, 1026. Portrait of, 1128. Rice, 1683. Posthumous paper by, 1344. plant water, 1973. Successor to, 1105. Strawberry, 2363. Walshia amorphella, 1127, 2356. Weevils. Aniseed vs. grain, 1742. Wanted. Information, 1111. Raspberries destroyed by, 1788. War on corn-worms, 1522. Well-known animals, Little known facts about, Warbles, 898. 2071. Washington Entomological Society. Address as Wells. White worms in, 1015. president, 2355. West. Canker worms at the, 1539. Wasp. Ichneumon-fly mistaken for, 477. Economic investigations in the South and, and parasites. Mud, 1827. spider-egg nest. Mud, 1847. Genuine army worm in the, 2009. Wasps, 1736. Hop growing in the, 235, 279. Digger, 309. Important observations on the grasshopand their habits, 375, 543, 615. per pest of the, 1571. Social, 616. Indies. Report on the cotton crop and its

enemies in, 2343.

Locusts in the, 2044.

Western cricket, 1959, 2267.

Winged pests of the, 2313.

Sucking organs of, 2182.

Gigantic, 534.

Water bug, 797, 1191.

larva, 1205.

Western cricket, Bruner, L. Observations on the, | Wheat. Leaf-hoppers injuring, 1766. Histology of, 1959. governors on the Rocky Mountain locust, 1557. grasshopper, 1998. Missouri. Ravages of young locusts in, striped cut-worm, 1059. Westward progress of the imported cabbageworm, 1694, 1721. What are army-worms ! 1400. becomes of bumble bees ! 1032. Wheat. Beetles supposed to be feeding on, 1746. Bruchus. A new insect'in, 19. Cause of smut in, 1461. cut-worm, 1059. Cut-worms destroying recently sown, 455. Damage to, 1870. fly. Companion, 2394. head army-worm, 1570, 2418. Hessian fly in seed, 494. insects, 1506. WHEAT. INSECTS AFFECTING. Aphis avenæ = Nectarophora granaria. Asopia farinalis, 980. Blissus leucopterus, 886, 888, 894, 895, 2363. Brachytaraus variegatus, 1461, 1479. Bruchus sp , 19. Calandra remotepunctata, 15. Calocoris rapidus, 2363. Cecidomyia destructor, 494, 1506. tritici = Diplosis tritici. Cecidomyid, 18. Chlorops sp., 2060. Cicadula exitiosa, 1766. Cut-worm, 1059. Diedrocephala flaviceps, 1766, 1767, 2363. Diplosis tritici, 109, 110, 142, 216, 280, 292, 372, 428, 711, 1512, 2238, 2267, 2363. Dræocoris rapidus = Calocoris rapidus. Euschistus fissilis, 2363. Gortyna nitela, 1589, 1870. Isosoma grande, 2288, 2291, 2363, 2394. hordei, 2394. nigrum = I. hordei. tritici, 2060, 2063, 2128, 2363, 2394. Jaseus sexnotatus, 1766. sp., 1766. Leucania albilinea, 1507, 1570. Longitarens ep., 636. Lygus pratensis, 2363. Meromyza americana, 727, 1058, 1506, 1589, 1848, 1875, 2394. Nectarophora granaria, 1806. Noctuid, 455. Oscinus sp., 2394. Pyralis farinalis = Asopia farinalis. Silvanus surinamensis, 1259. Sitophilus remotepunctate = Calandra remotepunctata. Tenebrio molitor, 980. Wire-worm, 2418. Wheat Isosoma, 2063, 2119. just before it ripens, Meromyza ameri-

cana attacking, 1058.

maggots, 727. midge, 109, 110, 142, 216, 280, 292, 428, 711, 1512, 2238, 2291, 2363. A new enemy to, 1507. insect injurious to, 2288. oats, and rye. Beetles working in, 1250. plant. Klippart's (Review), 186. rust and Hessian fly, 1605. Smut in, 1479. stalk-worm on Pacific coast, 2123. stalks. New depredator infesting, 2000, 2063, straw Isosoma, 2291, 2394. Larger, 2291, 2394. Webster, F. M. Insects affecting, 2394. fall, 2301, 2363, 2394. wire-worm, 2418, Worm in joints of, 1848. worms, 980. injuring, 1875. in reference to, 1589. White ant. Inquilines in galleries of, 1729. blast, 2119. clover. Webster, F. M. Insects affecting, 2394. grub, 68, 410, 1020, 1059, 2238, 2363, 2394. fungus, 594, 1064, 1430, 1436, 1399, 1803, 1823. again, 640. information wanted, 1072. parasite, 1363. grube, 5+1. in strawberry-beds, 1236. heart hickory Gelechia, 2363. lined morning Sphinx, 630, 1198, 1301, 1643, marked tussock-moth, 1059, 1363, 2238, 2378. again, 535. caterpillar, 1227. 'eggs, 480, 690. oak. Leaf-miner on, 1879. Woolly gall on, 739. pine plant-louse, 320. scale, 97. trees killed by borers, 1049. weevil, 781, 2363. worm, 465, 956. Abbot's, 1570. scale on acacias. Large, 1730. Means against 2374. waxy secretion on stems of bitter-sweet, 1913. willow insects, 907. worm, 917, 1211. worms in wells, 1015. Why noxious insects increase upon us, 766. Wier, D. B. Native plums, 2388. Wier's apple-worm trap. Thomas, 1312. trap. The coddling-moth, 1334. Wild cherry. Caterpillar's nest on, 242. cherry. Cocoon on, 259. grape-vine. Conical galls on leaves of, 1077. sage. Galls growing on, 1347. Will unimpregnated eggs hatch ! 1029.

Willow. Enemies of the black walnut and, 2333. Worm. In joint of wheat, 1848. gall. Pod-like, 1170. question, 909. galls, 46. and remedy. Strawberry, 1586. of Cecidemyids on, 46, 197. anake, 1662. Gracilaria. Purple, 2363. Tobacco, 885, 2288. WILLOW. INSECTS AFFECTING. Tomato, 152, 1059, 1558, 1661. Cecidomyia salicis-ailiquæ, 1170. An unknown, 986. Cecidomyidæ, 46, 197. Worms. Bee bread devoured by, 1298. Cimbex americana, 1380, 2291, 2333. in cabbage, 1915. isportei = C. americana. Clover, 83. 675, 948, 1132. Clostera americana = Ichthyura inclusa. in cottonwood, 1459. Coleopteron, 197. Currant, 864, 882, 1696. Dipteron, 197. on Dutchman's pipe, 1321. Gracilaria, 2363. feeding on hawthorn, 1051. Hymenopteron, 197. in flour and rye, 72. Ichthyura inclusa, 856. on horse-chestnut. Gregarious, 1192. Nematus ventralis, 907, 917, 1211. How to free wells of, 1015. Saperda, 2267. injuring wheat, 1875. Tenthredinidæ, 197. in joint wheat, 1848. Teras, 2363. under mulch hay, 1161. Vanessa antiopa, 907, 1234, 1609. in osage orange seed, 597. Willow insects. White, 907. in potatoes. Wire, 846. Teras, 2363. in wells, 1015. worm. Large, 1380. Wire, 224, 892, 982, 1030, 2238. White, 917, 1121, and their work. Three, 1504. worms. Gregarious, 856. Worthlessness of the sparrow as an insect killer, Winged pests of the West, 2313. 2413. Phylloxera in California, 1895. Wyckoff's silk-industry in the United States (Re-Wire-worms, 224, 892, 932, 1030, 2238. view), 2268. Means against, 103. Wyoming, Montana, etc., in 1881. Bruner, L. in potatoes, 846. The Rocky Mountain locust in, 2267. Wits jump together. How great, 567. Xanthoptera ridingsii n. sp., 1411 = Exyra riding. W-marked cut-worm, 1059, 2291. Wonder. Back-rolling, 1363. semicrocea, 1385, 1390. Wood-borers. Food habits of, 1902. Xiphidium, 118. Rearing, 1829. Xyleborus cælatus, 2267. gall on white-oak, 732. obesus, 1583. lice on grape-vine roots, 1900. Packard, A. S. Development of, 2267. nymph. Beautiful, 1127, 1363. pyri, 2, 149. Pearl, 1127, 1301, 1363. Xyleutes robinize = Cossus robinize. Woolly elm-tree louse, 1059. Xylina cinerea n. sp., 1301 = Lithophane antenlice on the beech, 449. nata. slug-like worm on apple, 796. Xylocopa carolina = X. virginica. Work in entomology, 202, 418. virginica, 372, 1111. Three worms and their, 1504 Xyloryctes satyrus, 40, 471. Worker bees! Do toads eat, 544. Xylotrechus colonus, 2267. Workers among hymenoptera, 311. Yama-mai silkworm, 1311. Works on North American microlepidoptera, 1975. Year. Destructive insects of the, 2322. World. Insect, 1466, 1467. Entomological notes of the, 2235. Worm. Apple, 1666. Insects of the, 2289, 2331. in apple, 177. Is this a grasshopper, 1565. Army, 11, 17, 120, 670, 876, 906, 1127, 1442, Yeast ferment in insects. Experiments with, 1930. 1482, 1551, 1570, 2119, 2239, 2267, 2269. Yellow-bear. Common, 1301. on bark of walnut tree, 757. fever-fly, 1950. boring into cucumber, 808, 843. headed cut-worm, 1059. peach, 1182. locust, 2363. in wheat-stalk, 1870. tailed Tachina-fly, 1127, 1625. Canker, 86, 172, 1021, 1066, 1127, 1363, 2012. swallow-tail, 361. in corn, 181. Yellows in peach trees. Supposed cause of, 515. Cotton, 1127, 1363, 1649, 1702, 1719, 2077, Yersin. Function of the nervous system of arti-2119, 2130, 2295, culates, 1959. Currant, 877, 1204. Young grasshoppers, 2377. eating in green tomatoes, 734. locusts. Destruction of, 1577. Hundred legged, 219. Ditching for, 1488. infesting meal sacks, 1896.

Habits of, 1578.

Young locusts in western Missouri. Ravages of,

pecan trees girdled, 489. tobacco-plants. Flea-beetle on, 1782. trout. Death-web of, 1138, 1160, 1174. So-called web-worm of, 1283.

Yucca, 1602, 1603.

angustifolis fertilized by Pronuba yuccasella, 1804.

borer, 1420, 1482, 1570.

Notes on the, 1465.

Fertilization of, 1329, 1336, 2171.

filamentosa. Prodoxus decipiens ovipositing in, 2049.

Further notes on the pollination of, 2000.

YUCCA. INSECTS AFFECTING.

Elaphidion tectum, 1602.

Megathymus yuccæ, 1420, 1465, 1482, 1570, 1602.

Prodoxus decipiens, 2000, 2049.

Pronuba yuccasella, 1329, 1336, 1387, 1354, 1363, 160(, 2000, 2171, 2391.

Yucca. Insects affecting—Continued. Sycophorus yucce, 1602.

Yucca moth, 1363.

Oviposition of the, 1354.

Professor Riley and the, 2068.

True and bogus, 1804.

Pollination of, 1603, 1804, 2000, 2371.

Pronuba and its connection with the pollination of, 2391.

Zarsea inflata, 2360.

Zarhipia, 2397.

Further notes on Phengodes and, 2411. Phengodes and, 2408.

Zebra cabbage-worm, 2232.

caterpillar, 179, 1127.

Zeller. Death of Professor, 2179.


Zerene catenaria, 1140.

Zimmermann's flea-beetle, 2291.

Zoo-geographical map of North America. Packard, A, S., 2267.

Zygena minos, 2224.


• • . .



. .

٠ ٠.

. • . -.



This book should be returned to the Library on or before the last date stamped below.

A fine of five cents a day is incurred by retaining it beyond the specified time.

Please return promptly.

